

SYMPLECTIC BIRATIONAL TRANSFORMATIONS OF FINITE ORDER ON O'GRADY'S SIXFOLDS

ANNALISA GROSSI, CLAUDIO ONORATI, AND DAVIDE CESARE VENIANI

ABSTRACT. We prove that any symplectic automorphism of finite order on a manifold of type OG6 acts trivially on the Beauville–Bogomolov–Fujiki lattice and that any birational transformation of finite order acts trivially on its discriminant group. Moreover, we classify all possible invariant and coinvariant sublattices.

1. INTRODUCTION

By a result of Rapagnetta [35], the Beauville–Bogomolov–Fujiki lattice $H^2(X, \mathbb{Z})$ of an irreducible holomorphic symplectic manifold X belonging to the sporadic deformation type of dimension 6 constructed by O'Grady [34] is isomorphic to

$$\mathbf{L} := 3\mathbf{U} \oplus 2[-2].$$

In this paper we answer the question of which isometries on \mathbf{L} can be induced by a symplectic birational transformation of finite order on a manifold of type OG6.

Our first theorem is the classification of all isometries on \mathbf{L} induced by symplectic automorphisms of finite order. Quite surprisingly, this turns out to be trivial.

Theorem 1.1 (§3.2). *Any symplectic automorphism of finite order on a manifold of type OG6 acts trivially on the Beauville–Bogomolov–Fujiki lattice \mathbf{L} .*

This phenomenon has never been observed before in other deformation types of irreducible holomorphic symplectic manifolds. In fact, there exist K3 surfaces, manifolds of type $K3^{[n]}$ and of type $Kum^{[n]}$ which admit symplectic automorphisms of finite order with nontrivial action (see for instance [25, 27, 30]).

Theorem 1.1 can be rephrased by saying that any symplectic automorphism of finite order is contained in the kernel of the representation map $\eta_* : \text{Bir}(X) \rightarrow \text{O}(\mathbf{L})$, where $\eta : H^2(X, \mathbb{Z}) \rightarrow \mathbf{L}$ is a marking (see §2.6). By [12, Theorem 2.1], the kernel is a deformation invariant, and in the case of manifolds of type OG6 it is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^8$ (see [28]).

Our second theorem considers the induced action on the discriminant group $\mathbf{L}^\sharp := \mathbf{L}^\vee/\mathbf{L}$.

Theorem 1.2 (§3.4). *Any symplectic birational transformation of finite order on a manifold of type OG6 acts trivially on the discriminant group \mathbf{L}^\sharp of the lattice \mathbf{L} .*

Date: February 20, 2024.

2020 Mathematics Subject Classification. 14J42 (14J50 14E07).

Key words and phrases. Irreducible holomorphic symplectic manifolds, symplectic birational transformations, symplectic automorphisms.

The second author gratefully acknowledges the Research Council of Norway project 250104 for financial support.

We remark that nonsymplectic automorphisms of manifolds of type OG6 can indeed have nontrivial action on both \mathbf{L} and \mathbf{L}^\sharp , as shown by the first author [10].

Finally, our third theorem classifies all possible isometries on \mathbf{L} induced by symplectic birational transformations of finite order on manifolds of type OG6. We denote the order of an isometry $g \in \mathrm{O}(\mathbf{L})$ by $|g|$ and we use the notation \mathbf{L}^g and \mathbf{L}_g for the invariant and coinvariant sublattices, respectively.

Theorem 1.3 (§3.6). *An isometry $g \in \mathrm{O}(\mathbf{L})$ of finite order is induced by a symplectic birational transformation on a manifold of type OG6 if and only if*

$$(1) \quad |g| \in \{1, 2, 3, 4, 5, 6, 8, 10, 12\}$$

and the pair $(\mathbf{L}^g, \mathbf{L}_g)$ appears in Table 1. In this case, the following equality holds:

$$(2) \quad |\det(\mathbf{L}^g)| = |\det(\mathbf{L}) \cdot \det(\mathbf{L}_g)|.$$

Heuristically, given an action of a group G of birational transformations on an irreducible holomorphic symplectic manifold (not necessarily of type OG6), a coinvariant sublattice of smaller rank corresponds to a larger family of manifolds admitting such an action. More precisely, the number of moduli is given by $\mathrm{rk}(\mathbf{L}^G) - 2$ (see [1, §4]). For this reason, in Table 1 the pairs $(\mathbf{L}^g, \mathbf{L}_g)$ are ordered by increasing $\mathrm{rk}(\mathbf{L}_g)$. The column ‘example’ provides a matrix relative to a standard system of generators of $3\mathbf{U} \oplus 2[-2]$ representing an isometry g with given pair $(\mathbf{L}^g, \mathbf{L}_g)$.

The existence of birational transformations with given (co)invariant sublattice in Theorem 1.3 is ensured by the Torelli theorem for manifolds of type OG6 (Theorem 2.15). It would be very interesting to produce explicit examples of symplectic birational transformations inducing the isometries described in Table 1. Although there are ways to construct nonsymplectic automorphisms geometrically (see for instance [9]), the lack of explicit families of manifolds of type OG6, beside desingularized moduli spaces of sheaves, makes this task difficult. We plan to study a special class of examples in a future work.

1.1. Historical note. For a K3 surface X the groups $\mathrm{Aut}(X)$ and $\mathrm{Bir}(X)$ coincide. All possible finite groups G of symplectic automorphisms were classified by Nikulin [31, 32] in the abelian case and then characterized by Mukai [30] in the general case as certain subgroups of the Mathieu group M_{23} . Alternative proofs were given later by Xiao [39] and Kondō [18]. The invariant and coinvariant sublattices for G of prime order were classified by Morrison [29] and by Garbagnati and Sarti [8]. Their classification was extended to all other groups by Hashimoto [11]. (For a survey, see [16, Chapter 15] or [19]).

Finite groups of symplectic automorphisms on $\mathrm{K3}^{[2]}$ fourfolds were studied first by Camere [5] and Mongardi [22, 24] in the case of prime order, and then by Höhn and Mason [13] in the general case. Finite groups of symplectic automorphisms on manifolds of type $\mathrm{K3}^{[n]}$ with $n > 2$ were treated by Huybrechts [17] and Mongardi [23, 25], who in particular conjectured that all of them appear as certain subgroups of the Conway group Co_1 . Partial results on symplectic automorphisms of $\mathrm{Kum}^{[n]}$ manifolds were obtained by Mongardi, Tari and Wandel [27].

1.2. Contents of the paper. The definitions and the basic results used in this paper are recalled in §2, where we also explain how the geometrical problem is translated into

an arithmetical one thanks to the Torelli theorem for irreducible holomorphic symplectic manifolds. The proofs of the statements above are carried out in §3.

Acknowledgments. We wish to warmly thank Fabio Bernasconi, Samuel Boissière, Simon Brandhorst, Chiara Camere, Alberto Cattaneo, Christian Lehn and Giovanni Mongardi for sharing their insights with us. A special acknowledgment goes to Maxim Smirnov, without whom this paper would probably not exist.

TABLE 1. Isometries $g \in \mathbf{O}(\mathbf{L})$ induced by symplectic birational transformations of finite order on manifolds of type OG6.

$ g $	\mathbf{L}^g	\mathbf{L}_g	example
1	\mathbf{L}	0	id
2	$2\mathbf{U} \oplus [2] \oplus 2[-2]$	$[-2]$	$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
2	$\mathbf{U} \oplus 2[2] \oplus 2[-2]$	$2[-2]$	$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$
2	$3[2] \oplus 2[-2]$	$3[-2]$	$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$
2	$3[2] \oplus [-2]$	\mathbf{D}_4	$\begin{pmatrix} 3 & 4 & 2 & -2 & 2 & -2 & 2 & 0 \\ 4 & 3 & 2 & -2 & 2 & -2 & 2 & 0 \\ -2 & -2 & -1 & 2 & -1 & 1 & -1 & 0 \\ 2 & 2 & 2 & -1 & 1 & -1 & 1 & 0 \\ -2 & -2 & -1 & 1 & -1 & 2 & -1 & 0 \\ 2 & 2 & 1 & -1 & 2 & -1 & 1 & 0 \\ -4 & -4 & -2 & 2 & -2 & 2 & -3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
3	$2\mathbf{U} \oplus [6] \oplus [-2]$	\mathbf{A}_2	$\begin{pmatrix} 1 & 1 & 1 & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
3	$\mathbf{U} \oplus 2[6]$	$2\mathbf{A}_2$	$\begin{pmatrix} 1 & 1 & 1 & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$

Continues on next page

Table 1, follows from previous page

$ g $	\mathbf{L}^g	\mathbf{L}_g	example
4	$\mathbf{U} \oplus [2] \oplus [4] \oplus [-2]$	\mathbf{A}_3	$\begin{pmatrix} -1 & 2 & -1 & -1 & -1 & -1 & 0 & 0 \\ 2 & -1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & -1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & -1 & 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 1 & 1 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
4	$3[2] \oplus [-2]$	\mathbf{D}_4	$\begin{pmatrix} 0 & 1 & -1 & -1 & 0 & 0 & -1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & -1 & 3 & 2 & 1 & -1 & 2 & 0 \\ 1 & -1 & 2 & 1 & 0 & 0 & 1 & 0 \\ -1 & 1 & -2 & -1 & 0 & 1 & -1 & 0 \\ 1 & -1 & 2 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
4	$3[2] \oplus [-4]$	$\mathbf{A}_3 \oplus [-2]$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & -1 & -1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
4	$2[2] \oplus [4]$	\mathbf{D}_5	$\begin{pmatrix} 3 & 4 & 2 & 2 & 2 & 2 & 2 & 4 \\ 4 & 3 & 2 & 2 & 2 & 2 & 2 & 4 \\ 2 & 2 & 2 & 2 & 1 & 2 & 1 & 3 \\ 2 & 2 & 2 & 2 & 2 & 1 & 1 & 3 \\ 2 & 2 & 2 & 1 & 2 & 2 & 1 & 3 \\ 2 & 2 & 1 & 2 & 2 & 2 & 1 & 3 \\ -4 & -4 & -2 & -2 & -2 & -2 & -3 & -4 \\ -8 & -8 & -6 & -6 & -6 & -6 & -4 & -11 \end{pmatrix}$
5	$\mathbf{U} \oplus \begin{pmatrix} 4 & 2 \\ 2 & 6 \end{pmatrix}$	\mathbf{A}_4	$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 2 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & -1 & -1 & 0 & -1 \\ 0 & 0 & -1 & 2 & 2 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & -2 & -2 & 0 & -1 & 0 \\ 0 & 0 & 0 & -2 & -2 & -2 & -2 & -1 \end{pmatrix}$
6	$\mathbf{U} \oplus [2] \oplus [6] \oplus [-2]$	$\mathbf{A}_2 \oplus [-2]$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
6	$3[2] \oplus [-2]$	\mathbf{D}_4	$\begin{pmatrix} 4 & 3 & 2 & -2 & 2 & -2 & 2 & 0 \\ 3 & 3 & 1 & -1 & 2 & -2 & 2 & 0 \\ -2 & -1 & 0 & 1 & -1 & 1 & -1 & 0 \\ 2 & 1 & 1 & 0 & 1 & -1 & 1 & 0 \\ -2 & -2 & -1 & 1 & -1 & 2 & -1 & 0 \\ 2 & 2 & 1 & -1 & 2 & -1 & 1 & 0 \\ -4 & -4 & -2 & 2 & -2 & 2 & -3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
6	$2[2] \oplus [6] \oplus [-2]$	$\mathbf{A}_2 \oplus 2[-2]$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$

Continues on next page

Table 1, follows from previous page

$ g $	\mathbf{L}^g	\mathbf{L}_g	example
6	$2[2] \oplus [6]$	$2\mathbf{A}_2 \oplus [-2]$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & -1 \end{pmatrix}$
8	$2[2] \oplus [4]$	\mathbf{D}_5	$\begin{pmatrix} 2 & 1 & 1 & 1 & 2 & 1 & 1 & 2 \\ 2 & 2 & 1 & 2 & 2 & 2 & 1 & 3 \\ 1 & 1 & 1 & 2 & 2 & 2 & 1 & 2 \\ 2 & 1 & 2 & 2 & 2 & 2 & 1 & 3 \\ 2 & 1 & 1 & 2 & 3 & 3 & 2 & 3 \\ 2 & 1 & 1 & 2 & 3 & 2 & 1 & 3 \\ -2 & 0 & 0 & -2 & -2 & -2 & -1 & -2 \\ -6 & -4 & -4 & -6 & -8 & -6 & -4 & -9 \end{pmatrix}$
10	$[2] \oplus \begin{pmatrix} 4 & 2 \\ 2 & 6 \end{pmatrix}$	$\mathbf{A}_4 \oplus [-2]$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 & -1 \end{pmatrix}$
12	$2[2] \oplus [4]$	\mathbf{D}_5	$\begin{pmatrix} 2 & 2 & 2 & 1 & 2 & 2 & 1 & 3 \\ 2 & 2 & 1 & 2 & 2 & 2 & 1 & 3 \\ 1 & 2 & 2 & 2 & 2 & 2 & 1 & 3 \\ 2 & 1 & 2 & 2 & 2 & 2 & 1 & 3 \\ 2 & 2 & 2 & 2 & 3 & 3 & 1 & 4 \\ 2 & 2 & 2 & 2 & 3 & 4 & 2 & 4 \\ -2 & -2 & -2 & -2 & -2 & -4 & -1 & -4 \\ -6 & -6 & -6 & -6 & -8 & -8 & -4 & -11 \end{pmatrix}$

2. PRELIMINARIES

This section not only explains our conventions and fixes the notation used throughout the paper, but it also reviews some minor results – certainly known to experts – for which we could not find a satisfactory treatment in the literature.

In §2.1 we introduce our conventions on lattices, following Nikulin [33] and Miranda and Morrison [21]. In §2.2 we explain the algorithm, based on Nikulin's work, that we use to find all primitive embeddings of two given lattices. In §2.3 we collect various results related to lattice isometries. In §2.4 we introduce the notion of m -elementary lattice and we explain how to enumerate such lattices. In §2.5 we recall the definition of Hodge structure and other related concepts, as given by Morrison [29] (also cf. [16, §3.1]). Finally, in §2.6 we state the Torelli theorem in the case of manifolds of type OG6, which is the key to characterize groups of automorphisms and birational transformations in an arithmetical way.

2.1. Lattices. In this paper, a *lattice* of rank r is a free finitely generated \mathbb{Z} -module $L \cong \mathbb{Z}^r$ endowed with a nondegenerate symmetric bilinear pairing $L \otimes L \rightarrow \mathbb{Z}$ denoted $e \otimes f \mapsto e \cdot f$. We write $e^2 = e \cdot e$. We say that L is *even* if $e^2 \in 2\mathbb{Z}$ for each $e \in L$. The lattice $L(n)$ is obtained by composing the pairing with multiplication by $n \in \mathbb{Z} \setminus \{0\}$. The

rank, *signature* and *determinant* of L are denoted $\mathrm{rk}(L)$, $\mathrm{sign}(L)$, $\det(L)$, respectively. A lattice is *unimodular* if $|\det(L)| = 1$, and it is *hyperbolic* if $\mathrm{sign}(L) = (1, \mathrm{rk}(L) - 1)$.

The *dual* of L is defined as $L^\vee = \{x \in L \otimes \mathbb{Q} \mid x \cdot y \in \mathbb{Z} \text{ for each } y \in L\}$. The group

$$L^\sharp := L^\vee / L$$

is called the *discriminant group* of L . It is a finite abelian group of order $|L^\sharp| = |\det(L)|$. For even L , the \mathbb{Q} -linear extension of the pairing on L endows L^\sharp with a $\mathbb{Q}/2\mathbb{Z}$ -valued quadratic form. There is a natural homomorphism $\mathrm{O}(L) \rightarrow \mathrm{O}(L^\sharp)$. The image of a subgroup $G \subset \mathrm{O}(L)$ through this homomorphism is denoted G^\sharp . The *divisibility* of a vector $v \in L$ is defined as

$$(v, L) := \gcd\{v \cdot w \mid w \in L\}.$$

Note that $v/(v, L)$ defines an element of L^\vee , hence of L^\sharp .

The unimodular indefinite even lattice of rank 2 is denoted \mathbf{U} . The negative definite ADE lattices are denoted $\mathbf{A}_n, \mathbf{D}_n, \mathbf{E}_n$. The notation $[m]$ with $m \in \mathbb{Z} \setminus \{0\}$ denotes a lattice of rank 1 generated by a vector of square m .

Two even lattices L_1, L_2 are said to belong to the same *genus* if $\mathrm{sign}(L_1) = \mathrm{sign}(L_2)$ and $L_1^\sharp \cong L_2^\sharp$ as finite quadratic forms (cf. [33, Corollary 1.9.4]). All lattices appearing in this paper are unique in their genus. For indefinite lattices this follows from Nikulin's or Miranda and Morrison's criteria ([33, Theorem 3.6.2], [33, Theorem 1.14.2], [21, Corollary 7.8]), and for definite lattices from the Smith–Minkowski–Siegel formula (see [6]).

2.2. Primitive embeddings. An embedding of lattices $M \subset L$ is called *primitive* if L/M is a free abelian group. In this case we write $M \hookrightarrow L$ and we denote by $N := M^\perp$ the *orthogonal complement* of M in L . We refer to [33, Proposition 1.5.1 and Proposition 1.15.1] for more details on what follows.

A primitive embedding of even lattices $M \hookrightarrow L$ is given by a *gluing subgroup* $H \subset M^\sharp$ and a *gluing isometry* $\gamma: H \rightarrow H' \subset N^\sharp$. If Γ denotes the *gluing graph* of γ in $M^\sharp \oplus N(-1)^\sharp$, the following identification between finite quadratic forms holds (the quadratic form on the right hand side is the one induced by $M^\sharp \oplus N(-1)^\sharp$):

$$(3) \quad L^\sharp \cong \Gamma^\perp / \Gamma.$$

Two primitive embeddings $M \hookrightarrow L$ are equivalent under the action of $\mathrm{O}(M)$ and $\mathrm{O}(L)$ if and only if the corresponding groups H and H' are conjugate under the action of $\mathrm{O}(M)$ and $\mathrm{O}(N(-1)) = \mathrm{O}(N)$ in a way that commutes with the gluing isometries.

Assume now that L is unique in its genus. By [33, Proposition 1.15.1], an embedding $M \hookrightarrow L$ is equivalently given by a subgroup $K \subset L^\sharp$, which here we call *embedding subgroup*, and an isometry $\xi: K \rightarrow K' \subset M^\sharp$. If Ξ denotes the graph of ξ in $L^\sharp \oplus M(-1)^\sharp$, the following identification between finite quadratic forms holds (the quadratic form on the right hand side is the one induced by $L^\sharp \oplus M(-1)^\sharp$):

$$(4) \quad N^\sharp \cong \Xi^\perp / \Xi.$$

We define the *gluing* and *embedding index* to be $h := |H|$ and $k := |K|$, respectively. They satisfy

$$(5) \quad h^2 \cdot |\det(L)| = |\det(M) \cdot \det(N)|, \quad k^2 \cdot |\det(N)| = |\det(L) \cdot \det(M)|,$$

from which it follows that $hk = |\det(M)|$.

Given M and L , we proceed in the following way in order to find all primitive embeddings $M \hookrightarrow L$ up to the action of $O(L)$.

- (i) We determine the embedding index k from (5).
- (ii) We find all embedding subgroups $K \subset L^\sharp$ with an isometric counterpart $K' \subset M^\sharp$.
- (iii) For each isometry $\xi: K \rightarrow K'$ we compute $\text{sign}(M^\perp)$ and $(M^\perp)^\sharp$ from (4), determining the possible genera for M^\perp .
- (iv) We find all possible $N = M^\perp$ in these genera.
- (v) For each N , we compute the gluing index h of $M \hookrightarrow L$ with $N \cong M^\perp$.
- (vi) We find all gluing subgroups $H \subset M^\sharp$ with an isometric counterpart $H' \subset N(-1)^\sharp$ up to the action of $O(M)$ and $O(N(-1)) = O(N)$.

Given a vector $v \in M$, we are interested in computing (v, L) . Of course, (v, L) depends on the primitive embedding $M \hookrightarrow L$. In general, it only holds $(v, L) \mid (v, M)$. The following lemma provides a useful formula.

Lemma 2.1. *Let M, L be two lattices. If a primitive embedding $M \hookrightarrow L$ is defined by the gluing subgroup $H \subset M^\sharp$, then for each $v \in M$ it holds*

$$(v, L) = \max\{d \in \mathbb{N} \mid v/d \in H^\perp\}.$$

Proof. Let Γ be the gluing graph. The assertion follows from the identification (3) and the elementary observation that $(v, L) = \max\{d \in \mathbb{N} \mid v/d \in L^\sharp\}$, so

$$(v, L) = \max\{d \in \mathbb{N} \mid (v/d, 0) \in \Gamma^\perp\} = \max\{d \in \mathbb{N} \mid v/d \in H^\perp\}. \quad \square$$

Corollary 2.2. *Let $M \hookrightarrow L$ be a primitive embedding. If $|\det(M^\perp)| = |\det(L) \cdot \det(M)|$, then $(v, L) = 1$ for each $v \in M$.*

Proof. Indeed, by (5) the gluing index is equal to $|\det(M)|$, hence H^\perp is trivial. \square

2.3. Isometries. Given a lattice L and a subgroup $G \subset O(L)$, we denote by L^G the *invariant sublattice*. Its orthogonal complement $L_G := (L^G)^\perp \subset L$ is called the *coinvariant sublattice*. Given $g \in O(L)$, we denote by L^g and L_g the invariant and coinvariant sublattices, respectively, of the subgroup generated by g .

We adopt the convention that the (real) *spinor norm*

$$\text{spin}: O(L) \rightarrow \mathbb{R}^\times / (\mathbb{R}^\times)^2 = \{\pm 1\}$$

takes the value $+1$ on a reflection ρ_v induced by a vector v with $v^2 < 0$. (We refer the reader to [21, Chapter I.10], although there the opposite convention is used.) The kernel of the spinor norm is denoted $O^+(L)$. Equivalently, an isometry $g \in O(L)$ belongs to $O^+(L)$ if and only if g preserves the orientation of a positive definite subspace $V \subset L \otimes \mathbb{R}$ of maximal dimension (cf. [20, Lemma 4.1] and [21, Chapter I, Proposition 11.3]).

Lemma 2.3. *If L_G is negative definite, then $G \subset O^+(L)$.*

Proof. For any $g \in G$ it holds $L_g \subset L_G$, therefore L_g is negative definite. The isometry g , seen as an element of $O(L \otimes \mathbb{R})$, can be written as the product of reflections ρ_v for certain vectors $v \in (L \otimes \mathbb{R})_g$. It follows that $v^2 < 0$ for each such v , so $\text{spin}(g) = +1$. \square

Lemma 2.4. *If $g \in O(L)$ has prime order p , then $p - 1 \mid \text{rk}(L_g)$.*

Proof. Since g is defined over \mathbb{Z} , its eigenspaces in $L \otimes \mathbb{C}$ associated with the primitive p th roots of unity have the same dimension. \square

2.4. m -elementary lattices. Given $m \in \mathbb{N}$, we say that an abelian group A is m -elementary if $m\alpha = 0$ for each $\alpha \in A$. The length of A of an abelian group is denoted by $\ell(A)$, and its p -length by $\ell_p(A)$. We say that a lattice L is m -elementary if L^\sharp is an m -elementary abelian group. An m -elementary lattice L satisfies

$$(6) \quad |\det(L)| \leq m^{\ell(L^\sharp)} \leq m^{\text{rk}(L)}.$$

The following lemma is usually stated for m a prime number, but the same proof works for general m (see for instance [2, §5.3]). The details are left to the reader.

Lemma 2.5. *If Λ is a unimodular lattice and $f \in \text{O}(\Lambda)$ has order m , then Λ^f and Λ_f are m -elementary lattices. \square*

The following statements are a direct consequence of (4) and [4, Lemma 3.1].

Lemma 2.6. *If M is an m -elementary lattice, L is an l -elementary lattice and there exists a primitive embedding $M \hookrightarrow L$, then N is $\text{lcm}(m, l)$ -elementary. \square*

Lemma 2.7. *Let L be an m -elementary even lattice. If m is odd, then $\text{rk}(L)$ is even. \square*

Given $m, d \in \mathbb{N}$ and $s_+, s_- \in \mathbb{Z}_{\geq 0}$ we can enumerate all m -elementary lattices L of signature (s_+, s_-) and determinant $|\det(L)| \leq d$ in the following way.

- (1) We list all possible m -elementary abelian groups A of order $|A| \leq d$.
- (2) We list all possible finite quadratic forms $q: A \rightarrow \mathbb{Q}/2\mathbb{Z}$ using Miranda and Morrison's classification [21].
- (3) For each q , we consider the genus \mathfrak{g} of signature (s_+, s_-) and discriminant group q .
- (4) We list all lattices in \mathfrak{g} .

We carried out Step 3 using the `sageMath` function `genera` found in `sage.quadratic_forms.genera.genus` and implemented by Brandhorst [36]. For all computations in this paper, the last step is made easier by the fact that all lattices considered are unique in their genus.

Now let us consider a primitive embedding $L \hookrightarrow \Lambda$ with Λ a unimodular lattice. We write $R := L^\perp$ and we fix isomorphism $L^\sharp \cong R(-1)^\sharp$ and $\text{O}(L^\sharp) \cong \text{O}(R^\sharp)$. Given $f \in \text{O}(\Lambda)$ with $f(L) = L$, we can consider its restrictions $g = f|_L$ and $h = f|_R$. Then, $g^\sharp = h^\sharp$ through the isomorphism $\text{O}(L^\sharp) \cong \text{O}(R^\sharp)$. Conversely, by [33, Corollary 1.5.2], if there exist $g \in \text{O}(L)$ and $h \in \text{O}(R)$ such that g^\sharp and h^\sharp coincide through the isomorphism $\text{O}(L^\sharp) \cong \text{O}(R^\sharp)$, then there exists $f \in \text{O}(\Lambda)$ with $f(L) = L$, $f|_L = g$ and $f|_R = h$.

Lemma 2.8. *Let L be a lattice. If $g \in \text{O}(L)$ has order m and $g^\sharp = \text{id}$, then L_g is an m -elementary lattice.*

Proof. Let Λ be a unimodular lattice such that there exists a primitive embedding $L \hookrightarrow \Lambda$. (There always exists one with $\text{rk}(\Lambda) \leq 2\text{rk}(L)$, since the direct sum of $\text{rk}(L)$ copies of \mathbf{U} is an overlattice of $L \oplus L(-1)$.) Let $R = L^\perp$. Since $g^\sharp = \text{id}$, we can extend g to an isometry $f \in \text{O}(\Lambda)$ by $h = \text{id} \in \text{O}(R)$. As $R_h = 0$, it holds $L_g \cong \Lambda_f$. Given that $|f| = |g|$, we deduce from Lemma 2.5 that L_g is m -elementary. \square

Lemma 2.9. *Let L be a lattice and $g \in \text{O}(L)$ an isometry. Consider the restriction $g' = g|_{L_g} \in \text{O}(L_g)$. If*

$$|\det(L^g)| = |\det(L) \cdot \det(L_g)|,$$

then $g^\sharp = \text{id}$ and $(g')^\sharp = \text{id}$.

Proof. Indeed, by (5) the equation holds if and only if the gluing subgroup of $L_g \hookrightarrow L$ is the whole group L_g^\sharp . Hence, $(L^g)^\sharp = L^\sharp \oplus L_g^\sharp$. Obviously, the restriction of g to L^g is the identity, so its image in $O((L^g)^\sharp)$ is trivial. Therefore, also g^\sharp and $(g')^\sharp$ must be trivial. \square

The discriminant group L^\sharp of a lattice L is said have *parity* $\delta = 0$ if no element of order 2 in L^\sharp has square $\pm 1/2 \pmod{2\mathbb{Z}}$ (cf. [33, p. 144]). Otherwise, one says that L^\sharp has parity $\delta = 1$.

The following theorem, which we will use in §3.1 to deal with isometries of order 2, is easily derived from Nikulin's classification of 2-elementary lattices. An analogous theorem for groups of odd prime order p has been proven by Brandhorst and Cattaneo [3].

Theorem 2.10. *Let $l_+, l_-, t_+, t_-, a \in \mathbb{Z}_{\geq 0}, \delta \in \{0, 1\}$ such that $l_+ - l_- \equiv 0 \pmod{8}$. There exist a unimodular even lattice Λ of signature (l_+, l_-) and an involution $f \in O(\Lambda)$ with coinvariant lattice Λ_f of signature (t_+, t_-) , and discriminant group Λ_f^\sharp of length a and parity δ if and only if the following conditions hold:*

- (1) $t_+ \leq l_+$ and $t_- \leq l_-$;
- (2) $a \leq \min(t_+ + t_-, l_+ + l_- - t_+ - t_-)$;
- (3) $t_+ + t_- + a \equiv 0 \pmod{2}$;
- (4) if $\delta = 0$, then $t_+ - t_- \equiv 0 \pmod{4}$;
- (5) if $a = 0$, then $\delta = 0$ and $t_+ - t_- \equiv 0 \pmod{8}$;
- (6) if $a = 1$, then $t_+ - t_- \equiv \pm 1 \pmod{8}$;
- (7) if $a = 2$ and $t_+ - t_- \equiv 4 \pmod{8}$, then $\delta = 0$;
- (8) if $\delta = 0$, and $a = t_+ + t_-$ or $a = l_+ + l_- - t_+ - t_-$, then $t_+ - t_- \equiv 0 \pmod{8}$.

Proof. If such an involution $f \in O(\Lambda)$ exists, then it holds $\text{sign}(\Lambda_f) = (l_+ - t_+, l_- - t_-)$ and $(\Lambda^f)^\sharp \cong (\Lambda_f)^\sharp$. Nikulin's [33, Theorem 3.6.2] then implies that the conditions listed hold.

Conversely, if the conditions hold, then by the same theorem there exist 2-elementary even lattices M, N with $\text{sign}(M) = (l_+ - t_+, l_- - t_-)$, $\text{sign}(N) = (t_+, t_-)$ and $M^\sharp \cong N(-1)^\sharp$ of length a and parity δ . Therefore, $M \oplus N$ admits a primitive overlattice Λ . Moreover, id on M and $-\text{id}$ on N glue well to an involution $f \in O(\Lambda)$ with $M \cong \Lambda^f$ and $N \cong \Lambda_f$. \square

2.5. Hodge structures. A *Hodge structure of weight 2* on a lattice L is the choice of three complex subspaces $L^{2,0}, L^{1,1}, L^{0,2}$ of $L \otimes \mathbb{C}$ with $L \otimes \mathbb{C} = L^{2,0} \oplus L^{1,1} \oplus L^{0,2}$, $\overline{L^{2,0}} = L^{0,2}$, $\overline{L^{1,1}} = L^{1,1}$, and such that

$$\begin{aligned} v \cdot \bar{v} &> 0 \text{ for } v \in L^{2,0}, v \neq 0, \\ v \cdot w &= 0 \text{ for } v, w \in L^{2,0}, \\ v \cdot w &= 0 \text{ for } v \in L^{2,0} \oplus L^{0,2}, w \in L^{1,1}. \end{aligned}$$

If $\dim_{\mathbb{C}} L^{2,0} = 1$, the Hodge structure is said to be *of K3 type*. In this paper we will only consider Hodge structures of K3 type, which for the sake of simplicity we will just call Hodge structures.

An isometry of a lattice L with a Hodge structure is a *Hodge isometry* if its linear extension to \mathbb{C} preserves the Hodge structure. A Hodge isometry of L is a *symplectic*

isometry if its linear extension restricts to the identity on $L^{2,0}$. The subgroups of Hodge and symplectic isometries are denoted $O_{\text{hdg}}(L)$ and $O_{\text{sp}}(L)$, respectively.

A *signed Hodge structure* on L is a Hodge structure such that the restriction of the bilinear form to $L^{1,1} \cap (L \otimes \mathbb{R})$ has signature $(1, \dim L^{1,1} - 1)$, together with the choice of one connected component, called *positive cone* and denoted P , of the subset

$$\{v \in L^{1,1} \cap (L \otimes \mathbb{R}) \mid v^2 > 0\}.$$

The real subspace $(L^{2,0} \oplus L^{0,2}) \cap (L \otimes \mathbb{R})$ is always positive definite, so only lattices of signature $(3, \text{rk } L - 3)$ can admit a signed Hodge structure of weight 2.

Given a cone $C \subset L \otimes \mathbb{R}$, we set

$$O(L, C) := \{g \in O(L) \mid g(C) = C\}.$$

Note that $O^+(L) = O(L, P)$ if L is of signature $(3, \text{rk } L - 3)$.

Lemma 2.11. *Let L be a lattice. If $C \subset L \otimes \mathbb{R}$ is a convex cone, and $G \subset O(L, C)$ is a finite subgroup, then*

$$(L \otimes \mathbb{R})^G \cap C \neq \emptyset.$$

Proof. Take $\omega \in C$ and define $\omega_G := \sum_{g \in G} g(\omega)$. Clearly, $\omega_G \in (L \otimes \mathbb{R})^G$, but also $\omega_G \in C$, as $g(\omega) \in C$ for every $g \in G$ and C is a convex cone. \square

Given a lattice L with a signed Hodge structure, we define $O_{\text{sp}}^+(L) := O^+(L) \cap O_{\text{sp}}(L)$.

Lemma 2.12. *Let L be a lattice of signature $(3, r - 3)$, with $r \geq 3$, and $G \subset O(L)$ a finite subgroup. If there exists a signed Hodge structure such that $G \subset O_{\text{sp}}^+(L)$, then L_G is negative definite.*

Proof. The invariant real subspace $(L \otimes \mathbb{R})^G$ contains the 2-dimensional real subspace $(L^{2,0} \oplus L^{0,2}) \cap (L \otimes \mathbb{R})$, which is positive definite. Moreover, $(L \otimes \mathbb{R})^G$ intersects the convex cone $P \subset L^{1,1} \cap (L \otimes \mathbb{R})$ by Lemma 2.11. Therefore, $\text{sign } L^G = (3, \text{rk } L^G - 3)$, and L_G is consequently negative definite. \square

Lemma 2.13. *Let L be a lattice of signature $(3, r - 3)$, with $r \geq 3$, and $G \subset O(L)$ a finite subgroup. If L_G is negative definite, then there exists a signed Hodge structure on L such that $L^{1,1} \cap L = L_G$ and $G \subset O_{\text{sp}}^+(L)$.*

Proof. Since L_G is negative definite, $L^G \otimes \mathbb{R}$ contains a positive definite subspace of maximal dimension 3. This implies that

$$\dim_{\mathbb{C}}(L \otimes \mathbb{C})^G = \dim_{\mathbb{R}}(L \otimes \mathbb{R})^G \geq \dim_{\mathbb{R}}(L^G \otimes \mathbb{R}) \geq 3.$$

Therefore, we can find $\sigma \in (L \otimes \mathbb{C})^G$ with $\sigma^2 = 0$, $\sigma \cdot \bar{\sigma} > 0$. Choosing σ outside a certain countable union of hyperplanes, we can suppose that

$$\sigma^\perp \cap L = L_G.$$

In other words, putting $L^{2,0} = \mathbb{C}\sigma$, $L^{0,2} = \overline{L^{2,0}}$ and $L^{1,1} = (L^{2,0} \oplus L^{0,2})^\perp \subset L \otimes \mathbb{C}$ we define a signed Hodge structure on L (choosing P arbitrarily) such that $L^{1,1} \cap L = L_G$. In this way, $G \subset O_{\text{sp}}(L)$ by construction. By Lemma 2.3, it also holds $G \subset O^+(L)$, therefore $G \subset O_{\text{sp}}^+(L)$. \square

Fix a signed Hodge structure on L . Given a subset $\mathcal{W} \subset L$ of primitive vectors which is invariant under $O(L)$, a connected component of

$$(7) \quad \mathbb{P} \setminus \bigcup_{w \in \mathcal{W} \cap L^{1,1}} w^\perp$$

is called a *chamber defined by \mathcal{W}* . Note that every chamber is a convex cone and that $O_{\text{hdg}}^+(L)$ acts on the set of chambers. We put $O_{\text{sp}}^+(L, C) := O_{\text{sp}}^+(L) \cap O(L, C)$.

Lemma 2.14. *Let L be a lattice with a signed Hodge structure. If C is a chamber defined by a subset $\mathcal{W} \subset L$ and $G \subset O_{\text{sp}}^+(L, C)$ is a finite subgroup, then*

$$L_G \cap \mathcal{W} = \emptyset.$$

Proof. By Lemma 2.11, there exists $v \in (L \otimes \mathbb{R})^G \cap C$. Each $w \in L_G$ is orthogonal to v , hence $w \notin \mathcal{W}$. \square

2.6. Torelli theorem. An *irreducible holomorphic symplectic* manifold is a simply connected compact Kähler manifold X such that $H^0(X, \Omega_X^2)$ is generated by a nowhere degenerate holomorphic 2-form σ . An automorphism or a birational transformation of X is called *symplectic* if its pullback acts trivially on σ . The groups of automorphisms and birational transformations of X are denoted $\text{Aut}(X)$ and $\text{Bir}(X)$, respectively.

Let X be an irreducible holomorphic symplectic manifolds of type OG6. A *marking* is the choice of an isometry $\eta: H^2(X, \mathbb{Z}) \rightarrow \mathbf{L} := 3\mathbf{U} \oplus 2[-2]$. Given a *marked pair* (X, η) of type OG6, we consider the associated *representation map*

$$\eta_*: \text{Bir}(X) \rightarrow O(\mathbf{L}), \quad f \mapsto \eta \circ (f^{-1})^* \circ \eta^{-1}.$$

We say that an isometry $g \in O(\mathbf{L})$ is *induced* by an automorphism (resp. birational transformation) if there exists a marked pair (X, η) of type OG6 and an element $f \in \text{Aut}(X)$ (resp. $f \in \text{Bir}(X)$) such that $\eta_*(f) = g$.

We define

$$\begin{aligned} \mathcal{W}_{\text{OG6}}^{\text{pex}} &:= \{w \in \mathbf{L} \mid w^2 = -2, (w, \mathbf{L}) = 2\} \cup \{w \in \mathbf{L} \mid w^2 = -4, (w, \mathbf{L}) = 2\} \\ \mathcal{W}_{\text{OG6}} &:= \mathcal{W}_{\text{OG6}}^{\text{pex}} \cup \{w \in \mathbf{L} \mid w^2 = -2, (w, \mathbf{L}) = 1\}. \end{aligned}$$

Given a signed Hodge structure on \mathbf{L} , a chamber defined by $\mathcal{W}_{\text{OG6}}^{\text{pex}}$ is called an *exceptional chamber*, whereas a chamber defined by \mathcal{W}_{OG6} is called a *Kähler chamber*.

The Torelli theorem for irreducible holomorphic symplectic manifolds was first proven by Verbitsky [37, 38] (cf. also Huybrechts [14, 15] and Markman [20]). We state here a version which is convenient for our purposes.

Theorem 2.15 (Torelli theorem for manifolds of type OG6). *A subgroup $G \subset O(\mathbf{L})$ is induced by a group of symplectic automorphisms (resp. symplectic birational transformations) on a manifold of type OG6 if and only if there exist a signed Hodge structure on \mathbf{L} and a Kähler (resp. exceptional) chamber C such that*

$$G \subset O_{\text{sp}}^+(\mathbf{L}, C). \quad \square$$

Proof. Suppose that G is induced by a group of symplectic automorphisms on a manifold X of type OG6. It follows that G is contained in the monodromy group of X , which is $O^+(\mathbf{L})$ by a result of Mongardi and Rapagnetta [26, Theorem 5.5]. The Hodge decomposition and the canonical choice of the positive cone induce a signed Hodge structure on \mathbf{L} . All

automorphisms of X preserve the Kähler cone, which is a chamber C cut out by vectors in \mathcal{W}_{OG6} by [26, Theorem 1.2], i.e. a Kähler chamber. Therefore, $G \subset \text{O}_{\text{sp}}^+(\mathbf{L}, C)$.

Conversely, suppose there exist a signed Hodge structure on \mathbf{L} and a Kähler chamber C such that $G \subset \text{O}_{\text{sp}}^+(\mathbf{L}, C)$. Then, by Huybrechts' theorem on the surjectivity of the period map [14, Theorem 8.1] there exists a manifold X of type OG6 whose Hodge decomposition induces the given signed Hodge structure. Moreover, we can assume that C is the Kähler cone of X . As G consists of monodromy transformations and respects the Kähler cone, it is induced by automorphisms of X by Markman's Hodge theoretic version of the Torelli theorem [20, Theorem 1.3]. Obviously, the automorphisms must be symplectic.

The same proof applies for birational transformations considering the fundamental exceptional chamber of X ([20, Definition 5.2]), which is the distinguished exceptional chamber containing a Kähler class, instead of the Kähler cone, by [20, Corollary 5.7]. \square

Theorem 2.16. *A finite subgroup $G \subset \text{O}(\mathbf{L})$ is induced by a group of symplectic automorphisms on a manifold of type OG6 if and only if \mathbf{L}_G is negative definite and*

$$(8) \quad \mathbf{L}_G \cap \mathcal{W}_{\text{OG6}} = \emptyset.$$

Proof. The two conditions are necessary by Lemma 2.12 and Lemma 2.14.

Conversely, assume that they hold. By Lemma 2.13 there exists a signed Hodge structure on \mathbf{L} such that $\mathbf{L}^{1,1} \cap \mathbf{L} = \mathbf{L}_G$ and $G \subset \text{O}_{\text{sp}}^+(\mathbf{L})$. Since $\mathbf{L}^{1,1} \cap \mathcal{W}_{\text{OG6}} = \mathbf{L}_G \cap \mathcal{W}_{\text{OG6}} = \emptyset$, the decomposition (7) is trivial: there is only one Kähler chamber, namely \mathbf{P} itself. Therefore, $\text{O}_{\text{sp}}^+(\mathbf{L}) = \text{O}_{\text{sp}}^+(\mathbf{L}, \mathbf{P})$ and we conclude by Theorem 2.15. \square

Theorem 2.17. *A finite subgroup $G \subset \text{O}(\mathbf{L})$ is induced by a group of symplectic birational transformations on a manifold of type OG6 if and only if \mathbf{L}_G is negative definite and*

$$(9) \quad \mathbf{L}_G \cap \mathcal{W}_{\text{OG6}}^{\text{pex}} = \emptyset.$$

Proof. An analogous argument as in Theorem 2.16 applies: there exists a signed Hodge structure on \mathbf{L} such that \mathbf{P} itself is the only exceptional chamber and $G \subset \text{O}_{\text{sp}}^+(\mathbf{L}) = \text{O}_{\text{sp}}^+(\mathbf{L}, \mathbf{P})$. \square

3. PROOFS

Recall that $\mathbf{L} := 3\mathbf{U} \oplus 2[-2]$. We set $\mathbf{A} := 5\mathbf{U}$, $\mathbf{R} := 2[2]$ and we also fix a primitive embedding $\mathbf{L} \hookrightarrow \mathbf{A}$, necessarily with $\mathbf{L}^\perp \cong \mathbf{R}$. From now on, $g \in \text{O}(\mathbf{L})$ will denote an isometry of order m .

We first consider the case in which m is a prime number in §3.1. This is enough to prove Theorem 1.1 in §3.2. We then go on to classify the cases in which m is a power of 2 in §3.3, which we need for the proof of Theorem 1.2 in §3.4. Finally, we treat the remaining cases in §3.5, and we finish off the proof of Theorem 1.3 in §3.6.

3.1. Prime order. We start with two preliminary observations.

Lemma 3.1. *If $g \in \text{O}(\mathbf{L})$ has prime order p and \mathbf{L}_g is negative definite, then $p \in \{2, 3, 5\}$.*

Proof. Since $\text{sign}(\mathbf{L}) = (3, 5)$, we have $\text{rk}(\mathbf{L}_g) \leq 5$, so the statement follows from Lemma 2.4. \square

Lemma 3.2. *The group $\text{O}(\mathbf{L}^\sharp)$ has order 2.*

Proof. Indeed, $O(\mathbf{L}^\sharp)$ is generated by the involution exchanging the two generators of \mathbf{L}^\sharp whose square is equal to $3/2 \pmod{2\mathbb{Z}}$. \square

For a lattice N we introduce the set

$$\mathcal{C}(N) := \{v \in N \mid v^2 = -2 \text{ or } v^2 = -4\}.$$

Note that if N is negative definite, the set $\mathcal{C}(N)$ is finite and can be explicitly determined, for instance using the `sageMath` function `short_vector_list_up_to_length` found in `sage.quadratic_forms.quadratic_form` [36]. The following technical lemma turns out to be of the utmost importance in dealing with condition (9) of [Theorem 2.17](#).

Lemma 3.3. *Let N be a lattice such that for every $\alpha \in N^\sharp$ with*

$$\alpha^2 \in \{3/2, 1\} \pmod{2\mathbb{Z}}$$

there exists $v \in \mathcal{C}(N)$ with $2 \mid (v, N)$ such that the class of $v/2$ in N^\sharp is α . Then, for every primitive embedding $N \hookrightarrow \mathbf{L}$ it holds

$$N \cap \mathcal{W}_{\text{OG6}}^{\text{pex}} = \emptyset \quad \text{if and only if} \quad |\det(N^\perp)| = |\det(\mathbf{L}) \cdot \det(N)|.$$

Proof. Let $H \subset N^\sharp$ be the gluing subgroup of a primitive embedding $N \hookrightarrow \mathbf{L}$ and $M = N^\perp$. The condition $|\det(M)| = |\det(\mathbf{L}) \cdot \det(N)|$ is equivalent by (5) to $H = N^\sharp$, which in turn is equivalent to $H^\perp = 0$.

If this holds, each vector $v \in N$ satisfies $(v, \mathbf{L}) = 1$ by [Corollary 2.2](#), so $N \cap \mathcal{W}_{\text{OG6}}^{\text{pex}} = \emptyset$.

Conversely, take a nonzero element $\alpha \in H^\perp$. Then, thanks to (3), we can identify the element $(0, \alpha) \in M^\sharp \oplus N^\sharp$ with a nonzero element $\beta \in \mathbf{L}^\sharp$. Therefore,

$$\alpha^2 = (0, \alpha)^2 = \beta^2 \in \{3/2, 1\} \pmod{2\mathbb{Z}},$$

so by hypothesis there exists $v \in \mathcal{C}(N)$ such that $v/2 = \alpha \in N^\sharp$. Then, $(v, \mathbf{L}) = 2$ by [Lemma 2.1](#), hence $v \in N \cap \mathcal{W}_{\text{OG6}}^{\text{pex}}$. \square

Remark 3.4. Notably for our purposes, the lattices $[-2]$, $[-4]$, \mathbf{A}_3 , \mathbf{D}_4 and \mathbf{D}_5 satisfy the hypothesis of [Lemma 3.3](#), as one can readily check by inspection.

Proposition 3.5. *If an isometry $g \in O(\mathbf{L})$ of order 2 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to either $[-2]$, $2[-2]$, $3[-2]$ or \mathbf{D}_4 , and condition (2) holds.*

Proof. Let $N = \mathbf{L}_g$ and suppose first that $g^\sharp = \text{id}$.

By [Lemma 2.8](#), N is an even 2-elementary negative definite lattice of rank ≤ 5 . Applying [33, Theorem 3.6.2], we see that N is isometric to one of the following lattices:

$$[-2], 2[-2], 3[-2], \mathbf{D}_4, 4[-2], \mathbf{D}_4 \oplus [-2], 5[-2].$$

By inspection, we see that all these lattices satisfy the hypothesis of [Lemma 3.3](#). Therefore, condition (9) of [Theorem 2.17](#) implies condition (2). This excludes the cases in which $N \cong 4[-2], \mathbf{D}_4 \oplus [-2], 5[-2]$, since they do not admit such a primitive embedding.

Now we turn to the case $g^\sharp \neq \text{id}$.

Consider the embedding $\mathbf{L} \hookrightarrow \mathbf{A}$. We can extend the involution g to an isometry $f \in O(\mathbf{A})$, also with $|f| = 2$, by the involution $h \in O(\mathbf{R})$ exchanging the generators on \mathbf{R} .

Since $\mathbf{\Lambda}_f$ is an even 2-elementary hyperbolic lattice of rank ≤ 6 , [Theorem 2.10](#) yields the following possibilities for its isometry class:

$$\begin{aligned} & [2], \mathbf{U}, \mathbf{U}(2), [2] \oplus [-2], \mathbf{U} \oplus [-2], \mathbf{U}(2) \oplus [-2], \mathbf{U} \oplus 2[-2], \mathbf{U}(2) \oplus 2[-2], \\ & \mathbf{U} \oplus 3[-2], \mathbf{U}(2) \oplus 3[-2], \mathbf{U} \oplus \mathbf{D}_4, \mathbf{U} \oplus 4[-2]. \end{aligned}$$

We then compute the possible embeddings $\mathbf{R}_h \cong [4] \hookrightarrow \mathbf{\Lambda}_f$ to find $N \cong (\mathbf{R}_h)^\perp$. The result of this computation is that N is isometric to one of the following lattices:

$$\begin{aligned} & [-4], [-2] \oplus [-4], \mathbf{A}_3, 2[-2] \oplus [-4], \mathbf{A}_3 \oplus [-2], 3[-2] \oplus [-4], \\ & \mathbf{D}_5, \mathbf{D}_4 \oplus [-4], \mathbf{A}_3 \oplus 2[-2], 4[-2] \oplus [-4]. \end{aligned}$$

By inspection, we see again that all these lattices satisfy the hypothesis of [Lemma 3.3](#). Therefore, condition (9) of [Theorem 2.17](#) implies condition (2), which in turn implies implies that $g^\sharp = \text{id}$ by [Lemma 2.9](#), contradicting the assumption. \square

In the case of odd prime order $p = |g| > 2$, we have $g^\sharp = \text{id}$. Hence, we can use the identity on \mathbf{R} to extend g to an isometry $f \in \mathbf{O}(\mathbf{\Lambda})$. By [\[3, Theorem 1.1\]](#), we can classify the possible sublattices $\mathbf{L}_g \cong \mathbf{\Lambda}_f$, which are p -elementary (see also [\[7, Chapter 15, Theorem 13\]](#)). Since $\det(\mathbf{L}_g)$ is odd, the embedding subgroup of $\mathbf{L}_g \hookrightarrow \mathbf{L}$ is trivial. Thus, condition (2) holds automatically and we obtain the following results.

Proposition 3.6. *If an isometry $g \in \mathbf{O}(\mathbf{L})$ of order 3 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to either \mathbf{A}_2 or $2\mathbf{A}_2$, and condition (2) holds. \square*

Proposition 3.7. *If an isometry $g \in \mathbf{O}(\mathbf{L})$ of order 5 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to \mathbf{A}_4 , and condition (2) holds. \square*

3.2. Proof of [Theorem 1.1](#). Let us assume that a nontrivial isometry $g \in \mathbf{O}(\mathbf{L})$ of finite order is induced by a symplectic automorphism. Up to passing to a power of g , we can suppose without loss of generality that $p = |g|$ is a prime number.

By [Theorem 2.16](#) the coinvariant sublattice \mathbf{L}_g is negative definite, hence [Lemma 3.1](#) applies. A group of automorphisms is in particular a group of birational transformations. Hence, by [Proposition 3.5](#), [3.6](#) and [3.7](#), we know that \mathbf{L}_g must contain a vector w of square -2 .

Since $(w, \mathbf{L}) \mid w^2$, either $(w, \mathbf{L}) = 1$ or $(w, \mathbf{L}) = 2$. In both cases, $w \in \mathcal{W}_{\text{OG6}}$. Therefore, $\mathbf{L}_g \cap \mathcal{W}_{\text{OG6}} \neq \emptyset$, which contradicts the second condition of [Theorem 2.16](#). \square

3.3. Powers of 2. We now examine the cases in which $|g| = 2^k$ for $k > 1$.

Lemma 3.8. *If $g \in \mathbf{O}(\mathbf{L})$ has order m with $4 \mid m$, then \mathbf{L}_g is an m -elementary lattice.*

Proof. If $g^\sharp = \text{id}$, [Lemma 2.8](#) applies immediately. If $g^\sharp \neq \text{id}$, we can argue as follows.

We consider again the embedding $\mathbf{L} \hookrightarrow \mathbf{\Lambda}$ and the extension of g to an isometry $f \in \mathbf{O}(\mathbf{\Lambda})$ by the involution $h \in \mathbf{O}(\mathbf{R})$ exchanging the generators of \mathbf{R} . By [Lemma 2.5](#) the lattice $\mathbf{\Lambda}_f$ is m -elementary. Hence, the claim holds by [Lemma 2.6](#), because $\mathbf{R}_h \cong [4]$ and $\mathbf{L}_g \cong \mathbf{R}_h^\perp \subset \mathbf{\Lambda}_f$. \square

Proposition 3.9. *If an isometry $g \in \mathbf{O}(\mathbf{L})$ of order 4 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to either $\mathbf{A}_3, \mathbf{D}_4, \mathbf{A}_3 \oplus [-2]$ or \mathbf{D}_5 , and condition (2) holds.*

Proof. The lattice $N := \mathbf{L}_g$ is 4-elementary by [Lemma 3.8](#), and negative definite by [Theorem 2.17](#). We can list all (nonzero) 4-elementary negative definite lattices satisfying $\text{rk}(N) \leq 5$ and $|\det(N)| \leq 4^5 = 1024$ by [\(6\)](#). Since N admits a primitive embedding $N \hookrightarrow \mathbf{L}$, a computation shows that N must be isomorphic to one of the following lattices:

$$\begin{aligned} &[-2], [-4], 2[-2], [-2] \oplus [-4], 2[-4], \mathbf{A}_3, 3[-2], 2[-2] \oplus [-4], [-2] \oplus 2[-4], 3[-4], \mathbf{D}_4, \\ &\mathbf{A}_3 \oplus [-2], \mathbf{A}_3 \oplus [-4], 4[-2], 3[-2] \oplus [-4], 2[-2] \oplus 2[-4], [-2] \oplus 3[-4], \mathbf{D}_5, \mathbf{D}_4 \oplus [-2], \\ &\mathbf{D}_4 \oplus [-4], \mathbf{A}_3 \oplus 2[-2], \mathbf{A}_3 \oplus [-2] \oplus [-4], 5[-2], 3[-2] \oplus 2[-4], 2[-2] \oplus 3[-4]. \end{aligned}$$

For all these lattices, the hypothesis of [Lemma 3.3](#) is satisfied. Therefore, condition [\(9\)](#) forces condition [\(2\)](#). The only lattices admitting such a primitive embedding are the following ones:

$$\begin{aligned} &[-2], [-4], 2[-2], [-2] \oplus [-4], 2[-4], \mathbf{A}_3, 3[-2], 2[-2] \oplus [-4], [-2] \oplus 2[-4], 3[-4], \\ &\mathbf{D}_4, \mathbf{A}_3 \oplus [-2], \mathbf{D}_5. \end{aligned}$$

Finally, we observe that the restriction $g' = g|_N \in \text{O}(N)$ satisfies $|g'| = 4$, $N^{g'} = 0$ and $(g')^\sharp = \text{id}$ by [Lemma 2.9](#). A direct computation shows that only the lattices listed in the statement admit such an isometry $g' \in \text{O}(N)$. \square

Corollary 3.10. *If an isometry $g \in \text{O}(\mathbf{L})$ of order divisible by 4 is induced by a symplectic birational transformation, then there exists a primitive embedding $\mathbf{A}_3 \hookrightarrow \mathbf{L}_g$.*

Proof. Let g' be a power of g with $|g'| = 4$. By inspection of all cases in [Proposition 3.9](#), we see that $\mathbf{A}_3 \hookrightarrow \mathbf{L}_{g'}$. Therefore, \mathbf{L}_g contains a sublattice isomorphic to \mathbf{A}_3 . Such an embedding is necessarily primitive, as \mathbf{A}_3 does not admit even overlattices of index > 1 , so we conclude. \square

Lemma 3.11. *A lattice N admitting primitive embeddings $\mathbf{A}_n \hookrightarrow N \hookrightarrow \mathbf{L}$ satisfies*

$$\ell_p(N^\sharp) \leq \begin{cases} \text{rk}(N) - n & \text{if } p \nmid n + 1 \\ \text{rk}(N) - n + 1 & \text{if } p \mid n + 1. \end{cases}$$

Moreover, if $p \neq 2$, then

$$\ell_p(N^\sharp) \leq 8 - \text{rk}(N).$$

Proof. Let $B := \mathbf{A}_n^\perp \subset N$. From the identification [\(3\)](#) for the embedding $\mathbf{A}_n \hookrightarrow N$ and the fact that $\mathbf{A}_n^\sharp \cong \mathbb{Z}/(n+1)\mathbb{Z}$ we see that

$$\ell_p(N^\sharp) \leq \ell_p(\mathbf{A}_n^\sharp) + \ell_p(B^\sharp) \leq \ell_p(\mathbf{A}_n^\sharp) + \text{rk}(B) \leq \begin{cases} 0 + (\text{rk}(N) - n) & \text{if } p \nmid n + 1 \\ 1 + (\text{rk}(N) - n) & \text{if } p \mid n + 1. \end{cases}$$

Suppose now that $p \neq 2$ and let $M = N^\perp \subset \mathbf{L}$. From the identification [\(4\)](#) for the embedding $N \hookrightarrow \mathbf{L}$ we obtain

$$\ell_p(N^\sharp) \leq \ell_p(\mathbf{L}^\sharp) + \ell_p(M^\sharp) \leq 0 + \text{rk}(M) = 8 - \text{rk}(N). \quad \square$$

Proposition 3.12. *If an isometry $g \in \text{O}(\mathbf{L})$ of order 8 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to \mathbf{D}_5 , and condition [\(2\)](#) holds.*

Proof. Let $N := \mathbf{L}_g$. By inspection of all cases in [Proposition 3.9](#), we see that there exists a primitive embedding $\mathbf{A}_3 \hookrightarrow \mathbf{L}_{g^2}$. Hence, thanks to [Theorem 2.17](#), [Lemma 3.8](#), and the existence of $\mathbf{L}_{g^2} \hookrightarrow N$, we infer that N is an 8-elementary negative definite lattice admitting primitive embeddings $\mathbf{A}_3 \hookrightarrow N \hookrightarrow \mathbf{L}$. In particular, $\text{rk}(N) \leq 5$. By [Lemma 3.11](#) we have $\ell_2(N^\sharp) \leq 3$, so $|\det(N)| \leq 8^3 = 512$.

A computation with the algorithm explained in [§2.4](#) shows that N must be isomorphic to either \mathbf{D}_4 , \mathbf{D}_5 or $\mathbf{D}_4 \oplus [-2]$. By [Lemma 3.3](#) and condition (9), we can exclude the case $N \cong \mathbf{D}_4 \oplus [-2]$. Finally, the restriction $g' = g|_N$ satisfies $|g'| = 8$, $N^{g'} = 0$ and $(g')^\sharp = \text{id}$. There is no such isometry $g' \in \text{O}(\mathbf{D}_4)$, hence only $N \cong \mathbf{D}_5$ is possible. \square

Proposition 3.13. *An isometry $g \in \text{O}(\mathbf{L})$ induced by a symplectic birational transformation cannot have order 16.*

Proof. Suppose such an isometry g exists. Then, $\mathbf{L}_{g^2} \cong \mathbf{D}_5$ by [Proposition 3.12](#). Now, \mathbf{L}_{g^2} embeds primitively into \mathbf{L}_g and \mathbf{L}_g is negative definite. Therefore,

$$5 = \text{rk}(\mathbf{L}_{g^2}) \leq \text{rk}(\mathbf{L}_g) \leq 5,$$

which forces $\text{rk}(\mathbf{L}_g) = 5$ and $\mathbf{L}_g \cong \mathbf{L}_{g^2} \cong \mathbf{D}_5$. But \mathbf{D}_5 admits no isometry of order 16, hence such g cannot exist. \square

3.4. Proof of [Theorem 1.2](#). Let $g \in \text{O}(\mathbf{L})$ be induced by a symplectic birational transformation and suppose that $g^\sharp \neq \text{id}$. Thanks to [Lemma 3.2](#), up to passing to an odd power of g we can suppose without loss of generality that $|g| = 2^k$ for some $k \in \mathbb{N}$.

From [Proposition 3.13](#) it follows that $k \leq 3$. Therefore, condition (2) holds because of [Proposition 3.5](#), [3.9](#) or [3.12](#). Consequently, it must be $g^\sharp = \text{id}$ by [Lemma 2.9](#), contradicting the assumption. \square

3.5. Other orders. [Theorem 1.2](#) and [Lemma 2.8](#) imply together that \mathbf{L}_g is $|g|$ -elementary if g is induced by a symplectic birational transformation, a fact which we will not repeat in the rest of the paper. In the next proof, the following lattice appears:

$$\tilde{\mathbf{A}} = \begin{pmatrix} -2 & 1 & -1 & -1 \\ 1 & -2 & 1 & 1 \\ -1 & 1 & -2 & 0 \\ -1 & 1 & 0 & -4 \end{pmatrix}.$$

Proposition 3.14. *If an isometry $g \in \text{O}(\mathbf{L})$ of order 6 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to one of the following lattices: $\mathbf{A}_2 \oplus [-2]$, \mathbf{D}_4 , $\mathbf{A}_2 \oplus 2[-2]$, $2\mathbf{A}_2 \oplus [-2]$. Moreover, condition (2) holds.*

Proof. Let $N := \mathbf{L}_g$. By [Proposition 3.6](#), it holds $\mathbf{L}_{g^2} \cong \mathbf{A}_2$ or $\mathbf{L}_{g^2} \cong 2\mathbf{A}_2$. Since $\mathbf{L}_{g^2} \hookrightarrow \mathbf{L}_g$, in both cases there must exist a (necessarily primitive) embedding $\mathbf{A}_2 \hookrightarrow N$. Hence, it holds $\ell_2(N^\sharp) \leq 3$ and $\ell_3(N^\sharp) \leq 3$ by [Lemma 3.11](#) and the fact that $\text{rk}(N) \leq 5$ because N is negative definite; consequently, $|\det(N)| \leq 6^3 = 216$. A computation shows that N must be isomorphic to one of the following lattices:

$$\begin{aligned} & \mathbf{A}_2, \mathbf{A}_2 \oplus [-2], \mathbf{A}_2 \oplus [-6], \mathbf{D}_4, 2\mathbf{A}_2, \tilde{\mathbf{A}}, \mathbf{A}_2 \oplus 2[-2], \mathbf{A}_2 \oplus \mathbf{A}_2(2), \mathbf{A}_2 \oplus [-2] \oplus [-6], \\ & \mathbf{A}_2 \oplus 2[-6], \mathbf{D}_5, \mathbf{D}_4 \oplus [-2], 2\mathbf{A}_2 \oplus [-2], \tilde{\mathbf{A}} \oplus [-2], \mathbf{A}_2 \oplus 3[-2], \mathbf{D}_4 \oplus [-6], 2\mathbf{A}_2 \oplus [-6], \\ & \mathbf{A}_2 \oplus 2[-2] \oplus [-6], \tilde{\mathbf{A}} \oplus [-6], \mathbf{A}_2 \oplus \mathbf{A}_2(2) \oplus [-2], \mathbf{A}_2 \oplus [-2] \oplus [-6], \mathbf{A}_2 \oplus \mathbf{A}_2(2) \oplus [-6]. \end{aligned}$$

Since 3 does not divide $\det(\mathbf{L})$, all elements of order 3 in N^\sharp must belong to the gluing subgroup H of $N \hookrightarrow \mathbf{L}$. Therefore, the restriction $g' = g|_N \in \mathrm{O}(N)$ satisfies $|g'| = 6$, $N^{g'} = 0$, and induces the identity on the elements of order 3 in N^\sharp . By inspection we see that only the following lattices admit such an isometry h :

$$\mathbf{A}_2 \oplus [-2], \mathbf{D}_4, \mathbf{A}_2 \oplus 2[-2], \mathbf{D}_5, \mathbf{D}_4 \oplus [-2], \mathbf{A}_2 \oplus [-2].$$

Lemma 3.3 implies that condition (2) must hold in order for $\mathcal{W}_{\mathrm{OG6}}^{\mathrm{pex}} \cap N = \emptyset$ to be true, hence $(g')^\sharp = \mathrm{id}$ because of **Lemma 2.9**. This excludes \mathbf{D}_5 and $\mathbf{D}_4 \oplus [-2]$. \square

Proposition 3.15. *An isometry $g \in \mathrm{O}(\mathbf{L})$ induced by a symplectic birational transformation cannot have order 9.*

Proof. By looking at g^3 and **Proposition 3.6**, we see that $N := \mathbf{L}_g$ is a 9-elementary negative definite lattice admitting primitive embeddings $\mathbf{A}_2 \hookrightarrow N \hookrightarrow \mathbf{L}$. Hence, it holds $\ell_3(N^\sharp) \leq 3$ by **Lemma 3.11**; consequently, $|\det(N)| \leq 9^3 = 729$. We compute that N can be either $\mathbf{A}_2, 2\mathbf{A}_2$ or $\mathbf{A}_2 \oplus \mathbf{A}_2(3)$, but none of these lattices admits an isometry of order 9. \square

Proposition 3.16. *If an isometry $g \in \mathrm{O}(\mathbf{L})$ of order 10 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to $\mathbf{A}_4 \oplus [-2]$, and condition (2) holds.*

Proof. Arguing as above, $N := \mathbf{L}_g$ must be a 10-elementary negative definite lattice admitting primitive embeddings $\mathbf{A}_4 \hookrightarrow N \hookrightarrow \mathbf{L}$. Thus, if $\mathrm{rk}(N) = 4$ then $N \cong \mathbf{A}_4$, but this lattice does not have any isometry $g \in \mathrm{O}(N)$ of order 10 acting trivially on the discriminant group such that $N^h = 0$.

Suppose that $\mathrm{rk}(N) = 5$. By **Lemma 3.11** it holds $\ell_2(N^\sharp) \leq 1$ and $\ell_5(N^\sharp) \leq 2$; consequently, $|\det(N)| \leq 2 \cdot 5^2 = 50$. We compute that $N \cong \mathbf{A}_4 \oplus [-2]$ or $N \cong \mathbf{A}_4 \oplus [-10]$, but the second lattice does not have admit any isometry $g' \in \mathrm{O}(N)$ of order 10 acting trivially on the elements of N^\sharp of order 5 such that $N^{g'} = 0$. \square

Proposition 3.17. *If an isometry $g \in \mathrm{O}(\mathbf{L})$ of order 12 is induced by a symplectic birational transformation, then \mathbf{L}_g is isomorphic to \mathbf{D}_5 , and condition (2) holds.*

Proof. Let $N := \mathbf{L}_g$, which is a 12-elementary negative definite lattice. Looking at g^3 , we see that there exists primitive embeddings $\mathbf{A}_3 \hookrightarrow N$ (by inspection of the cases in **Proposition 3.9**). We infer from **Lemma 3.11** that $\ell_2(N^\sharp) \leq 3$ and $\ell_3(N^\sharp) \leq 2$, so $|\det(N)| \leq 4^3 \cdot 3^2 = 576$. Therefore, N is isomorphic to one of the following lattices:

$$\begin{aligned} &\mathbf{A}_3 \oplus [-2], \mathbf{D}_5, \mathbf{A}_5, \mathbf{D}_4 \oplus [-2], \mathbf{A}_3 \oplus \mathbf{A}_2, \mathbf{A}_3 \oplus 2[-2], \tilde{\mathbf{A}} \oplus [-2], \mathbf{A}_3 \oplus [-2] \oplus [-4], \\ &\mathbf{A}_3 \oplus [-2] \oplus [-6], \mathbf{A}_3 \oplus [-2] \oplus [-12]. \end{aligned}$$

Since there must exist $g' \in \mathrm{O}(N)$ with $|g'| = 12$, we see that N is isomorphic to either \mathbf{D}_5 or $\mathbf{D}_4 \oplus [-2]$. But then condition (2) must hold because of **Lemma 3.3**, so it should also be $(g')^\sharp = \mathrm{id}$. Such an isometry g' exists only if $N \cong \mathbf{D}_5$. \square

Proposition 3.18. *An isometry $g \in \mathrm{O}(\mathbf{L})$ induced by a symplectic birational transformation cannot have order 15 or 25.*

Proof. Let $m \in \{15, 25\}$. The coinvariant sublattice $N := \mathbf{L}_g$ is an m -elementary negative definite lattice admitting primitive embeddings $\mathbf{A}_4 \hookrightarrow N \hookrightarrow \mathbf{L}$ by **Proposition 3.7**. Since

m is odd, we have necessarily $\text{rk}(N) = 4$ by [Lemma 2.7](#), hence $N \cong \mathbf{A}_4$, but there is no isometry in $O(\mathbf{A}_4)$ of order m . \square

Proposition 3.19. *An isometry $g \in O(\mathbf{L})$ induced by a symplectic birational transformation cannot have order 20 or 24.*

Proof. Suppose $|g| = 20$. Considering g^2 , we see from [Proposition 3.16](#) that it should be $\mathbf{L}_g \cong \mathbf{A}_4 \oplus [-2]$, but this lattice does not admit any isometry of order 20. An analogous argument holds when $|g| = 24$. \square

3.6. Proof of Theorem 1.3. Suppose that an isometry $g \in O(\mathbf{L})$ of finite order m is induced by a symplectic birational transformation on a manifold of type OG6. By [Theorem 2.17](#) the coinvariant sublattice \mathbf{L}_g is negative definite, hence by [Lemma 3.1](#) only 2, 3 and 5 appear as prime factors of m . As the orders 16, 9 and 25 are not possible by [Proposition 3.13](#), [3.15](#) and [3.18](#), respectively, it holds $m \mid 2^3 \cdot 3 \cdot 5 = 120$, which means

$$m \in \{1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120\}.$$

Now, [Proposition 3.18](#) excludes 15, 30, 60, 120, and [Proposition 3.19](#) excludes 20, 24, 40. Therefore, condition (1) holds.

From [Proposition 3.5](#), [3.6](#), [3.7](#), [3.9](#), [3.12](#), [3.14](#), [3.16](#) and [3.17](#), we know the isomorphism class of \mathbf{L}_g . Moreover, equation (2) always holds, which means that the embedding subgroup of $\mathbf{L}_g \hookrightarrow \mathbf{L}$ is trivial. This enables us to reconstruct the genus of \mathbf{L}^g , hence \mathbf{L}^g itself, being always unique in its genus. As a result of our computations, the pair $(\mathbf{L}^g, \mathbf{L}_g)$ appears in [Table 1](#).

Conversely, one can check that the column ‘example’ provides isometries with the given invariant and coinvariant lattices. By [Corollary 2.2](#), condition (9) of [Theorem 2.17](#) holds automatically. Therefore, all isometries contained in [Table 1](#) are indeed induced by a symplectic birational transformation on some marked pair of type OG6. This concludes the proof of the theorem. \square

REFERENCES

1. Samuel Boissière, Chiara Camere, and Alessandra Sarti, *Classification of automorphisms on a deformation family of hyper-Kähler four-folds by p -elementary lattices*, *Kyoto J. Math.* **56** (2016), no. 3, 465–499. MR 3542771
2. Samuel Boissière, Marc Nieper-Wißkirchen, and Alessandra Sarti, *Smith theory and irreducible holomorphic symplectic manifolds*, *J. Topol.* **6** (2013), no. 2, 361–390. MR 3065180
3. Simon Brandhorst and Alberto Cattaneo, *Prime order isometries of unimodular lattices and automorphisms of ihs manifolds*, preprint, [arXiv:1912.07119v3](#) (2020).
4. Simon Brandhorst, Serkan Sonel, and Davide Cesare Veniani, *Idoneal genera and K3 surfaces covering an Enriques surface*, preprint, [arXiv:2003.08914v6](#) (2022).
5. Chiara Camere, *Symplectic involutions of holomorphic symplectic four-folds*, *Bull. Lond. Math. Soc.* **44** (2012), no. 4, 687–702. MR 2967237
6. John H. Conway and Neil J. A. Sloane, *Low-dimensional lattices. IV. The mass formula*, *Proc. Roy. Soc. London Ser. A* **419** (1988), no. 1857, 259–286. MR 965484
7. ———, *Sphere packings, lattices and groups*, third ed., Grundlehren Math. Wiss., vol. 290, Springer-Verlag, New York, 1999, with additional contributions by E. Bannai, R. E. Borcherds, J. Leech, S. P. Norton, A. M. Odlyzko, R. A. Parker, L. Queen and B. B. Venkov. MR 1662447
8. Alice Garbagnati and Alessandra Sarti, *Symplectic automorphisms of prime order on K3 surfaces*, *J. Algebra* **318** (2007), no. 1, 323–350. MR 2363136

9. Annalisa Grossi, *Induced birational transformations on O'Grady's sixfolds*, J. London Math. Soc. **105** (2022), no. 1, 665–690.
10. ———, *Nonsymplectic automorphisms of prime order on O'Grady's sixfolds*, Rev. Mat. Iberoam. (2022), to appear, [doi:10.4171/RMI/1341](https://doi.org/10.4171/RMI/1341), [arXiv:2004.12694v3](https://arxiv.org/abs/2004.12694v3).
11. Kenji Hashimoto, *Finite symplectic actions on the K3 lattice*, Nagoya Math. J. **206** (2012), 99–153. MR 2926486
12. Brendan Hassett and Yuri Tschinkel, *Hodge theory and Lagrangian planes on generalized Kummer fourfolds*, Mosc. Math. J. **13** (2013), no. 1, 33–56, 189. MR 3112215
13. Gerald Höhn and Geoffrey Mason, *Finite groups of symplectic automorphisms of hyperkähler manifolds of type K3^[2]*, Bull. Inst. Math. Acad. Sin. (N.S.) **14** (2019), no. 2, 189–264. MR 3973459
14. Daniel Huybrechts, *Compact hyper-Kähler manifolds: basic results*, Invent. Math. **135** (1999), no. 1, 63–113. MR 1664696
15. ———, *A global Torelli theorem for hyperkähler manifolds [after M. Verbitsky]*, Astérisque (2012), no. 348, Exp. No. 1040, x, 375–403, Séminaire Bourbaki: Vol. 2010/2011. Exposés 1027–1042. MR 3051203
16. ———, *Lectures on K3 surfaces*, Cambridge Studies in Advanced Mathematics, vol. 158, Cambridge University Press, Cambridge, 2016. MR 3586372
17. ———, *On derived categories of K3 surfaces, symplectic automorphisms and the Conway group*, Development of moduli theory—Kyoto 2013, Adv. Stud. Pure Math., vol. 69, Math. Soc. Japan, [Tokyo], 2016, pp. 387–405. MR 3586514
18. Shigeyuki Kondō, *Niemeyer lattices, Mathieu groups, and finite groups of symplectic automorphisms of K3 surfaces*, Duke Math. J. **92** (1998), no. 3, 593–603, with an appendix by Shigeru Mukai. MR 1620514
19. ———, *A survey of finite groups of symplectic automorphisms of K3 surfaces*, J. Phys. A **51** (2018), no. 5, 053003, 17. MR 3755569
20. Eyal Markman, *A survey of Torelli and monodromy results for holomorphic-symplectic varieties*, Complex and differential geometry, Springer Proc. Math., vol. 8, Springer, Heidelberg, 2011, pp. 257–322. MR 2964480
21. Rick Miranda and David R. Morrison, *Embeddings of integral quadratic forms*, preliminary draft, web.math.ucsb.edu/~drm/manuscripts/eiqf.pdf, 2009.
22. Giovanni Mongardi, *Symplectic involutions on deformations of K3^[2]*, Cent. Eur. J. Math. **10** (2012), no. 4, 1472–1485. MR 2925616
23. ———, *On natural deformations of symplectic automorphisms of manifolds of K3^[n] type*, C. R. Math. Acad. Sci. Paris **351** (2013), no. 13–14, 561–564. MR 3095106
24. ———, *On symplectic automorphisms of hyper-Kähler fourfolds of K3^[2] type*, Michigan Math. J. **62** (2013), no. 3, 537–550. MR 3102529
25. ———, *Towards a classification of symplectic automorphisms on manifolds of K3^[n] type*, Math. Z. **282** (2016), no. 3–4, 651–662. MR 3473636
26. Giovanni Mongardi and Antonio Rapagnetta, *Monodromy and birational geometry of O'Grady's sixfolds*, J. Math. Pures Appl. (9) **146** (2021), 31–68. MR 4197280
27. Giovanni Mongardi, Kévin Tari, and Malte Wandel, *Prime order automorphisms of generalised Kummer fourfolds*, Manuscripta Math. **155** (2018), no. 3–4, 449–469. MR 3763414
28. Giovanni Mongardi and Malte Wandel, *Automorphisms of O'Grady's manifolds acting trivially on cohomology*, Algebr. Geom. **4** (2017), no. 1, 104–119. MR 3592467
29. David R. Morrison, *On K3 surfaces with large Picard number*, Invent. Math. **75** (1984), no. 1, 105–121. MR 728142
30. Shigeru Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, Invent. Math. **94** (1988), no. 1, 183–221. MR 958597
31. Viacheslav V. Nikulin, *Finite groups of automorphisms of Kählerian surfaces of type K3*, Uspehi Mat. Nauk **31** (1976), no. 2(188), 223–224 (Russian). MR 0409904
32. ———, *Finite groups of automorphisms of Kählerian K3 surfaces*, Trudy Moskov. Mat. Obshch. **38** (1979), 75–137 (Russian), English translation: Trans. Moscow Math. Soc. **38** (1980), no. 2, 71–35. MR 544937

33. ———, *Integer symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no. 1, 111–177 (Russian), English translation: Math USSR-Izv. **14** (1979), no. 1, 103–167 (1980). MR 525944
34. Kieran G. O’Grady, *A new six-dimensional irreducible symplectic variety*, J. Algebraic Geom. **12** (2003), no. 3, 435–505. MR 1966024
35. Antonio Rapagnetta, *Topological invariants of O’Grady’s six dimensional irreducible symplectic variety*, Math. Z. **256** (2007), no. 1, 1–34. MR 2282256
36. William A. Stein et al., *Sage Mathematics Software (Version 9.0)*, The Sage Development Team, 2020, www.sagemath.org.
37. Misha Verbitsky, *Mapping class group and a global Torelli theorem for hyperkähler manifolds*, Duke Math. J. **162** (2013), no. 15, 2929–2986, Appendix A by Eyal Markman. MR 3161308
38. ———, *Errata for “Mapping class group and a global Torelli theorem for hyperkähler manifolds” by Misha Verbitsky*, Duke Math. J. **169** (2020), no. 5, 1037–1038. MR 4079420
39. Gang Xiao, *Galois covers between K3 surfaces*, Ann. Inst. Fourier (Grenoble) **46** (1996), no. 1, 73–88. MR 1385511

LEHRSTUHL FÜR ALGEBRA UND ZAHLENTHEORIE, INSTITUT FÜR MATHEMATIK, UNIVERSITÄT AUGSBURG, UNIVERSITÄTSSTR. 14, 86135 AUGSBURG, GERMANY

Current address: Fakultät für Mathematik, Technische Universität Chemnitz, Reichenhainer Str. 39, 09126 Chemnitz, Germany

Email address: annalisa.grossi@math.tu-chemnitz.de

MATEMATISK INSTITUTT, UNIVERSITETET I OSLO, POSTBOKS 1053, BLINDERN, 0316 OSLO, NORWAY

Current address: Dipartimento di Matematica, Università degli studi di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Roma, Italy

Email address: onorati@mat.uniroma2.it

INSTITUT FÜR DISKRETE STRUKTUREN UND SYMBOLISCHES RECHNEN, UNIVERSITÄT STUTTGART, PFAFFENWALDRING 57, 70569 STUTTGART, GERMANY

Email address: davide.veniani@mathematik.uni-stuttgart.de