

RESEARCH ARTICLE

Open Access

Cyclic AMP induces apoptosis in multiple myeloma cells and inhibits tumor development in a mouse myeloma model

Virginie Follin-Arbelet^{1*}, Peter O Hofgaard², Harald Hauglin³, Soheil Naderi¹, Anders Sundan⁴, Rune Blomhoff³, Bjarne Bogen² and Heidi K Blomhoff¹

Abstract

Background: Multiple myeloma is an incurable disease requiring the development of effective therapies which can be used clinically. We have elucidated the potential for manipulating the cAMP signaling pathway as a target for inhibiting the growth of multiple myeloma cells.

Methods: As a model system, we primarily used the murine multiple myeloma cell line MOPC315 which can be grown both *in vivo* and *in vitro*. Human multiple myeloma cell lines U266, INA-6 and the B-cell precursor acute lymphoblastic leukemia cell line Reh were used only for *in vitro* studies. Cell death was assessed by flow cytometry and western blot analysis after treatment with cAMP elevating agents (forskolin, prostaglandin E2 and rolipram) and cAMP analogs. We followed tumor growth *in vivo* after forskolin treatment by imaging DsRed-labelled MOPC315 cells transplanted subcutaneously in BALB/c nude mice.

Results: In contrast to the effect on Reh cells, 50 μ M forskolin more than tripled the death of MOPC315 cells after 24 h *in vitro*. Forskolin induced cell death to a similar extent in the human myeloma cell lines U266 and INA-6. cAMP-mediated cell death had all the typical hallmarks of apoptosis, including changes in the mitochondrial membrane potential and cleavage of caspase 3, caspase 9 and PARP. Forskolin also inhibited the growth of multiple myeloma cells in a mouse model *in vivo*.

Conclusions: Elevation of intracellular levels of cAMP kills multiple myeloma cells *in vitro* and inhibits development of multiple myeloma *in vivo*. This strongly suggests that compounds activating the cAMP signaling pathway may be useful in the field of multiple myeloma.

Background

Multiple myeloma (MM) is a B-cell malignancy characterized by accumulation of plasma cells in the bone marrow, osteolytic bone lesions, and immunodeficiency [1]. It accounts for ~10% of hematological malignancies [2] with a median survival of 4 years [3]. Despite the progress made the last decades in the development of new therapies, multiple myeloma remains an incurable disease for which a constant search for new treatment strategies must continue.

Cyclic adenosine monophosphate (cAMP) is an intracellular messenger formed in response to diverse extracellular stimuli including hormones or neurotransmitters. It is generated from ATP by adenylyl cyclases, and is degraded by phosphodiesterases (PDE) into adenosine-5'-monophosphate. The main targets of cAMP are protein kinase A (PKA) [4], cAMP-gated ion channels [5] and exchange proteins directly activated by cAMP (EPAC) [6]. cAMP affects numerous cellular processes, such as cell differentiation, cell cycle progression and apoptosis, both in a PKA-dependent and PKA-independent manner [7-9]. In many cancer tissues and cell lines, alterations in cAMP signaling pathway including changes in intracellular levels of cAMP [10,11] and PKA isoforms ratio

* Correspondence: virginie.follin-arbelet@medisin.uio.no

¹Department of Biochemistry, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

Full list of author information is available at the end of the article

switch [12-15], have been observed. Consequently, there is a growing interest in manipulating the cAMP signaling pathway as a strategy for the treatment of cancer, and in particular a renewed interest for the potential of combining PDE inhibitors and glucocorticoids for treatment of hematological malignancies [16].

We have previously shown that cAMP blocks the G1/S phase transition and DNA synthesis in lymphoid cells [17-19]. More recently, we demonstrated that elevation of intracellular cAMP alone has no effect on cell death in B-cell precursor acute lymphoblastic leukemia (BCP-ALL) cells, but that it prevents apoptosis and accumulation of p53 in the cells subjected to γ -irradiation (γ -IR) [20]. In the present paper, we have explored the role of cAMP in multiple myeloma by primarily using the multiple myeloma cell line MOPC315. This cell line was chosen as it is a suitable mouse model [21,22] for studying the effect of cAMP on development of multiple myeloma *in vivo*. Elevation of intracellular levels of cAMP in the multiple myeloma cells did not prevent γ -IR-mediated death of the cells *in vitro*, but interestingly, cAMP alone efficiently killed the myeloma cells. More importantly, we could demonstrate that cAMP prevents the growth of multiple myeloma cells *in vivo*.

Methods

Chemicals, Antibodies

Forskolin and rolipram (Sigma; Saint Louis, MO, USA) were diluted in dimethyl sulfoxide (DMSO), 8CPT-cAMP (Biolog, Bremen, Germany) was diluted in distilled water, whereas prostaglandin E₂ (Cayman, Ann Arbor, MI, USA) was diluted in ethanol. Propidium iodide, DMSO, saponin, paraformaldehyde and bovine serum albumin (BSA) were purchased from Sigma. The cationic fluorescent carbocyanine dye 5,5',6,6'-Tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide (JC-1) was from Calbiochem (San Diego, CA, USA).

Antibodies against caspase 3 (8G10), caspase 9 (the mouse-specific 9504 and the human-specific 9502) and PARP were purchased from Cell Signaling Technologies (Danvers, MA, USA). P53 (fl393) antibody was purchased from Santa Cruz Biotechnology (Fremont, CA, USA). Antibody against GAPDH (Sigma) was used as a loading control. Anti-goat and anti-mouse HRP-conjugated secondary antibodies were purchased from Bio-Rad (Hercules, CA, USA).

Irradiation of the cells

Cells were irradiated using a ¹³⁷Cs source at 4.3 Gy/min.

Cell lines and cell culture

The BCP-ALL cell line Reh [23] was cultured as previously described [19]. The transplantable BALB/c

mineral oil-induced plasmacytoma cell line, MOPC315 [21], was used to generate a subline, MOPC315.4, that grew well *in vitro* and *in vivo* [24]. A subline of MOPC315.4, MOPC315.BM (Bogen et al., unpublished), was used for the present experiments. Some experiments employed MOPC315.BM labeled with the fluorescent protein DsRed. For simplicity, the MOPC315.BM subline will be referred to as MOPC315 throughout the paper. The cells were cultured *in vitro* in RPMI 1640 (Invitrogen, Paisley, UK) containing 2 mM L-glutamine, supplemented with MEM non essential amino acid (Sigma), 1 mM sodium pyruvate (Sigma), 50 μ M monothioglycerol (Sigma), 12 μ g/ml gentamycin (Sigma) and 10% heat-inactivated FBS (Lonza, Verviers, Belgium). The human multiple myeloma cell line, INA-6 cells, was a kind gift from Dr. M. Gramatzki (Erlangen, Germany) and were cultured in RPMI 1640 (Invitrogen) containing 2 mM L-glutamine (Invitrogen), supplemented with 1 ng/ml Il-6 (Invitrogen), 12 μ g/ml gentamycin (Sigma) and 10% heat-inactivated FBS (Lonza). The U266 cell line was purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA) and was cultured in RPMI 1640 (Invitrogen) containing 2 mM L-glutamine (Invitrogen), supplemented with 15% heat-inactivated FBS (Sigma), 100 U/ml penicillin (Invitrogen), and 100 μ g/ml streptomycin (Invitrogen).

Flow cytometry

Flow cytometry analysis was performed on a FACS Calibur (Becton-Dickinson). For determination of cell viability by exclusion of propidium iodide (PI), 500 μ l of cell culture were incubated with 20 μ g/ml PI for 10 min at room temperature prior to analysis. The cationic fluorescent carbocyanine dye, 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide (JC-1) was used to assess changes in the mitochondrial membrane potential ($\Delta\Psi$ m) observed in apoptotic cells. Cells were incubated for 15 min at 37°C with 15 μ g/ml JC-1 before analysis. For determination of apoptotic cells, TUNEL assays were performed by using an In Situ Cell Death Detection Kit, Fluorescein from Roche (Mannheim, Germany). Briefly, cells were washed in ice cold PBS before being fixed with 4% paraformaldehyde and permeabilized with 0.1% saponin. Cells were washed in ice cold PBS before incubation in the TUNEL reaction mix for 1 h at 37°C. After washing the cells 3 times, the cells were analyzed by flow cytometry.

Immunoblot analysis

Cells were lysed in RIPA buffer (50 mM Tris [pH7.5], 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5 mM EDTA, 50 mM NaF, 10 mM β -glycerophosphate, 1 mM Na₃VO₄, 0.2 mM phenylmethylsulfonyl fluoride [PMSF], 10 μ g/ml leupeptin, and 0.5% aprotinin) and an equal amount

of proteins (50 µg) was separated by SDS-PAGE (Bio-Rad) electrophoresis. After transfer to a nitrocellulose membrane (GE Healthcare) using a semidry transfer cell (Bio-Rad), proteins were detected by standard immunoblotting procedures. In brief, the nitrocellulose membranes were washed in Tris buffered saline and 0.1% Tween (TBST) and incubated in blocking solution (5% non-fat dry milk in TBST or 5% BSA in TBST) at room temperature. After washing, the membranes were incubated overnight at 4°C with primary antibodies diluted in blocking solution. After washing in TBST, the membranes were incubated for 1 h with HRP-conjugated secondary antibody diluted in blocking solution, followed by a final washing at room temperature. Immunoreactive proteins were visualized with the enhanced chemiluminescence detection system (ECL, Amersham Pharmacia Biotech, UK) or the SuperSignal[®] west Dura Extended Duration substrate (Thermo Scientific, Rockford, IL, USA) according to the manufacturer's protocol.

Mouse model for multiple myeloma

Adult BALB/c nude mice (purchased from Charles River, Germany) were injected subcutaneously in the interscapular region with 5×10^5 tumor MOPC315. DsRed cells suspended in 100 µL PBS. Two days after injection of the cells, 5 mice were injected intraperitoneally with 4-5 mg/kg forskolin diluted in a PBS/DMSO solution (15:0.1), and 5 mice were injected with the vehicle. In a separate experiment, forskolin (or vehicle) was injected 3 times on days 2, 4 and 6. Tumor growth was followed daily by palpation and imaging. Mice with tumor diameters of 15-20 mm were killed by cervical dislocation. The study was approved by the National Committee for Animal Experiments.

In vivo imaging of mice

Mice were anaesthetized with 2.5% isoflurane (Baxter As, Norway). Immediately afterwards, they were placed in a light-sealed imaging chamber and kept anaesthetized throughout the imaging period.

Images were acquired using a combination of excitation (30 nm passband) and emission (20 nm passband) filters on an IVIS Spectrum Imaging System (Caliper Life Sciences). The following spectral channels were used (excitation:emission center wavelength in nm): 465:540, 465:580, 535:600 and 570:620. Spectral images were recorded in units of photons/second/cm²/sr and imported as 32 bit floating point TIFF files into Mathematica 5.2 (Wolfram Research) for further processing. Images were scaled with an excitation light correction factor [25] yielding normalized fluorescence efficiency (NFE) images for further processing. Background reference autofluorescence spectrum was recorded from the interscapular region on day 0 before MOPC315

injection. A reference MOPC315.DsRed spectrum was determined from a region containing a localized tumor (day 5) with the reference autofluorescence subtracted. MOPC315.DsRed specific signal was determined by linear (pseudo-inverse) unmixing [26], yielding DsRed fluorescence maps, which were thresholded, intensity color-coded and overlaid a white light illuminated image. Quantification of MOPC315.DsRed fluorescence was done by computing the total DsRed fluorescence for above-threshold pixels for each animal.

Statistical analysis


The paired-samples t-test was applied to check the significance in cell line experiments, using the PASW Statistic 18 software for windows. In all the figures, histograms show mean values of the indicated number of experiments, with error bars corresponding to SEM values. For *in vivo* experiments, the Wilcoxon signed-rank test was used to determine significant differences between 2 groups of mice.

Results

Elevation of cAMP levels by forskolin induces death of multiple myeloma cells

We have previously shown that in B-cell precursor acute lymphoblastic leukemia (BCP-ALL) cells, elevated intracellular levels of cAMP prevent apoptosis induced by a variety of DNA-damaging cytotoxic agents, including ionizing radiation (IR). We have also demonstrated that destabilization of p53 is a key feature in this process [20]. Now we have compared the effects of the adenylyl cyclase-activating diterpine forskolin [27] on IR-mediated cell death of the BCP-ALL cell line Reh and the plasmacytoma cell line MOPC315. Figure 1A (left panel) shows that forskolin (50 µM) alone had no effect on the viability of the Reh cells, but unsurprisingly prevented IR-induced apoptosis, as measured 24 h later. In sharp contrast, forskolin did not prevent IR-mediated death of the myeloma cells (Figure 1A, right panel), but rather potentiated the effect of irradiation. More intriguingly, after 24 hours of sole forskolin treatment, the percentage of dead MOPC315 cells increased from ~16% to ~50%.

To confirm the differential effect of forskolin on IR-treated Reh cells and MOPC315 cells, the effect of elevating cAMP on p53 expression was analyzed. In accordance with our previous result [20], forskolin prevented the IR-induced stabilization of p53 in Reh cells (Figure 1B). In contrast forskolin had no effect on p53 induced by IR in MOPC315 cells. Importantly, forskolin alone decreased the p53 levels in MOPC315 cells, indicating that induction of p53 is not involved in cAMP-mediated cell death of MOPC315 cells. Myeloma cells were notably more sensitive to irradiation than Reh cells; only 2



Gy was used to obtain similar death in MOPC315 cells compared to 10 Gy in Reh cells.

Dose- and time-dependent effects of forskolin on death of MOPC315 cells are mediated via cAMP

MOPC315 cells were treated for 24 h with increasing doses of forskolin, or with 50 µM of forskolin, at various time points. Cell death was measured by incorporation of PI. Forskolin induced death of MOPC315 cells was both dose- and time-dependent (Figure 2A and 2B, respectively). Death occurred at doses as low as 0.1 µM forskolin, and statistically significant death could be detected already after 8 h with 50 µM forskolin.

In addition to the activation of adenylyl cyclase, forskolin has been reported to modulate other cellular processes, such as ion channels [28,29]. We tested the effect of other cAMP increasing agents to verify that forskolin-induced cell death was mediated by

intracellular accumulation of cAMP. This included the cell membrane permeable cAMP analog 8-chlorophenylthio-cAMP (8CPT-cAMP) and prostaglandin E2 (PGE2), which increases intracellular levels of cAMP through the activation G-protein-coupled receptors [30,31]. MOPC315 cells were killed in a dose-dependent manner by 8CPT-cAMP or PGE2 (Figure 2C and 2D, respectively), supporting the notion that forskolin kills the MOPC315 cells via induction of cAMP.

cAMP induces death of human multiple myeloma cells

To verify that the killing of murine multiple myeloma cells by cAMP was applicable to human myeloma cells, the human multiple myeloma cell line U266 and the IL-6-dependent human myeloma cell line INA-6 were included. The cells were treated for 24 h with increasing doses of forskolin or 8CPT-cAMP, and cell death was assessed by PI incorporation. Both human multiple




Figure 2 cAMP induces cell death in MOPC315 cells. MOPC315 cells were incubated for 24 h with the indicated concentrations of forskolin (A), with vehicle (open circles) or 50 μM forskolin (closed circles) for the indicated incubation times (B), or with the stated concentrations of 8-CPT-cAMP (C) or prostaglandin E2 (D) for 24 h. Cell death was measured by PI-staining, and the results are presented as percentage of PI-positive cells +/-SEM. A,C and D, n = 4; B, n = 3. *P < 0.05 relative to cells treated with vehicle.

myeloma cell lines were sensitive to intracellular elevation of cAMP (Figure 3A, B and 3C), indicating that elevation of intracellular levels of cAMP indeed also induces cell death in human multiple myeloma cells. By using a cAMP Biotrack enzymeimmunoassay (EIA), we verified that forskolin increased the intracellular cAMP concentrations in these two cell lines (data not shown). To further confirm the involvement of the cAMP signaling pathway in killing multiple myeloma cells, INA-6 cells were treated for 24 hours with a low dose of

forskolin in combination with rolipram, an inhibitor of PDE4. Rolipram or a low dose of forskolin alone induced little or no cell death, whereas a combination of the two compounds markedly increased cell death (Figure 3D).

cAMP induces apoptotic cell death in multiple myeloma cells

To ascertain whether cAMP induces cell death by apoptosis, MOP315 cells treated with forskolin or 8CPT-cAMP




Figure 3 cAMP induces cell death in human multiple myeloma cell line. U266 cells and INA-6 cells were incubated for 24 h with the stated doses of forskolin (A and B) or 8-CPT-cAMP(C) or rolipram and forskolin(D). Cell death was assessed by PI staining. The results are presented as the mean values \pm SEM of 3 experiments.

were analyzed for DNA fragmentation by TdT-mediated dUTP nick end labeling (TUNEL) technique, or by analysis of changes in mitochondrial membrane potential ($\Delta\Psi_m$) by staining the cells with JC-1. Forskolin and 8CPT-cAMP induced similar percentage of dead cells whether cell death was measured as percentage of cells with fragmented DNA (TUNEL assay), by changes in mitochondrial membrane potential (JC-1 staining), or by simple incorporation of PI (Figure 4A), clearly suggesting that cAMP induces apoptotic death of multiple myeloma cells.

To verify cAMP induced death of myeloma cells to be apoptotic, we investigated the downstream events

following mitochondrial depolarization. Mitochondrial outer membrane permeabilization results in the release of cytochrome C from the intermembrane space into the cytosol, triggering the assembly of the caspase-activating complex that mediates autocleavage and activation of caspase 9 [32]. Once activated, caspase 9 activates downstream effector caspases such as caspase 3 provoking the cleavage of several proteins, such as PARP, which ultimately leads to cell destruction [33]. MOPC315 cells and INA-6 cells were treated with 50 μ M forskolin or vehicle, and expression of cleaved caspase 3, caspase 9 and PARP were examined by western




Figure 4 cAMP induces apoptotic cell death. (A) MOPC315 cells were treated for 24 h with 200 μM 8-CPT-cAMP or 50 μM forskolin. Overall cell death, changes in mitochondrial membrane potential and DNA breaks were assessed by PI staining, JC-1 staining, and TUNEL analysis, respectively. The results are presented as mean values±SEM of 3 experiments. (B) MOPC315 cells (left panel) and INA-6 cells (right panel) were treated with or without 50 μM forskolin. At the indicated time points, cells were collected and subjected to western blot analysis with antibodies recognizing full length and cleaved products of caspase 9, caspase 3 and PARP. One of 3 representative experiments is shown.

blot analysis. Forskolin induced profound cleavage of caspase 9, caspase 3 and PARP in INA-6 cells, and to a lesser extent in MOPC315 cells (Figure 4B), confirming that cAMP indeed kills human and murine multiple myeloma cells by activating the apoptotic machinery

cAMP delays growth of multiple myeloma cells in vivo

Having shown that elevation of intracellular cAMP kills multiple myeloma cells in vitro, we explored the

therapeutic potential of cAMP-elevating compounds on tumor growth in vivo, taking advantage of a previously established mouse model for multiple myeloma based on subcutaneous injection of MOPC315 cells [24] pre-labeled with the fluorescent protein DsRed [34]. BALB/c nude mice were subcutaneously injected between the shoulders with 5×10^5 MOPC315 cells stably transfected with the gene encoding the fluorescent protein DsRed (MOPC315.DsRed cells). Two days after


inoculation of the cells, 5 mice were intraperitoneally injected with a single dose of forskolin (4-5 mg/kg), whereas 5 mice were injected with the same volume of vehicle. Tumor size was followed daily by *in vivo* imaging of DsRed fluorescence using an IVIS Spectrum Imaging System from Caliper Life Sciences. All 10 mice eventually developed tumors, but a single dose of forskolin substantially delayed the tumor growth *in vivo* (Figure 5A). Similar results were obtained in a separate experiment where mice were injected 3 times with forskolin or vehicle on days 2, 4, and 6 after tumor cells injection (Figure 5B). Statistical differences between vehicle treated and forskolin treated mice is achieved ($p < 0.05$) from day 6 after tumor cell injection. Figure 5C

shows *in vivo* images of mice taken at day 7. Together, these results suggest that cAMP-elevating compounds may indeed have a therapeutic potential in treatment of multiple myeloma.

Discussion

We have demonstrated that intracellular elevation of cAMP levels efficiently kills both murine and human multiple myeloma cells *in vitro*, and that the cAMP-elevating compound forskolin markedly delays the *in vivo* growth of multiple myeloma cells in a mouse model.

Modulation of intracellular cAMP by directly increasing the level of cAMP in the cell or by inhibiting PDE has become an interesting approach to cancer therapy



[16,35,36 for reviews]. In a phase-II study, theophylline, a methylxanthine that inhibits PDEs, proved to be effective in patients with chronic lymphocytic leukemia [37]. Activation of the cAMP pathways may either induce or inhibit cell proliferation or cell death depending on the cell type, and from our own research it is clear that the effect of cAMP also varies between different types of lymphoid cells. Thus, whereas elevation of intracellular cAMP inhibits DNA-damage induced apoptosis and p53 stabilization in BCP-ALL cells and normal B- and T cells [20], no such effects were seen in myeloma cells. It is possible that the inability of cAMP to prevent the IR-induced stabilization of p53 in myeloma cells could explain why cAMP is unable to counteract IR-mediated apoptosis in these cells.

Why myeloma cells and not BCP-ALL cells are so efficiently killed by solely elevating the level of cAMP is, however, unclear. The different players in the cAMP signaling pathway are highly compartmentalized in the cells, with G-coupled receptors, adenylyl cyclases, PKAs, Epacs, and phosphodiesterases all being brought in close proximity in distinct signalosomes within the cells [38]. It is possible that the activity of distinct signalosomes might contribute to localized, yet physiological significant differences in response to activating the cAMP signal in different lymphoid subpopulations. We also observed variations in the sensitivity to forskolin between the different myeloma cell lines used. This could presumably be due to variations in level and/or activity of the various components of the cAMP/PKA pathways in the different cell lines.

In an early paper [39], it was shown that cAMP analogs including 8-chloro-cAMP, dibutyryl-cAMP and 8-bromo-cAMP inhibited cell growth and induced cell death in glucocorticoid sensitive and resistant multiple myeloma cell lines. However, it was subsequently concluded that 8-chloro-cAMP mediated the cytotoxicity via its metabolite 8-chloro-adenosine (8Cl-AD) and not via the cAMP pathway [40,41]. Therefore, the potential for cAMP-elevating compounds in therapy of multiple myeloma was not further pursued. Recently, however, in an interesting study by Rickles and coworkers using a high throughput screening (cHTS) platform to identify new drugs to combine with existing therapeutic strategies for multiple myeloma [42], it was discovered that the agonist of the adenosine A_{2A} receptor as well as phosphodiesterase (PDE) inhibitors synergized with glucocorticoids to inhibit cell proliferation and induce death of multiple myeloma cells [42], thereby supporting our present results.

A key finding in the present study was the novel demonstration of the ability of the cAMP elevating agent forskolin to inhibit the *in vivo* growth of multiple myeloma cells in a mouse model. It is not yet clear whether this reduced

tumor growth is due to induced tumor cell death. Tumors eventually also developed in forskolin-treated mice, which could be due to the outgrowth of a small portion of forskolin-resistant cells. Attempts to give 3 doses of forskolin spaced 2 days apart did not markedly improve the effect on tumor growth compared to a single dose. A combination of cAMP-elevating compounds and conventional therapeutic agents could probably improve the outcome. The enhanced killing of myeloma cells we observed *in vitro* by combining forskolin and γ -irradiation supports this strategy. Based on the findings by Rickles et al [42], it will also be interesting to test the combination of cAMP elevating agents, phosphodiesterase inhibitors and glucocorticoids on the *in vivo* growth of multiple myeloma cells. It is clear that the potential for cAMP in the field of multiple myeloma is revitalized.

Conclusion

Stimulation of the cAMP-signaling pathway not only kills human and murine multiple myeloma cells *in vitro*, but it also reduces *in vivo* growth of multiple myeloma cells in a mouse model. Elevation of cAMP kills the cells via classical apoptotic mechanisms involving mitochondrial membrane-changes and activation of caspases. These results support the potential use of cAMP elevating agents as targets against multiple myeloma.

Abbreviations used

BCP-ALL: B-cell precursor acute lymphoblastic leukemia; cAMP: Cyclic adenosine monophosphate; IR: Ionizing radiation; MOPC: Mineral Oil-induced Plasmacytoma; MM: Multiple myeloma; PDE: phosphodiesterases; PGE₂: prostaglandin E₂; PKA: Protein kinase A; PI: Propidium Iodide; TUNEL: TdT-mediated dUTP nick end labeling; 8CPT-cAMP: 8-chlorophenylthio-cAMP.

Acknowledgements

The authors are grateful to Britt Fux-Nilsen, Camilla Solberg and Hanne Hella for excellent technical help. The work was supported by the Norwegian Cancer Society (AS), the Jahre Foundation, The Blix Family Foundation, and Rachel and Otto Kr. Bruum's Foundation.

Author details

¹Department of Biochemistry, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway. ²Centre for Immune Regulation, Institute of Immunology, University of Oslo and Oslo University Hospital, Rikshospitalet, Oslo, Norway. ³Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway. ⁴Department of Cancer Research and Molecular Medicine, NTNU, Trondheim, Norway.

Authors' contributions

VFA designed the research, performed experiments, analyzed data and wrote the paper; POH helped designing, performing, analyzing data for *in vivo* research and helped in writing the paper; HH designed and analyzed *in vivo* imaging data and helped in writing the paper, SN helped designing the research, analyzing the data and writing the paper. AS provided material and helped in writing of the paper; RB helped designing the research and writing the paper; BB provided material, helped designing the research and writing the paper; HKB designed the research, analyzed data, and wrote the paper. All authors have read and approved the manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 14 January 2011 Accepted: 18 July 2011
Published: 18 July 2011

References

1. Kyle RA, Rajkumar SV: **Multiple myeloma.** *Blood* 2008, **111**:2962-2972.
2. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ: **Cancer statistics, 2009.** *CA Cancer J Clin* 2009, **59**:225-249.
3. Kumar SK, Rajkumar SV, Dispenzieri A, Lacy MQ, Hayman SR, Buadi FK, Zelderust SR, Dingli D, Russell SJ, Lust JA, Greipp PR, Kyle RA, Gertz MA: **Improved survival in multiple myeloma and the impact of novel therapies.** *Blood* 2008, **111**:2516-2520.
4. Tasken K, Aandahl EM: **Localized effects of cAMP mediated by distinct routes of protein kinase A.** *Physiol Rev* 2004, **84**:137-167.
5. Kaupp UB, Seifert R: **Cyclic nucleotide-gated ion channels.** *Physiol Rev* 2002, **82**:769-824.
6. de Rooij J, Zwartkruis FJ, Verheijen MH, Cool RH, Nijman SM, Wittinghofer A, Bos JL: **Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP.** *Nature* 1998, **396**:474-477.
7. Grandoch M, Roscioni SS, Schmidt M: **The role of Epac proteins, novel cAMP mediators, in the regulation of immune, lung and neuronal function.** *Br J Pharmacol* 2010, **159**:265-284.
8. Stork PJ, Schmitt JM: **Crosstalk between cAMP and MAP kinase signaling in the regulation of cell proliferation.** *Trends Cell Biol* 2002, **12**:258-266.
9. Kobsar A, Heeg S, Krohne K, Opitz A, Walter U, Bock M, Gambaryan S, Eigenthaler M: **Cyclic nucleotide-regulated proliferation and differentiation vary in human hematopoietic progenitor cells derived from healthy persons, tumor patients, and chronic myelocytic leukemia patients.** *Stem Cells Dev* 2008, **17**:81-91.
10. Marko D, Romanakis K, Zankl H, Furstenberger G, Steinbauer B, Eisenbrand G: **Induction of apoptosis by an inhibitor of cAMP-specific PDE in malignant murine carcinoma cells overexpressing PDE activity in comparison to their nonmalignant counterparts.** *Cell Biochem Biophys* 1998, **28**:75-101.
11. Monahan TM, Marchand NW, Fritz RR, Abell CW: **Cyclic adenosine 3':5'-monophosphate levels and activities of related enzymes in normal and leukemic lymphocytes.** *Cancer Res* 1975, **35**:2540-2547.
12. Cho-Chung YS, Nesterova MV: **Tumor reversion: protein kinase A isozyme switching.** *Ann N Y Acad Sci* 2005, **1058**:76-86.
13. McDaid HM, Cairns MT, Atkinson RJ, McAleer S, Harkin DP, Gilmore P, Johnston PG: **Increased expression of the R1alpha subunit of the cAMP-dependent protein kinase A is associated with advanced stage ovarian cancer.** *Br J Cancer* 1999, **79**:933-939.
14. Mantovani G, Bondioni S, Lania AG, Rodolfo M, Peverelli E, Polentarutti N, Veliz Rodriguez T, Ferrero S, Bosari S, Beck-Peccoz P, Spada A: **High expression of PKA regulatory subunit 1A protein is related to proliferation of human melanoma cells.** *Oncogene* 2008, **27**:1834-1843.
15. Neary CL, Nesterova M, Cho YS, Cheadle C, Becker KG, Cho-Chung YS: **Protein kinase A isozyme switching: eliciting differential cAMP signaling and tumor reversion.** *Oncogene* 2004, **23**:8847-8856.
16. Lerner A, Epstein PM: **Cyclic nucleotide phosphodiesterases as targets for treatment of hematological malignancies.** *Biochem J* 2006, **393**:21-41.
17. Blomhoff HK, Smeland EB, Beiske K, Blomhoff R, Ruud E, Bjoro T, Pfeifer-Ohlsson S, Watt R, Funderud S, Godal T: **Cyclic AMP-mediated suppression of normal and neoplastic B cell proliferation is associated with regulation of myc and Ha-ras protooncogenes.** *J Cell Physiol* 1987, **131**:426-433.
18. Blomhoff HK, Blomhoff R, Stokke T, deLange DC, Brevik K, Smeland EB, Funderud S, Godal T: **cAMP-mediated growth inhibition of a B-lymphoid precursor cell line Reh is associated with an early transient delay in G2/M, followed by an accumulation of cells in G1.** *J Cell Physiol* 1988, **137**:583-587.
19. Naderi S, Wang JY, Chen TT, Gutzkow KB, Blomhoff HK: **cAMP-mediated inhibition of DNA replication and S phase progression: involvement of Rb, p21Cip1, and PCNA.** *Mol Biol Cell* 2005, **16**:1527-1542.
20. Naderi EH, Findley HW, Ruud E, Blomhoff HK, Naderi S: **Activation of cAMP signaling inhibits DNA damage-induced apoptosis in BCP-ALL cells through abrogation of p53 accumulation.** *Blood* 2009, **114**:608-618.
21. Eisen HN, Simms ES, Potter M: **Mouse myeloma proteins with antihapten antibody activity. The protein produced by plasma cell tumor MOPC-315.** *Biochemistry* 1968, **7**:4126-4134.
22. Potter M: **Immunoglobulin-producing tumors and myeloma proteins of mice.** *Physiol Rev* 1972, **52**:631-719.
23. Rosenfeld C, Goutner A, Choquet C, Venuat AM, Kayibanda B, Pico JL, Greaves MF: **Phenotypic characterisation of a unique non-T, non-B acute lymphoblastic leukaemia cell line.** *Nature* 1977, **267**:841-843.
24. Lauritzsen GF, Bogen B: **The role of idiotype-specific, CD4+ T cells in tumor resistance against major histocompatibility complex class II molecule negative plasmacytoma cells.** *Cell Immunol* 1993, **148**:177-188.
25. Troy T, Jekic-McMullen D, Sambucetti L, Rice B: **Quantitative comparison of the sensitivity of detection of fluorescent and bioluminescent reporters in animal models.** *Mol Imaging* 2004, **3**:9-23.
26. Zimmermann T, Rietdorf J, Pepperkok R: **Spectral imaging and its applications in live cell microscopy.** *FEBS Lett* 2003, **546**:87-92.
27. Seamon KB, Padgett W, Daly JW: **Forskolin: unique diterpene activator of adenylate cyclase in membranes and in intact cells.** *Proc Natl Acad Sci USA* 1981, **78**:3363-3367.
28. Ono K, Fozzard HA, Hanck DA: **A direct effect of forskolin on sodium channel bursting.** *Pflugers Arch* 1995, **429**:561-569.
29. Laurenza A, Sutkowski EM, Seamon KB: **Forskolin: a specific stimulator of adenylate cyclase or a diterpene with multiple sites of action?** *Trends Pharmacol Sci* 1989, **10**:442-447.
30. Coleman RA, Smith WL, Narumiya S: **International Union of Pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes.** *Pharmacol Rev* 1994, **46**:205-229.
31. Negishi M, Sugimoto Y, Ichikawa A: **Prostanoid receptors and their biological actions.** *Prog Lipid Res* 1993, **32**:417-434.
32. Riedl SJ, Salvesen GS: **The apoptosome: signalling platform of cell death.** *Nat Rev Mol Cell Biol* 2007, **8**:405-413.
33. Taylor RC, Cullen SP, Martin SJ: **Apoptosis: controlled demolition at the cellular level.** *Nat Rev Mol Cell Biol* 2008, **9**:231-241.
34. Heikal AA, Hess ST, Baird GS, Tsien RY, Webb WW: **Molecular spectroscopy and dynamics of intrinsically fluorescent proteins: coral red (dsRed) and yellow (Citrine).** *Proc Natl Acad Sci USA* 2000, **97**:11996-12001.
35. Naviglio S, Caraglia M, Abbruzzese A, Chiosi E, Di Gesto GD, Marra M, Romano M, Sorrentino A, Sorvillo L, Spina A, Illiano G: **Protein kinase A as a biological target in cancer therapy.** *Expert Opin Ther Targets* 2009, **13**:83-92.
36. Savai R, Pullamsetti SS, Banat GA, Weissmann N, Ghofrani HA, Grimminger F, Schermuly RT: **Targeting cancer with phosphodiesterase inhibitors.** *Expert Opin Investig Drugs* 2010, **19**:117-131.
37. Wiernik PH, Paietta E, Goloubeva O, Lee SJ, Makower D, Bennett JM, Wade JL, Ghosh C, Kaminer LS, Pizzolo J, Tallman MS: **Phase II study of the theophylline in chronic lymphocytic leukemia: a study of the Eastern Cooperative Oncology Group (E4998).** *Leukemia* 2004, **18**:1605-1610.
38. Jarnaess E, Tasken K: **Spatiotemporal control of cAMP signalling processes by anchored signalling complexes.** *Biochem Soc Trans* 2007, **35**:931-937.
39. Krett NL, Zell JL, Halgren RG, Pillay S, Traynor AE, Rosen ST: **Cyclic adenosine-3',5'-monophosphate-mediated cytotoxicity in steroid sensitive and resistant myeloma.** *Clin Cancer Res* 1997, **3**:1781-1787.
40. Halgren RG, Traynor AE, Pillay S, Zell JL, Heller KF, Krett NL, Rosen ST: **8Cl-cAMP cytotoxicity in both steroid sensitive and insensitive multiple myeloma cell lines is mediated by 8Cl-adenosine.** *Blood* 1998, **92**:2893-2898.
41. Gandhi V, Ayres M, Halgren RG, Krett NL, Newman RA, Rosen ST: **8-chloro-cAMP and 8-chloro-adenosine act by the same mechanism in multiple myeloma cells.** *Cancer Res* 2001, **61**:5474-5479.
42. Rickles RJ, Pierce LT, Giordano TP, Tam WF, McMillin DW, Delmore J, Laubach JP, Borisy AA, Richardson PG, Lee MS: **Adenosine A2A receptor agonists and PDE inhibitors: a synergistic multitarget mechanism discovered through systematic combination screening in B-cell malignancies.** *Blood* 2010, **116**:593-602.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2407/11/301/prepub>

doi:10.1186/1471-2407-11-301

Cite this article as: Follin-Arbelet et al.: **Cyclic AMP induces apoptosis in multiple myeloma cells and inhibits tumor development in a mouse myeloma model.** *BMC Cancer* 2011 **11**:301.