

## Original Article

# Malignancies in Patients with Inflammatory Bowel Disease: Results from 20 Years of Follow-up in the IBSEN Study

Øistein Hovde,<sup>a,b</sup> Marte Lie Høivik,<sup>b</sup> Magne Henriksen,<sup>d</sup>  
Inger Camilla Solberg,<sup>c</sup> Milada Cvancarova Småstuen,<sup>c</sup> Bjørn A. Moum<sup>b,c</sup>

<sup>a</sup>Department of Gastroenterology, Innlandet Hospital Trust, Gjøvik, Norway <sup>b</sup>Institute of Clinical Medicine, University of Oslo, Oslo, Norway <sup>c</sup>Department of Gastroenterology, Oslo University Hospital, Oslo, Norway <sup>d</sup>Department of Gastroenterology, Østfold Hospital, Fredrikstad, Norway

Corresponding author: Øistein Hovde, MD, Department of Gastroenterology, Innlandet Hospital Trust, Kyrre Greppsgt. 19, 2819 Gjøvik, Norway. Tel.: +47 61157002; Fax: +47 61157439; email: [oistein.hovde@sykehuset-innlandet.no](mailto:oistein.hovde@sykehuset-innlandet.no)

## Abstract

**Background and Aims:** Whether patients with inflammatory bowel diseases [IBDs] have increased risk of developing cancer has been debated. The aims of the study were to determine the prevalence of intestinal/extraintestinal cancers in an IBD cohort 20 years after diagnosis and to assess whether these patients had an increased cancer-specific risk compared with a matched control population.

**Methods:** Patients with ulcerative colitis [UC] and Crohn's disease [CD] diagnosed 1990–1993 have been prospectively followed up for 20 years. Follow-up visits were carried out 1, 5, 10, and 20 years after inclusion. Data on all cancer cases, deaths, and causes of death were collected from the Cancer Registry of Norway and from the Norwegian Cause of Death Registry.

**Results:** In all, 756 patients [519 UC and 237 CD] were diagnosed with IBD. Increased risk of cancer was seen in UC patients (hazard ratio [HR] = 1.40, 95% confidence interval [CI] 1.08–1.81,  $p < 0.01$ ), but not in CD patients (HR = 1.23, 95% CI 0.80–2.03,  $p = 0.30$ ). Stratified by gender, our data revealed a statistically increased risk for all cancers only in male UC patients compared with the controls [HR = 1.51, 95% CI 1.08–2.11,  $p = 0.017$ ]. In both groups breast cancer was seen more often than expected.

**Conclusions:** Male UC patients display an increased risk of development of colorectal cancer and, also all cancers combined, compared with the controls. In both UC and CD, standardized incidence ratio for breast cancer was increased.

**Key Words:** Inflammatory bowel diseases; malignancies

## 1. Introduction

Crohn's disease [CD] and ulcerative colitis [UC] are chronic relapsing inflammatory diseases of the gastrointestinal tract collectively called inflammatory bowel disease [IBD]. The causes of the diseases remain unknown. Typically, IBD is diagnosed between the ages of 20 and 40 years and can cause a wide variety of clinical signs and symptoms.<sup>1–3</sup> Involvement of extraintestinal organs, such as joints, skin, bile ducts, and eyes, is not uncommon. The

highest reported prevalences of IBD are in Europe [827 per 100 000 inhabitants] and North America [578 per 100 000 inhabitants], and most time-trend analyses have shown a significantly increasing incidence, especially in infants and adolescents with CD. Robust epidemiological data from developing countries are lacking, but the incidence and prevalence are increasing throughout the world.<sup>4,5</sup> Therefore, it is important to obtain reliable data on the different disease outcomes.

IBD-related cancers are those attributable to chronic inflammation and/or to medical treatment of the diseases, i.e. immunosuppressive drugs and/or biologics.<sup>6</sup> Both intestinal and extraintestinal malignancies have been linked to IBD.<sup>7–12</sup> However, few longitudinal, prospective studies have focused on the association between IBD and the development of malignancies.

The aims of the present study were first to determine the prevalence of intestinal and extraintestinal cancers in a well-defined, population-based cohort of IBD patients 20 years after diagnosis and, second, to assess whether the IBD patients were at an increased cancer-specific risk compared with age- and gender-matched individuals from the normal population.

## 2. Materials and Methods

The Inflammatory Bowel South-Eastern Norway [IBSEN] study prospectively followed all patients [ $n = 756$ ] diagnosed with IBD from 1 January 1990, to 31 December 1993, in four geographically well-defined areas in south-eastern Norway. Each patient was matched with 25 individuals randomly selected from the general population and matched with respect to age, gender, and county of residence. On January 1, 1992, the total population in the area was 966 427. The criteria for inclusion in this population-based inception cohort have been described in detail elsewhere.<sup>13,14</sup>

Pre-scheduled follow-up visits were carried out at 1, 5, 10, and 20 [ $\pm 12$  months] years after inclusion. All visits included a clinical examination, a structured interview, and laboratory tests. Colonoscopies were performed at all of the visits, unless the patients objected, and otherwise performed as required. Data on all cancer cases, deaths, and causes of death were collected from the Cancer Registry of Norway and from the Norwegian Causes of Death Registry. The Cancer Registry of Norway contains detailed information on each case of cancer. Cancer information comes from several independent sources, thus ensuring a completeness of approximately 98.8%.<sup>15</sup> Moreover, all Norwegian citizens are assigned a unique digital identification number, which makes it possible to link data from several registries and enables highly reliable epidemiological research. All medical doctors in Norway are, by law, obliged to report new and suspected cancers to the Cancer Registry and whether the cancer is first diagnosed by autopsy. With regard to malignancies, all individuals in the study were observed from the date of IBD diagnosis until the first occurrence of cancer, death, or end of follow-up, whichever came first. The International Classification of Diseases [ICD] is the standard diagnostic tool for epidemiology, clinical purposes, and health management. All cancer diagnoses from the Cancer Registry of Norway used in this study were coded according to the ICD-10, and the first occurrence of cancer in each patient and in each of the controls was the diagnosis used in the registration. Our

registration did not allow us to distinguish between colonic and rectal cancer, as the diagnoses used included C18 [neoplasma malignum coli] and C19 [neoplasma malignum junctionis rectosigmoidalis], whereas C20 [neoplasma malignum recti] was not applied.

### 2.1. Definitions and disease classifications

Initially, the patients were classified as having UC, CD, indeterminate colitis [IC], or possible IBD as described previously.<sup>13,16</sup> Internationally accepted criteria were used for determining the diagnoses and subclassifications.<sup>14,17</sup>

The patients' diagnoses were systematically re-evaluated at the pre-scheduled visits at 1, 5, and 10 years after diagnosis. The diagnosis at the 10-year visit was set as the final IBD diagnosis.

Medical treatment was given in accordance with established clinical practice. During the first part of the study period, the patients were given 5-aminosalicylates [5-ASA], glucocorticoids, and thiopurines in a traditional step-up fashion,<sup>18,19</sup> whereas during the second half of the study, treatment with TNF [tumour necrosis factor]- $\alpha$  inhibitors with or without concomitant immunomodulators was introduced in the treatment of patients with chronic active or severe disease [Table 1].

### 2.2. Statistical analysis

Data were described with median and range. All IBD patients were age- and gender-matched with individuals selected at random from the general Norwegian population living in the same geographical region [county] [called controls]. Follow-up time was defined from the date of IBD diagnosis to the first occurrence of any cancer/specified cancer, death or end of follow-up, whichever came first. Event was defined as the occurrence of any cancer or the first occurrence of a specific cancer. All cancers and specific cancer risks for patients compared with controls were modelled using the Cox proportional model stratified by a matched set. Results are expressed as hazard ratios [HRs] with 95% confidence intervals [CIs]. In addition, for specified cancer groups SIRs [standardized incidence ratios] were computed as a ratio between observed and expected numbers. Expected numbers were derived from gender-stratified tables on cancer incidence in Norway provided by the Cancer Registry of Norway. CIs for the SIRs were constructed using Wilson and Hilferty approximation.<sup>20</sup> Significance level was set to 5%. All analyses were performed using SPSS V. 22.0 for Windows [SPSS; Chicago, IL, USA].

### 2.3. Ethics

The Regional Committee for Medical Ethics approved the study. The confidentiality of patient identity and records was maintained using the guidelines suggested by the National Health Department.

**Table 1.** Patients' characteristics at diagnosis and cumulative use of immunosuppressive drugs/TNF- $\alpha$  inhibitors during the entire study period.

Characteristics	Crohn's disease	Ulcerative colitis
Age, median [range]	28 yr [8–845]	39 yr [4–89]
Gender [females/males]	118/119	252/267
Treated with AZA	93/193 [50.8%] Missing = 54	48/376 [12.8%] Missing = 143
Treated with MTX	8/201 [4.0%] Missing = 36	6/447 [1.3%] Missing = 72
Treated with TNF- $\alpha$ inhibitors	42/176 [23.9%] Missing = 61	12/376 [3.2%] Missing = 143

AZA, azathioprine; MTX, methotrexate; TNF- $\alpha$  inhibitors, tumour necrosis factor- $\alpha$  inhibitors; yr, years.  
Treated: patients treated at any time during the study period.

### 3. Results

Our cohort comprised 756 patients diagnosed with IBD and 18900 age- and gender-matched controls. Demographic data are provided in [Table 1](#). There were no differences in age or gender between the diagnosis groups [data not shown]. The median follow-up time from diagnosis was 21 years [range 1 to 24 years]. Neither the IBD patients nor the controls had been diagnosed with any type of cancer by the start of the study.<sup>21,22</sup>

A total of 105 individuals in the IBD group [13.9%] developed cancer compared with 1830 in the control group [9.7%]. One subject in the IBD group and three subjects in the control group developed two consecutive cancers. IBD patients had a higher risk of cancer development than their matched controls; however, this increased risk was statistically significant only for UC patients [HR = 1.40, 95% CI 1.08–1.81,  $p < 0.01$  for UC patients; HR = 1.23, 95% CI 0.80–2.03,  $p = 0.30$  for CD patients]. When stratified by gender, our data revealed a higher risk for all cancers only for male UC patients [HR = 1.51, 95% CI 1.08–2.11,  $p = 0.017$ ]. The cancer risk for female UC patients was elevated compared with that of the controls; however, this increased risk did not reach statistical significance [[Table 2](#)]. For CD patients, there was no statistically significant difference in cancer risk between the cases and the controls. The observed and expected numbers of the most common malignancies and SIRs in UC and CD during follow-up are summarized in [Tables 3](#) and [4](#). Cumulative incidences for cancer development and death for male and female UC and CD patients are shown in [Figures 1](#) and [2](#). For both genders, cumulative incidence of cancer was higher in IBD patients compared with their controls, also when adjusted for competing risk of death.

#### 3.1. Colorectal cancer

Among female UC patients, 4/252 [1.6%] developed colorectal cancer [CRC], and among male UC patients, the number was 10/267 [3.7%]. In the control group 1.1% of the females and 1.3% of the males developed CRC.

In the CD group, 2/237 patients [0.84%] developed CRC compared with 56/5925 [0.94%] in the control group. The overall risk of males with UC of developing CRC was 2-fold higher compared with that of their controls [HR = 2.41, 95% CI 1.09–.31,  $p = 0.029$ ]. For female UC and all CD patients, there were no statistically significant differences in the CRC-specific risk between the cases and their matched controls [[Table 2](#)].

#### 3.2. Extraintestinal malignancies

In both UC and CD, the SIR for breast cancer was significantly increased. Moreover, CD patients exhibited an increased SIR for

trachea/lung cancer compared with the controls. The risk of developing malignant melanoma in the UC group was higher than in the control group, but the difference did not reach statistical significance. Generally, very few malignancies in the lymphoid/haematopoietic system were found, and in some diagnostic groups, the CI was so wide that conclusions must be drawn with caution [[Tables 2](#) and [3](#)].

### 4. Discussion

The present population-based study comprising 756 IBD patients followed for 20 years showed that the UC patients, but not the CD patients, had an increased risk of developing cancer compared with the controls, and male UC patients had an increased risk of developing CRC compared with the controls. The increased risk of having CRC in this UC cohort is in agreement with the findings in a recent meta-analysis of population-based cohort studies in UC patients,<sup>9</sup> in which 1.6% of the UC patients were diagnosed with CRC 14 years after diagnosis; SIRs ranged between 1.05 and 3.1. Men with UC had a greater risk than women [SIRs 2.6 vs 1.9]. In our study, 3.7% [10/267] of male and 1.6% [4/252] of female patients with UC developed CRC during the first 20 years after diagnosis.

A Danish study<sup>10</sup> did not find an increased overall risk of CRC in UC patients compared with the general population during the first 10 years after diagnosis, which is in accordance with results from the IBSEN study 10 years after UC diagnosis.<sup>23</sup> However, a colectomy rate more than twice as high as in other European centres indicates that there may have been a different treatment approach in the Danish study.<sup>23,24</sup>

Important risk factors for CRC include the duration of the colitis, together with the degree and extent of inflammation, in addition to the presence of primary sclerosing cholangitis and family history of CRC.<sup>25</sup> The patients gave information on family cancer history, but such information is lacking in the controls, and, therefore is not analysed further. In our study, we found a high SIR for biliary tract cancers in UC patients compared with the controls but the CI for this risk was wide, making it difficult to draw any conclusions.

In a long-term, nationwide population-based Danish cohort study with 30 years of follow-up, the authors concluded that patients with CD and UC were at increased risk for developing gastrointestinal and extraintestinal malignancies.<sup>26</sup> The authors further concluded that the risk of gastrointestinal malignancy has decreased in recent years and that this decrease has occurred without an increase in the overall risk of malignancy. This finding suggests that the benefits of current treatment strategies outweigh the cancer-related risks.

**Table 2.** Risk of cancer/CRC for IBD patients compared with controls: conditional Cox regression.

Variable	UC			CD		
	HR	95% CI	<i>p</i> -value	HR	95% CI	<i>p</i> -value
	All cancers combined					
Pts vs controls	1.40	1.08–1.81	0.010	1.28	0.80–2.03	0.302
Male pts vs controls	1.51	1.08–2.11	0.017	1.45	0.74–2.85	0.282
Female pts vs controls	1.28	0.86–1.90	0.227	1.15	0.61–2.18	0.662
	Colorectal cancers [CRCs]					
Pts vs controls	1.90	0.99–3.65	0.054	1.09	0.26–4.54	0.904
Male pts vs controls	2.41	1.09–5.31	0.029	1.43	0.19–11.05	0.729
Female pts vs controls	1.27	0.39–4.10	0.689	0.88	0.12–6.54	0.903

IBD, inflammatory bowel disease; UC, ulcerative colitis; CD, Crohn's disease; H, hazard ratio; CI, confidence interval; pts, patients.

**Table 3.** Observed and expected numbers of the most common malignancies and standard incidence rates [SIRs] among patients with ulcerative colitis during the entire follow-up period.

Site [ICD-10]	Observed [n]	Expected [n]	SIR	95% CI	p-value
Colon/rectum [C18-19]	14	6.36	2.20	1.253.61	0.008
Biliary tract [C23-24]	2	0.48	4.17	0.7013.77	n.s.
Trachea/lungs [C33-34]	6	5.52	1.09	0.442.26	n.s.
Melanoma [C43]	6	3.2	1.86	0.763.90	n.s.
Basal cell carcinoma [C44]	5	2.2	2.27	0.835.04	n.s.
Breast [C50]	13	6.48	2.00	1.123.34	0.021
Prostate [C61]	6	8.56	0.70	0.281.46	n.s.
Thyroid [C73]	2	0.36	5.56	0.9318.35	n.s.
Lymphatic leukaemia [C91]	6	1.36	4.41	1.799.18	0.003

ICD, International Classification of Diseases; CI, confidence interval; n.s., not significant.

**Table 4.** Observed and expected numbers of the most common malignancies and standard incidence rates [SIRs] among patients with Crohn's disease during the entire follow-up period.

Site [ICD-10]	Observed [n]	Expected [n]	SIR	95% CI	p-value
Colon/rectum [C18-19]	2	2.24	0.89	0.152.95	n.s.
Biliary tract [C23-24]	0	0.04	Cannot be computed		
Trachea/lungs [C33-34]	5	1.72	2.91	1.076.78	0.039
Melanoma [C43]	1	0.8	1.25	0.066.17	n.s.
Basal cell carcinoma [C44]	1	0.72	1.39	0.076.85	n.s.
Breast [C50]	6	2.48	2.42	0.985.03	n.s.
Prostate [C61]	3	2.24	1.34	0.343.65	n.s.
Thyroid [C73]	1	0.2	5.00	0.2524.66	n.s.
Lymphatic leukaemia [C91]	0	0.48	Cannot be computed		


ICD, International Classification of Diseases; CI, confidence interval; n.s., not significant.

The risk of malignancies was also assessed in a large-scale registry study from Finland, where 21 964 patients were followed up for 10.8 years.<sup>27</sup> A higher incidence of cancer among male patients with IBD compared with the background population was found, with a higher SIR for CD than for UC [1.23, 95% CI, 1.09–1.38 vs 1.08, 95% CI 1.02–1.14]. The authors reported a SIR of 1.73 [95% CI 1.30–2.26] in male UC patients and 1.92 [95% CI 1.38–2.59] in female patients for colon cancer, and 1.65 [95% CI 1.18–2.26] and 1.96 [95% CI 1.26–2.91] for male and female patients, respectively, for rectal cancer. In CD, no overall increased risk for developing CRC was observed in either gender, which is in accordance with our findings. Our study included substantially fewer patients but involved a follow-up time that was almost twice as long, as well as including a control group. The study from Finland also showed that UC patients had an excess number of biliary tract cancers [SIR = 7.26, 95% CI 4.37–11.13] and that male patients with UC had an increased risk of Hodgkin lymphomas [SIR = 2.45, 95% CI 1.06–4.81], although no association with use of immunomodulators was observed.


In general, there is a higher percentage of smokers/ex-smokers in the CD population than in the general population and in the UC population.<sup>13,33</sup> Therefore, it was surprising that the Finnish study did not find an increased SIR for lung/tracheal cancers [SIR = 1.14, 95% CI, 0.71–1.74] in this group. The UC patients, however, were found to have a slightly reduced risk of developing lung/tracheal cancers [SIR = 0.79, 95% CI, 0.62–0.98]. In our study, five CD patients developed lung/tracheal cancers compared with 1.72 expected [SIR = 2.91, 95% CI 1.07–6.78,  $p = 0.039$ ], whereas UC patients and their controls had equal incidences of these cancer types, but the exact relation to smoking habits was not calculated.

In a meta-analysis of eight population-based cohort studies from 1996 to 2009, comprising approximately 17 000 patients with IBD, the patients were found to have the same overall risk of developing extraintestinal cancers as the background population.<sup>12</sup> In this earlier study, CD patients had an increased risk of cancer of the upper gastrointestinal [GI] tract, lungs, urinary bladder, and skin. UC patients, in contrast, had an increased risk of liver/biliary tract cancers and leukaemia but a decreased risk of pulmonary cancer, which most likely occurred because UC patients smoke less than the background population. Our study confirms the increased risk for cancer of the respiratory tract in patients with CD. Our study also found an increased risk for breast cancer in IBD patients compared with their controls. Breast cancer resistance protein [BCRP] is an example of efflux transporters protecting the enterocytes from toxic compounds. This transporter has been reported to be downregulated in patients with active UC. In newly diagnosed drug-naïve patients with UC, Gutmann *et al.*<sup>28</sup> found that BCRP was significantly reduced compared with control subjects without UC. The same was seen in patients not responding to treatment with 5-ASA or prednisone. These data suggest that the inflammatory process is responsible for the reduced levels of BCRP and that this might be relevant in the pathogenesis of breast cancer.<sup>29</sup>

IBD patients, especially those treated with thiopurines, seem to be at increased risk of developing nonmelanoma skin cancer [NMSC].<sup>30</sup> In meta-analyses of cohort studies and nationwide studies, IBD patients treated with thiopurines also seem to have a 3–5-fold increased risk of developing lymphoproliferative disorders [LDs] compared with the background population, especially three forms of LD: hepato-splenic T-cell lymphoma, Epstein Barr Virus-related post transplant-like LD, and post-mono-nucleosis



**Figure 1.** Cumulative incidences for cancer development and death for ulcerative colitis [UC] patients stratified by gender.



**Figure 2.** Cumulative incidences for cancer development and death for Crohn's disease [CD] patients stratified by gender.

lymphoproliferations.<sup>6,31,32</sup> The overall risk of developing lymphoma, however, is low, and a lack of association in our study may be present because relatively few patients received immunomodulators.

Interestingly, a pooled analysis of data from 1594 CD patients concluded that the incidence of malignancy with the TNF- $\alpha$  inhibitor adalimumab monotherapy was not greater than that of the general population, but co-administration of immunomodulator therapy was associated with a greater than expected incidence of malignancies other than NMSC [SIR = 3.04, 95% CI 1.08–11.06].<sup>33</sup>

A meta-analysis comprising a total of 17052 IBD patients<sup>12</sup> found that IBD patients overall were not at an increased risk of extraintestinal cancer, but CD patients had an increased risk of cancer in the upper gastrointestinal tract, lungs, urinary bladder, and skin [SIR = 2.87, 1.82, 2.03, and 2.35, respectively]. As shown in the Finnish study, UC patients were at an increased risk of liver/biliary cancer [SIR = 2.85] and were at a reduced risk of lung cancer [SIR = 0.39].

For decades, attention has been given to the possible increased risk for haematological malignancies in patients with IBD, especially those who are treated with immunomodulators and/or TNF- $\alpha$  inhibitors. In our study, six UC patients who had never been treated with

immunomodulators/TNF- $\alpha$  inhibitors developed lymphatic leukaemia. None of the CD patients was diagnosed with leukaemia [Tables 2 and 3]. Table 1 shows the cumulative use of immunomodulators/TNF- $\alpha$  inhibitors. The percentage of treated patients is most likely overestimated because most of the patients with missing data are the oldest/dead patients and were often not treated with immunomodulators/TNF- $\alpha$  inhibitors.

Until recently, only anecdotal findings of occurrence of thyroid cancer in UC patients have been published. The Finnish registry study<sup>27</sup> reported SIRs of 2.47 [95% CI 1.18–4.54] and 1.73 [95% CI 1.02–2.72] in male and female UC patients, respectively. In our study, we found two patients with this type of cancer compared with nine in the control group. Since the numbers are small, the interpretation is difficult.

#### 4.1. Strengths and limitations of the present study

Major strengths of the present study include the longitudinal, prospective approach in a well-defined population-based cohort and the accurate detection of cases of malignancies through the national registries. The limited inclusion period and the fact that dedicated specialists and hospitals were responsible for the diagnostic procedures

are factors that contribute to a uniform diagnostic and therapeutic approach. People in Norway do not tend to move house very often, and it is therefore easier to perform prospective long-term studies. All patients had the same duration of follow-up in the identical time period, and each patient was age- and gender-matched with controls from a national registry with very high completeness and quality.

All citizens in Norway are assigned a unique ID number, which enables easy and accurate identification and linkage of medical and death certificate records. There is no reason to assume that inclusion in this study influenced the classification of cancer in the IBD group compared with the background population.

There were some limitations of the study. Even with the high number of patients included in this prospective cohort study, the occurrence of cancers was low which implies a limited statistical power. At diagnosis, the median age of the UC and CD patients was 38 and 29 years, respectively,<sup>14,34</sup> indicating that the patients were relatively young, even 20 years later. Hence, it is to be expected that an extension of the observation period would reveal increased numbers of IBD-related cancers within this group. The levels of education and socioeconomic conditions were not taken into consideration, mainly due to a lack of data. These differences are not marked in Norway. Because the cohort was regularly examined and the control group was not, the probability of being diagnosed with cancer at an earlier stage may have been higher among the patients than among the controls.

## 5. Conclusion

This population-based inception cohort study revealed that patients with a 20-year history of IBD had an increased risk of cancer compared with matched controls from the general population. Most of the excess risk was derived from an increased risk for colorectal cancers in male UC patients. This study also reveals a statistically significant increase in risk for lymphatic leukaemia and breast cancer in UC patients and for trachea/lung cancer and breast cancer in CD patients, compared with the rates seen in the general population, but the numbers here are small.

## Funding

Innlandet Hospital Trust, MSD, Norway and Janssens Biologics B.V.

## Conflict of Interest

The authors have nothing to disclose.

## Acknowledgments

We thank all of the following members of the Inflammatory Bowel South-East Norway [IBSEN] Study Group for participating in this study: Ole Høie, Sørlandet Hospital, Arendal; Gert Huppertz-Hauss and Tomm Bernklev, Telemark Hospital, Skien; Iril Monstad, Oslo University Hospital, Oslo; Jørgen Jahnsen, Akershus University Hospital, Lørenskog; Eva Günther, Østfold Hospital, Moss/Fredrikstad; and Njaal Stray, Diakonhjemmet Hospital, Oslo. The study used data from the Cancer Registry of Norway. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Cancer Registry of Norway is intended, nor should it be inferred.

## Author Contributions

ØH: acquisition of data, analysis and interpretation of data, drafting of the manuscript, revision of the manuscript. MLH: analysis and interpretation of data, critical revision of the manuscript for important intellectual content. MH and

ICS: acquisition of data, analysis and interpretation of data, critical revision of the manuscript for important intellectual content. MCS: statistical analysis, critical revision of the manuscript for important intellectual content. BAM: study concept and design, analysis and interpretation of data, revision of the manuscript for important intellectual content, obtained funding, study supervision.

## References

- Loftus EV Jr, Schoenfeld P, Sandborn WJ. The epidemiology and natural history of Crohn's disease in population-based patient cohorts from North America: a systematic review. *Aliment Pharmacol Ther* 2002;16:51–60.
- Wolters FL, Russel MG, Stockbrugger RW. Has disease outcome in Crohn's disease changed during the last four decades? *Aliment Pharmacol Ther* 2004;20:483–96.
- Kim B, Park SJ, Hong SP, Kim TI, Kim WH, Cheon JH. Proximal disease extension and related predicting factors in ulcerative proctitis. *Scand J Gastroenterol* 2014;49:177–83.
- Molodecky NA, Soon IS, Rabi DM, et al. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 2012;142:46–54.e42; quiz e30.
- Hovde O, Moum BA. Epidemiology and clinical course of Crohn's disease: results from observational studies. *World J Gastroenterol* 2012;18:1723–31.
- Magro F, Peyrin-Biroulet L, Sokol H, et al. Extra-intestinal malignancies in inflammatory bowel disease: results of the 3rd ECCO Pathogenesis Scientific Workshop [III]. *J Crohns Colitis* 2014;8:31–44.
- Canavan C, Abrams KR, Mayberry J. Meta-analysis: colorectal and small bowel cancer risk in patients with Crohn's disease. *Aliment Pharmacol Ther* 2006;23:1097–104.
- Eaden JA, Abrams KR, Mayberry JF. The risk of colorectal cancer in ulcerative colitis: a meta-analysis. *Gut* 2001;48:526–35.
- Jess T, Rungoe C, Peyrin-Biroulet L. Risk of colorectal cancer in patients with ulcerative colitis: a meta-analysis of population-based cohort studies. *Clin Gastroenterol Hepatol* 2012;10:639–45.
- Jess T, Simonsen J, Jorgensen KT, Pedersen BV, Nielsen NM, Frisch M. Decreasing risk of colorectal cancer in patients with inflammatory bowel disease over 30 years. *Gastroenterology* 2012;143:375–81.e1; quiz e13–4.
- Jess T, Gamborg M, Matzen P, Munkholm P, Sorensen TI. Increased risk of intestinal cancer in Crohn's disease: a meta-analysis of population-based cohort studies. *Am J Gastroenterol* 2005;100:2724–9.
- Pedersen N, Duricova D, Elkjaer M, Gamborg M, Munkholm P, Jess T. Risk of extra-intestinal cancer in inflammatory bowel disease: meta-analysis of population-based cohort studies. *Am J Gastroenterol* 2010;105:1480–7.
- Solberg IC, Vatn MH, Hoie O, et al. Clinical course in Crohn's disease: results of a Norwegian population-based ten-year follow-up study. *Clin Gastroenterol Hepatol* 2007;5:1430–8.
- Moum B, Vatn MH, Ekbom A, et al. Incidence of inflammatory bowel disease in southeastern Norway: evaluation of methods after 1 year of registration. Southeastern Norway IBD Study Group of Gastroenterologists. *Digestion* 1995;56:377–81.
- Larsen IK, Smastuen M, Johannesen TB, et al. Data quality at the Cancer Registry of Norway: an overview of comparability, completeness, validity and timeliness. *Eur J Cancer* 2009;45:1218–31.
- Henriksen M, Jahnsen J, Lygren I, et al. Change of diagnosis during the first five years after onset of inflammatory bowel disease: results of a prospective follow-up study [the IBSEN Study]. *Scand J Gastroenterol* 2006;41:1037–43.
- Lennard-Jones JE. Classification of inflammatory bowel disease. *Scand J Gastroenterol Suppl* 1989;170:2–6; discussion 16–9.
- Henriksen M, Jahnsen J, Lygren I, et al. Ulcerative colitis and clinical course: results of a 5-year population-based follow-up study [the IBSEN study]. *Inflamm Bowel Dis* 2006;12:543–50.
- Henriksen M, Jahnsen J, Lygren I, et al. Clinical course in Crohn's disease: results of a five-year population-based follow-up study [the IBSEN study]. *Scand J Gastroenterol* 2007;42:602–10.
- Wilson EB, Hilferty MM. The distribution of chi-square. *Proc Natl Acad Sci U S A* 1931;17:684–8.

21. Hovde O, Kempfski-Monstad I, Smastuen MC, *et al.* Mortality and causes of death in Crohn's disease: results from 20 years of follow-up in the IBSEN study. *Gut* 2014;**63**:771–5.
22. Hovde O, Smastuen MC, Hoivik ML, *et al.* Mortality and causes of death in ulcerative colitis: results from 20 years of follow-up in the IBSEN Study. *Inflamm Bowel Dis* 2016;**22**:141–5.
23. Solberg IC, Lygren I, Jahnsen J, *et al.* Clinical course during the first 10 years of ulcerative colitis: results from a population-based inception cohort [IBSEN Study]. *Scand J Gastroenterol* 2009;**44**:431–40.
24. Hoie O, Wolters FL, Riis *et al.* Low colectomy rates in ulcerative colitis in an unselected European cohort followed for 10 years. *Gastroenterology* 2007;**132**:507–15.
25. Xie J, Itzkowitz SH. Cancer in inflammatory bowel disease. *World J Gastroenterol* 2008;**14**:378–89.
26. Kappelman MD, Farkas DK, Long MD, *et al.* Risk of cancer in patients with inflammatory bowel diseases: a nationwide population-based cohort study with 30 years of follow-up evaluation. *Clin Gastroenterol Hepatol* 2014;**12**:265–73.e1.
27. Jussila A, Virta LJ, Pukkala E, Farkkila MA. Malignancies in patients with inflammatory bowel disease: a nationwide register study in Finland. *Scand J Gastroenterol* 2013;**48**:1405–13.
28. Gutmann H, Hruz P, Zimmermann C, *et al.* Breast cancer resistance protein and P-glycoprotein expression in patients with newly diagnosed and therapy-refractory ulcerative colitis compared with healthy controls. *Digestion* 2008;**78**:154–62.
29. Englund G, Jacobson A, Rorsman F, Artursson P, Kindmark A, Ronnblom A. Efflux transporters in ulcerative colitis: decreased expression of BCRP [ABCG2] and Pgp [ABCB1]. *Inflamm Bowel Dis* 2007;**13**:291–7.
30. Peyrin-Biroulet L, Khosrotehrani K, Carrat F, *et al.* Increased risk for non-melanoma skin cancers in patients who receive thiopurines for inflammatory bowel disease. *Gastroenterology* 2011;**141**:1621–8.e1-5.
31. Beaugerie L, Brousse N, Bouvier AM, *et al.* Lymphoproliferative disorders in patients receiving thiopurines for inflammatory bowel disease: a prospective observational cohort study. *Lancet* 2009;**374**:1617–25.
32. Kandiel A, Fraser AG, Korelitz BI, Brensinger C, Lewis JD. Increased risk of lymphoma among inflammatory bowel disease patients treated with azathioprine and 6-mercaptopurine. *Gut* 2005;**54**:1121–5.
33. Osterman MT, Sandborn WJ, Colombel JF, *et al.* Increased risk of malignancy with adalimumab combination therapy, compared with monotherapy, for Crohn's disease. *Gastroenterology* 2014;**146**:941–9.
34. Moum B, Vatn MH, Ekbo A, *et al.* Incidence of Crohn's disease in four counties in southeastern Norway, 1990–93. A prospective population-based study. The Inflammatory Bowel South-Eastern Norway [IBSEN] Study Group of Gastroenterologists. *Scand J Gastroenterol* 1996;**31**:355–61.