

Table 2. EDE-Q and CIA scores reported by gender and diagnostic categories

	Total Sample ¹		AN ²		BN ³		EDNOS ⁴		ANOVA F		Post Hoc Scheffé
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
EDE-Q global score	4.00 (1.31)	3.48 (1.53)	3.64 (1.47)	3.26 (1.49)	4.16 (1.24)	3.58 (1.41)	4.10 (1.24)	3.49 (1.64)	8.37***	.10	<i>a***, b**</i>
<i>Restraint</i>	3.66 (1.66)	3.28 (1.91)	3.52 (1.82)	3.43 (1.98)	3.84 (1.46)	3.35 (1.54)	3.60 (1.71)	3.21 (2.12)	1.93	-	-
<i>Eating concern</i>	3.50 (1.44)	2.88 (1.69)	3.07 (1.48)	3.11 (1.68)	3.76 (1.39)	3.11 (1.67)	3.56 (1.41)	2.70 (1.74)	11.38***	-	<i>a***, b**</i>
<i>Shape concern</i>	4.71 (1.41)	4.15 (1.49)	4.38 (1.55)	3.82 (1.40)	4.78 (1.39)	4.40 (1.39)	4.88 (1.30)	3.93 (1.64)	12.89***	-	<i>a***, b***</i>
<i>Weight concern</i>	4.13 (1.57)	3.60 (1.67)	3.61 (1.73)	2.66 (1.36)	4.25 (1.53)	3.43 (1.79)	4.37 (1.41)	3.43 (1.79)	12.90***	-	<i>a*, b**</i>
CIA global score	32.57 (10.33)	28.28 (12.95)	32.19 (10.79)	34.60 (6.51)	33.65 (9.46)	26.95 (12.46)	31.90 (10.69)	27.24 (14.28)	1.57	.99	-
<i>Personal impairment</i>	14.01 (4.31)	12.17 (5.61)	13.63 (4.29)	14.86 (2.04)	14.72 (4.26)	12.15 (5.21)	13.67 (4.31)	11.46 (6.33)	-	-	-
<i>Social impairment</i>	9.54 (3.97)	8.89 (5.00)	9.90 (4.27)	11.00 (3.91)	9.50 (3.61)	7.69 (5.02)	9.32 (4.04)	9.92 (5.22)	-	-	-
<i>Cognitive impairment</i>	8.97 (3.81)	7.20 (3.91)	8.61 (4.10)	8.57 (3.41)	9.39 (3.51)	7.11 (3.70)	8.86 (3.81)	6.87 (4.20)	-	-	-

Note. * p<.05, ** p<.01, *** p<.001

Results of post hoc Scheffé tests : *a* = AN vs. BN; *b* = AN vs. EDNOS; *c* = BN vs. EDNOS

¹ = [Female] N(EDE-Q)=620, N(CIA)=553; ¹ = [Male] N(EDE-Q)=47, N(CIA)=46; ² = [Female] N(EDE-Q)=161, N(CIA)=155; ² = [Male] N=7; ³ = [Female] N (EDE-Q)=217, N(CIA)=186; ³ = [Male] N =13; ⁴ = [Female] N(EDE-Q)=242, N(CIA)= 212; ⁴ = [Male] N(EDE-Q)=26, N(CIA) = 26;