

Table 1: Demographics and injury related variables of uncomplicated and complicated MTBIs at 8 weeks post injury

Variables	Uncomplicated MTBI (n=33)	Complicated MTBI (n=29)	p-value
Age (years) mean (SD)	39.9 (12.8)	40.1 (13.6)	0.95
Gender (n, %) - male - female	21 (64) 12 (36)	21 (72) 8 (28)	0.46
Handness (n, %) - right - left	29 (88) 4 (12)	29 (100) 0 (0)	0.12
Education (years) mean (SD)	14.7 (3.7)	14.4 (2.2)	0.65
Employment (n, %) - Yes - No	26 (79) 7 (21)	26 (90) 3 (10)	0.31
Mechanism of injury (n, %) - Traffic accidents - Falls - Violence - Other	19 (58) 8 (24) 4 (12) 2 (6)	7 (24) 17 (59) 4 (14) 1 (3)	0.02
GCS score (n, %) -15 -14 -13	26 (79) 7 (21) 0 (0)	21 (72) 6 (21) 2 (7)	0.42
Isolated TBI (n, %) - yes - no	21 (64) 12 (36)	18 (62) 11 (38)	0.90
LOC (n, %) - yes (<5 min) - yes (>=5 min) - no - unknown	21 (64) 0 (0) 6 (18) 6 (18)	19 (66) 1 (3) 5 (17) 4 (14)	0.87
PTA (n, %) - no amnesia - < 1 h - >1 < 24 h - unknown	1 (3) 23 (70) 0 (0) 9 (27)	4 (14) 19 (66) 0 (0) 6 (20)	0.19
Length of acute hospital stay (days) Median (25%p, 75%p)	1.0 (1.0, 2.0)	2.0 (1.0, 5.5)	0.001
Time between MRI-scan (days) Mean (SD)	468 (126)	447 (138)	0.53

p-values: T-test for continuous variables; Chi square and Fisher's exact test for categorical variables