

Liquidity induced asset bubbles via flows of ELMMs

Francesca Biagini^{*†} Andrea Mazzon^{*} Thilo Meyer-Brandis^{*}

March 23, 2018

Abstract

We consider a constructive model for asset price bubbles, where the market price W is endogenously determined by the trading activity on the market and the fundamental price W^F is exogenously given, as in [32]. To justify W^F from a fundamental point of view, we embed this constructive approach in the martingale theory of bubbles, see [30] and [11], by showing the existence of a flow of equivalent martingale measures for W , under which W^F equals the expectation of the discounted future cash flow.

As an application, we study bubble formation and evolution in a financial network.

Keywords: Bubbles, Equivalent martingale measures, Financial networks, Liquidity based model

1 Introduction

The formation of asset price bubbles has been thoroughly investigated from an economical point of view in many contributions, see Tirole [54], Allen and Gale [3], Choi and Douady [16], [17], Harrison and Kreps [26], Kaizoji [33], Earl et al. [21], DeLong, Shleifer, Summers and Waldmann [20], Scheinkman and Xiong [50], [51], Xiong [59], Abreu and Brunnermeier [1], Föllmer, Horst, and Kirman [23], Miller [39], Zhuk [60].

Different causes have been indicated as triggering factors for bubble birth, such as heterogeneous beliefs between interacting agents (as in [23], [26], [50], [51], [59], [60]), a breakdown of the dynamic stability of the financial system ([16], [17]), the diffusion of new investment decision rules from a few expert investors to larger population of amateurs (see [21]), the tendency of traders to choose the same behavior as the other traders' behavior as thoroughly as possible (see [33]), the presence of short-selling constraints (see [39]).

From the mathematical point of view, one of the main approaches is given by the martingale

^{*}Workgroup Financial and Insurance Mathematics, Department of Mathematics, Ludwig-Maximilians Universität, Theresienstraße 39, 80333 Munich, Germany. Emails: biagini@math.lmu.de, mazzon@math.lmu.de, meyer-brandis@math.lmu.de.

[†]Secondary affiliation: Department of Mathematics, University of Oslo, Box 1053, Blindern, 0316, Oslo, Norway.

theory of bubbles as introduced by Cox and Hobson [19] and Loewenstein and Willard [34] and mainly developed by Jarrow, Protter et al. [27], [28], [29], [30], [31]. See Protter [48] for an overview. In this setting a Q -bubble is defined as the difference between the market price of a given financial asset and its fundamental value, given by the expectation of the future cash flows under an equivalent local martingale measure Q .

Defined in this way, the bubble is a non-negative local martingale under Q , and it is strictly positive if and only if the market wealth W is a strict Q -local martingale (for a complete analysis, see for example [11], [19], [29], [30], [34], [48]).

In a complete market (see [29]), where only one equivalent local martingale measure (ELMM) exists, only two possibilities are given: either no bubble appears at all, or a bubble is already present at the beginning. This is a strong modeling withdraw, therefore in [30] and [11] incomplete markets have been taken into consideration: the birth and the evolution of a bubble are then determined by a flow of different ELMMs that gives rise to a corresponding shifting perception of the fundamental value of the asset. In [30] the underlying pricing measures may change only at certain stopping times, in [11] a continuous flow in the space of martingale measures is considered.

On the other hand, an alternative model is given by Jarrow, Protter and Roch in [32], where the fundamental value is exogenously given, whereas the market value is endogenously determined by the trading activity of investors, and studied through the analysis of the liquidity supply curve. For another constructive model, see also [10].

In this setting a bubble is still defined as the difference between the market value W and the fundamental value W^F , however it does not always coincide with the Q -bubble under a given equivalent martingale measure Q .

A natural question is then if it is possible to embed a constructive model, where the fundamental price is exogenous and the market price endogenous, in the martingale theory of bubbles, by determining a suitable flow of ELMMs for W under which W^F is justified from a fundamental point of view.

More precisely, given a liquidation time T for the financial asset, we look for a flow $(Q^t)_{t \in [0, T]}$ of ELMMs for the market wealth W such that the fundamental value of the asset is given as the expectation of the future cash flow as in equation (3.3). Note however that we do not obtain that W^F is also a (local) martingale under each measure of the flow, as thoroughly discussed in Remark 3.1.

Our main result is then that we can explicitly determine the form of such a flow of ELMMs in a liquidity driven model under very general assumptions, see Theorem 3.16. This requires a consistent technical effort, mostly devoted to guarantee the martingale property of the chosen flows of (eventual) probability densities. In this way we are able to directly connect the impact of the underlying macro-economic factors to the shift of the resulting pricing measure, which may change over time.

As an application of our method, we consider the evolution of a bubble in a financial network and compute the generating flow of ELMMs. However, this example is also of independent interest, as it studies how the interaction of market participants in a financial network can affect asset price formation and the consequent birth of a bubble. Different studies show

how contagion between investors and herding behavior may play an essential role when a bubble grows up: euphoria and exuberance can propagate among market participants, due to exchanges of ideas (see Lux [35]) or to the fact that investors may be attracted by the short period earnings of acquaintances investing in the bubbly asset, as observed by Bayer et al. in [8], where analyzing data from the housing bubble in L. A. in the 2000s the authors notice a strong contagion between neighbors.

Several contributions in the last years has been focusing on how some properties of the network, like mean degree or degree heterogeneity, can influence the contagion of failures and losses between banks during a financial crisis (see for example Acemoglu et al. [2], Allen and Gale [4], Amini et al. [5], Cont et al. [18], Gai and Kapadia [24], Newman et al. [41], Watts [56], Watts and Strogatz [57]). Some investigation has been proposed about how bubbles are generated at the microeconomic level by the interaction of market participants (see among others Lux [35], Scheinkman [50], Scheinkman and Xiong [51], Tirole [54], Zhuk [60]). However, only a few studies have been devoted to understand how the structure of a given financial network can influence the spread of contagion between investors that generates a bubble. In [35], for example, the author models the bubble as caused by a self-organizing process of infection between traders, expressed by a system of PDEs, leading to equilibrium prices that deviate from the fundamental value. However they consider a world in which everybody is connected with everybody, so that the network structure does not enter into play.

In our special case we focus on a model for the signed volume of market orders of X in dependence by some characteristics of the underlying networks of investors, such as the degree distribution. In particular we use some modeling approach deriving from the literature on infectious processes in a population by following the so called SIS model (see Pastor-Satorras and Vespignani [45] and [46]). We provide numerical simulations to investigate how different networks generate different contagion mechanisms and then to bubbles with different evolutions. In particular, it turns out that in more heterogenous networks (i.e. networks with a more right skewed degree distribution) contagion spread faster at the beginning so that the bubble builds up faster and bursts sooner: the nodes with high degree, which in average get infected faster, contribute with an higher weight in the more right skewed distributions.

The paper is therefore organized as follows: in Section 2 we describe the setting of the liquidity model, define the fundamental value of the asset and specify how the trading activity of investors influences the market price of the asset. In Section 3 we determine a possible flow $(Q^t)_{t \in [0, T]}$ of ELMMs satisfying (3.3) and show that the density process $(Z_{t,s})_{s \in [0, T]}$ with $Z_{t,s} = \frac{dQ^t}{dP} |_{\mathcal{F}_s}$ is a true martingale wrt s . In Section 4 we give an example showing how contagion between investors can develop the bubble in a network and compute the generating flow of ELMMs.

2 The Setting

Let (Ω, \mathcal{F}, P) be a probability space and $T > 0$ a random time on it, representing the maturity or liquidation time of the underlying risky asset as in the setting of [30]. We assume that (Ω, \mathcal{F}, P) is endowed with a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ satisfying the usual assumptions of completeness and right continuity.

On $(\Omega, \mathcal{F}, \mathbb{F}, P)$ we have (B^1, B^2, B^3, B^4, N) , where $B^i = (B_t^i)_{t \in [0, T]}$, $i = 1, 2, 3, 4$ are standard \mathbb{F} -Brownian motions and $N_t = \mathbb{1}_{\{\tau \leq t\}}$ is a jump process with τ totally inaccessible stopping time with intensity process $\pi = (\pi_t)_{t \in [0, T]}$. We assume that (B^1, B^2, B^3, B^4, N) are independent processes.

We follow the approach of [32] and study how trading activity may impact prices and generate the formation and bursting of speculative asset price bubbles. We consider a continuous time setting where a stock is traded through a limit order book. The *fundamental wealth or value* of the stock is given as a primitive of the model and represents the price process if market orders have no quantity impact on the price. The *market wealth* equals the fundamental value until market orders are executed. Market orders, which deplete or fill in the limit order book, produce a variation in the price over a small interval of time. If new market orders quickly enter before the price has time to decay again to the fundamental value, these short-term price variations may accumulate and result in a deviation from the fundamental wealth with a consequent bubble birth.

More specifically, we consider a financial asset whose fundamental wealth $W^F = (W_t^F)_{t \in [0, T]}$ (associated to the cumulative dividend process $(D_t)_{t \in [0, T]}$ and to the liquidation value F of the asset at time T) is given by

$$dW_t^F = W_t^F(ad t + b dB_t^1), \quad 0 \leq t \leq T, \quad (2.1)$$

with $W_0^F > 0$, $a \geq 0$ and $b > 0$.

We interpret τ as the time of birth of a bubble for this financial asset. The bubble follows the dynamics

$$d\beta_t = M_t \Lambda_t (-k \beta_t dt + 2dX_t + 2x W_t^F dN_t), \quad 0 \leq t < T, \quad (2.2)$$

where X is the signed volume of market orders (buy market orders minus sell market orders), x is the signed volume of market orders at τ and $M = (M_t)_{t \in [0, T]}$, $\Lambda = (\Lambda_t)_{t \in [0, T]}$ are respectively a measure of illiquidity and the so called *resiliency*. We put $\beta_\tau = 2x \Lambda_\tau M_\tau W_\tau^F$ for a given $x > 0$.

Remark 2.1. *As in [32], we assume that the supply curve for the stock is linear, i.e. that the variation induced by a market order of size y is proportional to y via the stochastic coefficients M and Λ . Here M quantifies the illiquidity level, while the resiliency coefficient Λ represents the degree to which the limit order book recovers from small trades. In this way $\frac{1}{2M_t \Lambda_t}$ gives the depth of the order book at time t , i.e. the size of the order required to change the price of an asset by one unit. This linearity assumption is better justified in the case of frequently traded and large volume stocks, see [12]. For less liquid companies, statistical*

analysis (see for example Cetin et al. [15]) shows that the supply is at best piece-wise linear. For more details about the economical motivation of this setting, we refer to [32].

We consider that X satisfies the following dynamics

$$\begin{aligned} X_t &= 0, & \text{for } 0 \leq t < \tau, \\ dX_t &= \mu_t dt + \sigma_t dB_t^2, & \text{for } \tau \leq t < T, \end{aligned} \quad (2.3)$$

where $\mu = (\mu_t)_{t \in [0, T]}$ and $\sigma = (\sigma_t)_{t \in [0, T]}$ are progressively measurable processes that a priori can also depend on X itself or on the bubble β .

In [32] the signed volume of market orders is modeled as in (2.3) with $\mu \equiv 0$ and $\sigma_t = \alpha \beta_t$. Here we introduce the drift μ in order to see the influence of the network on the size of the bubble, as we specify in Section 4.

Here the fundamental wealth process W^F is exogenously given, while the market wealth process $W = (W_t)_{t \in [0, T]}$ is endogenously determined as

$$W_t = W_t^F + \beta_t, \quad 0 \leq t < T.$$

At liquidation time T we have $W_T = W_T^F$: the asset is liquidated at time T at the estimated firm's value, i.e. at the fundamental value. In particular we require in the sequel that there exists an equivalent local martingale measure for W only on the open interval $[0, T)$, since around time T the liquidation procedure is not subjected to market equilibrium mechanisms.

Assumption 2.2. (i) $\int_{\tau}^T \mu_s^2 ds < \infty$ a.s.

(ii) $\int_{\tau}^T \sigma_s^2 ds < \infty$ a.s. and $\int_{\tau}^T \frac{1}{\sigma_s^4} ds < \infty$ a.s.

(iii) μ and σ are such that there exists a unique solution of (2.3) (see for example Theorem 7 in Chapter V.3 in [47]);

(iv) $M = (M_t)_{t \in [0, T]}$ is an adapted process that satisfies the dynamics

$$dM_t = \tilde{\mu}(M_t)dt + \tilde{\sigma}(M_t)dB_t^3, \quad 0 \leq t \leq T,$$

where $\tilde{\mu}$ and $\tilde{\sigma}$ are such that there exists a unique solution of (2.4) according to Theorem 7 in Chapter V.3 in [47]. Moreover $\int_a^b \tilde{\sigma}^{-4}(x)dx < \infty$ for every a, b such that $0 < a < b < \infty$.

(v) $\Lambda = (\Lambda_t)_{t \in [0, T]}$ satisfies the dynamics

$$d\Lambda_t = \mu'(\Lambda_t)dt + \sigma'(\Lambda_t)dB_t^4, \quad 0 \leq t \leq T,$$

$\Lambda_0 \in (\lambda, 1)$, with μ', σ' that satisfy conditions Theorem 7 in Chapter V.3 in [47]. Furthermore $\mu'(\lambda) > 0$, $\mu'(1) < 0$, $\sigma'(1) = 0$, $\sigma'(\lambda) = 0$ a.s., so that we obtain $\lambda \leq \Lambda_t \leq 1$, a.s. for all $t \in [0, T]$.

(vi) $\pi = (\pi_t)_{t \in [0, T]}$ is bounded, i.e. $|\pi_t| \leq \Pi < \infty$ a.s for all $t \in [0, T]$.

(vii) T is a bounded a.s. (possibly by a very large constant) \mathbb{F} -stopping time independent of (B^1, B^2, N) such that $\tau < T$ a.s.

Notice that we assume $\tau < T$ and T bounded a.s. for the sake of simplicity. The following results still hold without these conditions by imposing some integrability conditions on T . For example, it would be sufficient $T < \infty$ a.s., $\mathbb{E}_P[e^T | \mathcal{F}_t] < \infty$ and $\mathbb{E}_P[T - \tau | \mathcal{F}_t] > 0$ a.s. for $t \in [0, T]$.

Remark 2.3. Here we exclude that σ can depend on β . However the following results also hold for the case $\sigma_t = \alpha \beta_t$, $t \in [\tau, T]$, $\alpha \in \mathbb{R}$, considered in [32] to model the evolution of the bubble given by illiquidity effects. We refer to [36] for more details in this case.

Proposition 2.4. From the hypothesis on M it follows that $\int_0^T M_s^\alpha ds < \infty$ a.s. for all $\alpha \in \mathbb{R}$.

Proof. Following the same argument as in [38], we have that

$$\int_0^T M_s^\alpha ds = \int_0^T \frac{M_s^\alpha}{\tilde{\sigma}^2(M_s)} d[M, M]_s = \int_0^\infty \frac{x^\alpha}{\tilde{\sigma}^2(x)} L_T^x dx, \quad (2.4)$$

where L_T^x is the local time at T and the last equality follows by occupation time formula (see for example Corollary 1 in Chapter IV of [47]).

Then the integral is finite since, by the fact that $0 < M_s < \infty$ a.s. for each $s \in [0, T]$, we have that the occupation time L_T^a has compact support in $(0, \infty)$. \square

From Remark 2.3 we have that β satisfies the SDE

$$d\beta_t = 2\Lambda_t M_t [(-k\beta_t + \mu_t)dt + \sigma_t dB_t^2 + x W_t^F dN_t], \quad \tau \leq t < T.$$

The bubble takes therefore the following explicit expression:

$$\begin{aligned} \beta_t &= \beta_\tau e^{-k \int_\tau^t \Lambda_s M_s ds} + \int_\tau^t \mu_s \Lambda_s M_s e^{-k \int_s^t \Lambda_u M_u du} ds + \\ &+ \int_\tau^t \sigma_s \Lambda_s M_s e^{-k \int_s^t \Lambda_u M_u du} dB_s^2, \quad \tau \leq t < T. \end{aligned} \quad (2.5)$$

3 Flow of equivalent local martingale measures

Let $\mathcal{M}_{loc}(W)$ be the space of equivalent local martingale measures for $W = (W_t)_{t \in [0, T]}$. Given $Q \in \mathcal{M}_{loc}(W)$, a Q -bubble β^Q is defined as

$$\beta_t^Q = W_t - \mathbb{E}_Q[W_T | \mathcal{F}_t] \quad (3.1)$$

in the approach of [29] and [30]. In particular we have that the bubble introduced in (2.2) coincides with a Q -bubble if and only if

$$W_t^F = \mathbb{E}_Q[W_T | \mathcal{F}_t], \quad t \in [0, T] \quad (3.2)$$

for some $Q \in \mathcal{M}_{loc}(W)$.

This is of course not possible in our setting. However we can find a flow $(Q^t)_{t \in [0, T]} \subseteq \mathcal{M}_{loc}(W)$ such that

$$W_t^F = \mathbb{E}_{Q^t}[W_T | \mathcal{F}_t] = \mathbb{E}_{Q^t}[W_T^F | \mathcal{F}_t]. \quad (3.3)$$

In this way the bubble described in (2.2) is the result of the shift in the pricing measure induced by the change in the macro-economic and financial conditions in the market.

Remark 3.1. *We wish to point out the difference and relations between our constructive approach and the martingale theory of bubbles as [29], [30] and [11]. In our setting as well as under the approach of [32], the bubble β is defined as*

$$\beta_t = W_t - W_t^F, \quad (3.4)$$

where W^F is a primitive of the model. According to the martingale theory of bubbles as illustrated in [29] and [30], the market wealth W is given a priori and for a given $Q \in \mathcal{M}_{loc}(W)$ the Q -bubble process β^Q is defined as in (3.1), which also implies that β^Q is non-negative. The two definitions coincide if the fundamental wealth process W^F in (4.4) is also a (local) Q -martingale for $Q \in \mathcal{M}_{loc}(W)$, i.e. if (3.2) holds, otherwise they differ.

In our setting as well as in [32] (see Section 5), we have that $\mathcal{M}_{loc}(W) \cap \mathcal{M}_{loc}(W^F) = \emptyset$, so the bubble process cannot be a local martingale under any equivalent local martingale measure $Q \in \mathcal{M}_{loc}(W)$ for the wealth process W and may also assume negative values. Hence the appearance of negative bubbles is not in contrast with arbitrage theory in our approach.

However, while in the martingale approach the model is automatically arbitrage-free because $\mathcal{M}_{loc}(W) \neq \emptyset$ is assumed a priori, in our “constructive” model for bubbles we need to explicitly exclude arbitrage possibilities. Since in Theorem 3.16 we show the existence of a flow $(Q^t)_{t \in [0, T]} \subseteq \mathcal{M}_{loc}(W)$, i.e. that $\mathcal{M}_{loc}(W) \neq \emptyset$, we obtain that our market model is arbitrage-free, see also Remark 3.17.

It is then a challenging question whether our constructive model can be included in the more fundamental view of the martingale theory of bubble of [29] and [30] by following [11]. To this purpose we investigate the existence of a flow $(Q^t)_{t \in [0, T]} \subseteq \mathcal{M}_{loc}(W)$ which can “fundamentally explain” the a-priori given fundamental wealth, i.e. such that (3.3) holds. This is not in contrast with our comments above since now the measure Q^t is not fixed all over the interval $[0, T]$, but it may change in time. In fact (3.3) does not imply that W^F is a martingale under Q^t over the interval $[0, T]$ because (3.3) holds t -wise and in general it is not true that

$$W_s^F = \mathbb{E}_{Q^t}[W_T | \mathcal{F}_s]$$

for $s \neq t$, $s, t \in [0, T]$.

We now explicitly compute a flow $(Q^t)_{t \in [0, T]} \in \mathcal{M}_{loc}(W)$ justifying the existence of the bubble in (2.2) from a fundamental point of view.

Let $Q \in \mathcal{M}_{loc}(W)$. Then the density process $Z = (Z_t)_{t \in [0, T]}$ of Q with respect to P is given by

$$Z_t = \frac{dQ}{dP} \Big|_{\mathcal{F}_t} = \mathcal{E} \left(\int_0^t \alpha_s^1 dB_s^1 + \int_0^t \alpha_s^2 dB_s^2 + \int_0^t \alpha_s^3 d\tilde{N}_s + \int_0^t \alpha_s^4 dB_s^3 + \int_0^t \alpha_s^5 dB_s^4 + L_t \right),$$

$0 \leq t < T$, where $\tilde{N}_t = N_t - \int_0^{t \wedge \tau} n_s ds$, $t \in [0, T]$, L is a martingale strongly orthogonal to (B^1, B^2, B^3, B^4, N) and the processes α^i , $i = 1, \dots, 5$ are such that for $0 \leq s < T$ the following equality holds:

$$W_s^F (a + b\alpha_s^1) + 2\Lambda_s M_s (\mu_s + \sigma_s \alpha_s^2 - k\beta_s) \mathbb{1}_{\{s \geq \tau\}} + 2\pi_s x W_s^F \Lambda_s M_s (\alpha_s^3 + 1) \mathbb{1}_{\{s < \tau\}} = 0. \quad (3.5)$$

Since (3.5) does not involve α^4 , α^5 or L , we put $\alpha^4 \equiv \alpha^5 \equiv L \equiv 0$.

We can split (3.5) as

$$b\alpha_s^1 = -a - 2\pi_s x \Lambda_s M_s (\alpha_s^3 + 1) \quad \text{for } s < \tau \quad (3.6)$$

and

$$b\alpha_s^1 = -a + \frac{2\Lambda_s M_s}{W_s^F} (k\beta_s - \mu_s - \sigma_s \alpha_s^2) \quad \text{for } s \geq \tau. \quad (3.7)$$

We look for a flow of the form

$$Z_{t,s} = \frac{dQ^t}{dP} \Big|_{\mathcal{F}_s} = \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 + \int_0^s \alpha_u^{t,3} d\tilde{N}_u \right), \quad s \in [0, T], \quad (3.8)$$

since (3.5) does not involve conditions on $\alpha^{t,4}$, $\alpha^{t,5}$ and $\alpha^{t,6}$. In particular, we note that the laws of M , Λ and T are invariant under this change of measure.

If $\alpha^{t,1}$, $\alpha^{t,2}$ and $\alpha^{t,3}$ satisfy (3.6) and (3.7), the fundamental process under Q^t is given by

$$\frac{dW_s^F}{W_s^F} = \tilde{\mu}_s^t ds + b d\tilde{B}_s^t, \quad 0 \leq s \leq T, \quad (3.9)$$

where \tilde{B}^t denote the Q^t -standard Brownian motion given by

$$\tilde{B}_s^t = B_s^1 - \int_0^s \alpha_u^{t,1} du, \quad 0 \leq s \leq T,$$

and

$$\tilde{\mu}_s^t = \begin{cases} -2\pi_s x \Lambda_s M_s (\alpha_s^{t,3} + 1) & \text{for } s < \tau, \\ \frac{2\Lambda_s M_s}{W_s^F} (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) & \text{for } s \geq \tau. \end{cases} \quad (3.10)$$

If the condition

$$\mathbb{E}_{Q^t} \left[\int_t^T (W_s^F)^2 ds \right] < \infty \quad (3.11)$$

is satisfied, we have that (3.3) is equivalent to

$$\mathbb{E}_{Q^t} \left[\int_t^T W_s^F \tilde{\mu}_s^t ds \middle| \mathcal{F}_t \right] = 0,$$

that is

$$0 = \mathbb{E}_{Q^t} \left[\int_t^\tau W_s^F \pi_s x \Lambda_s M_s (\alpha_s^{t,3} + 1) ds + \int_\tau^T \Lambda_s M_s (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) ds \middle| \mathcal{F}_t \right] \quad (3.12)$$

for $t < \tau$ and

$$\mathbb{E}_{Q^t} \left[\int_t^T \Lambda_s M_s (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) ds \middle| \mathcal{F}_t \right] = 0 \quad (3.13)$$

for $t \geq \tau$.

To show the existence of the flow $(Q^t)_{t \in [0, T]} \subseteq \mathcal{M}_{loc}(W)$, we choose $\alpha^{t,2}$ and $\alpha^{t,3}$ so that the integrals inside the conditional expectation in (3.12) and (3.13) are zero almost surely. We show later on that a posteriori this choice ensures as well that (3.11) holds.

For $t \geq \tau$, let

$$\alpha_s^{t,2} = \frac{1}{\Lambda_s M_s \sigma_s} \left(s - \frac{\mathbb{E}[T|\mathcal{F}_t] + t}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2(\mathbb{E}[T|\mathcal{F}_t] - t)} \right) + \frac{k\beta_s}{\sigma_s} - \frac{\mu_s}{\sigma_s}, \quad t \leq s < T.$$

Notice that such $\alpha_s^{t,2}$ is well defined since from Assumption 2.2 it holds $\Lambda_s > 0$, $M_s > 0$, $\sigma_s > 0$ a.s. for every $s \in [0, T]$.

With this choice we have on $\{T > t\}$ that

$$\begin{aligned} & \mathbb{E}_{Q^t} \left[\int_t^T \Lambda_s M_s (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) ds \middle| \mathcal{F}_t \right] \\ &= \mathbb{E}_{Q^t} \left[\int_t^T \left(s - \frac{\mathbb{E}[T|\mathcal{F}_t] + t}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2(\mathbb{E}[T|\mathcal{F}_t] - t)} \right) ds \middle| \mathcal{F}_t \right] \\ &= \mathbb{E}_{Q^t} \left[\left(\frac{T^2 - t^2}{2} - (T - t) \frac{\mathbb{E}[T|\mathcal{F}_t] + t}{2} + (T - t) \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2(\mathbb{E}[T|\mathcal{F}_t] - t)} \right) \middle| \mathcal{F}_t \right] \\ &= \frac{\mathbb{E}[T^2|\mathcal{F}_t] - t^2}{2} - \frac{(\mathbb{E}[T|\mathcal{F}_t] - t)(\mathbb{E}[T|\mathcal{F}_t] + t)}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2} = 0, \end{aligned} \quad (3.14)$$

since by Assumption 2.2 the law of T does not change under Q^t .

For $t < \tau$ define

$$C_{t,\tau} := \int_t^\tau W_s^F \pi_s x \Lambda_s M_s (\alpha_s^{t,3} + 1) ds$$

and choose $\alpha_s^{t,2}$ to be such that

$$\mathbb{E}_{Q^t} \left[\int_\tau^T \Lambda_s M_s (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) ds \middle| \mathcal{F}_t \right] = -\mathbb{E}_{Q^t} [C_{t,\tau} | \mathcal{F}_t],$$

i.e.

$$\alpha_s^{t,2} = \frac{1}{\Lambda_s M_s \sigma_s} \left(s - \frac{\mathbb{E}_{Q^t}[C_{t,\tau}|\mathcal{F}_t]}{\mathbb{E}[T-\tau|\mathcal{F}_t]} - \frac{\mathbb{E}[T+\tau|\mathcal{F}_t]}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2\mathbb{E}[T-\tau|\mathcal{F}_t]} - \frac{\mathbb{E}^2[\tau|\mathcal{F}_t] - \mathbb{E}[\tau^2|\mathcal{F}_t]}{2\mathbb{E}[T-\tau|\mathcal{F}_t]} \right) + \frac{k\beta_s}{\sigma_s} - \frac{\mu_s}{\sigma_s}, \quad t \leq s < T,$$

so that

$$\begin{aligned} & \mathbb{E}_{Q^t} \left[\int_{\tau}^T \Lambda_s M_s (k\beta_s - \mu_s - \sigma_s \alpha_s^{t,2}) ds \middle| \mathcal{F}_t \right] \\ &= \mathbb{E}_{Q^t} \left[\int_{\tau}^T \left(s - \frac{\mathbb{E}_{Q^t}[C_{t,\tau}|\mathcal{F}_t]}{\mathbb{E}[T-\tau|\mathcal{F}_t]} - \frac{\mathbb{E}[T+\tau|\mathcal{F}_t]}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2\mathbb{E}[T-\tau|\mathcal{F}_t]} - \frac{\mathbb{E}^2[\tau|\mathcal{F}_t] - \mathbb{E}[\tau^2|\mathcal{F}_t]}{2\mathbb{E}[T-\tau|\mathcal{F}_t]} \right) ds \middle| \mathcal{F}_t \right] \\ &= \frac{\mathbb{E}[T^2 - \tau^2|\mathcal{F}_t]}{2} - \mathbb{E}_{Q^t}[C_{t,\tau}|\mathcal{F}_t] - \frac{\mathbb{E}[T-\tau|\mathcal{F}_t]\mathbb{E}[T+\tau|\mathcal{F}_t]}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2} \\ & \quad - \frac{\mathbb{E}^2[\tau|\mathcal{F}_t] - \mathbb{E}[\tau^2|\mathcal{F}_t]}{2} \\ &= -\mathbb{E}_{Q^t}[C_{t,\tau}|\mathcal{F}_t], \end{aligned}$$

and then (3.12) holds.

For $s < t \vee \tau$ we set $\alpha_s^{t,2} = 0$.

Summarizing:

$$\alpha_s^{t,2} = \begin{cases} 0 & \text{for } s < \tau \vee t, \\ \frac{1}{\Lambda_s M_s \sigma_s} (s - \eta_{t,\tau}) + \frac{k\beta_s}{\sigma_s} - \frac{\mu_s}{\sigma_s} & \text{for } s \geq \tau \vee t, \end{cases} \quad (3.15)$$

where

$$\begin{aligned} \eta_{t,\tau} &= \frac{\mathbb{E}_{Q^t}[\int_{t \wedge \tau}^{\tau} W_s^F \pi_s x \Lambda_s M_s (\alpha_s^{t,3} + 1) ds | \mathcal{F}_t]}{2\mathbb{E}[T-\tau \vee t | \mathcal{F}_t]} - \frac{\mathbb{E}[T+\tau \vee t | \mathcal{F}_t]}{2} + \frac{\mathbb{E}^2[T|\mathcal{F}_t] - \mathbb{E}[T^2|\mathcal{F}_t]}{2\mathbb{E}[T-\tau \vee t | \mathcal{F}_t]} \\ & \quad - \frac{\mathbb{E}^2[\tau \vee t | \mathcal{F}_t] - \mathbb{E}[(\tau \vee t)^2 | \mathcal{F}_t]}{2\mathbb{E}[T-\tau \vee t | \mathcal{F}_t]}. \end{aligned} \quad (3.16)$$

Remark 3.2. Notice that from Assumption 2.2 and from the fact that the integral in (3.16) is bounded, we have that $\eta_{t,\tau}$ is finite and \mathcal{F}_t -measurable, and that moreover $\mathbb{E}[\eta_{t,\tau}^\alpha] < \infty$ for all $\alpha \in \mathbb{R}$.

Choosing

$$\alpha_s^{t,3} = \begin{cases} 0 & \text{for } s < t \text{ or } s \geq \tau, \\ \frac{1}{(M_s+1)(W_s^F+1)} - 1 & \text{for } t \leq s < \tau, \end{cases} \quad (3.17)$$

and

$$\alpha_s^{t,1} = \begin{cases} 0 & \text{for } s < t, \\ -\frac{a}{b} - \frac{2}{b}\pi_s\Lambda_s\frac{M_s}{M_s+1}\frac{1}{W_s^F+1} & \text{for } t \leq s < \tau, \\ -\frac{a}{b} - \frac{2}{bW_s^F}(s - \eta_{t,\tau}) & \text{for } s \geq \tau \vee t. \end{cases} \quad (3.18)$$

we have that (3.12) and (3.13) hold.

Now we give the following

Proposition 3.3. *Let $\alpha^{t,1}$, $\alpha^{t,2}$ and $\alpha^{t,3}$ be as in (3.15)-(3.18). Then*

$$\mathbb{E}_{Q^t} \left[\int_t^T (W_s^F)^2 ds \right] < \infty, \quad t \in [0, T].$$

Proof. From (3.10) and from the expressions of $\alpha^{t,1}$, $\alpha^{t,2}$ and $\alpha^{t,3}$ in (3.15)-(3.18) we have that

$$\tilde{\mu}_s^t = \begin{cases} -2\pi_s x \Lambda_s \frac{M_s}{M_s+1} \frac{1}{W_s^F+1} & \text{for } s < \tau, \\ \frac{1}{W_s^F}(\eta_{t,\tau} - s) & \text{for } s \geq \tau, \end{cases}$$

where $\eta_{t,\tau}$ is given in (3.16). Then from (3.9) it holds that under Q^t

$$\begin{aligned} dW_s^F &= \psi_s ds + bW_s^F d\tilde{B}_s^t && \text{for } s < \tau, \\ dW_s^F &= (\eta_{t,\tau} - s) ds + bW_s^F d\tilde{B}_s^t && \text{for } s \geq \tau, \end{aligned}$$

where $\psi_s = -2\pi_s x \Lambda_s \frac{M_s}{M_s+1} \frac{1}{W_s^F+1}$.

Thus we have

$$W_s^F = \begin{cases} e^{b\tilde{B}_s^t - \frac{b^2}{2}s} \int_0^s \psi_u e^{-b\tilde{B}_u^t + \frac{b^2}{2}u} du & \text{for } s < \tau, \\ e^{b\tilde{B}_s^t - \frac{b^2}{2}s} \int_0^s (\eta_{t,\tau} - u) e^{-b\tilde{B}_u^t + \frac{b^2}{2}u} du & \text{for } s \geq \tau. \end{cases}$$

Then

$$\begin{aligned} & \mathbb{E}_{Q^t} \left[\int_t^T (W_s^F)^2 ds \right] \\ &= \mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^{\tau} \left(\int_0^s \psi_u e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^2 ds + \int_{\tau}^T \left(\int_0^s (\eta_{t,\tau} - u) e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^2 ds \right] \\ &\leq \mathbb{E}_{Q^t} \left[4\Pi^2 x^2 \int_{t \wedge \tau}^{\tau} \left(\int_0^s e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^2 ds + (|\eta_{t,\tau}| + T)^2 \int_{\tau}^T \left(\int_0^s e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^2 ds \right] \\ &\leq \left(4\Pi^2 x^2 + \mathbb{E} [(|\eta_{t,\tau}| + T)^4]^{\frac{1}{2}} \right) \mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^T \left(\int_0^s e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^4 ds \right]^{\frac{1}{2}}. \end{aligned}$$

Since T is bounded and the first term is finite by Remark 3.2, it remains to prove

$$\mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^T \left(\int_0^s e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^4 ds \right] < \infty. \quad (3.19)$$

We have that

$$\begin{aligned} \mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^T \left(\int_0^s e^{b(\tilde{B}_s^t - \tilde{B}_u^t) - \frac{b^2}{2}(s-u)} du \right)^4 ds \right] &= \mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^T \left(\int_0^s e^{b\tilde{B}_{s-u}^t - \frac{b^2}{2}(s-u)} du \right)^4 ds \right] \\ &= \mathbb{E}_{Q^t} \left[\int_{t \wedge \tau}^T \left(\int_0^s e^{b\tilde{B}_r^t - \frac{b^2}{2}r} dr \right)^4 ds \right] \leq \mathbb{E}_{Q^t} [(T - t \wedge \tau)^2]^{\frac{1}{2}} \mathbb{E}_{Q^t} \left[\left(\int_0^T e^{b\tilde{B}_r^t - \frac{b^2}{2}r} dr \right)^8 \right]^{\frac{1}{2}}. \end{aligned}$$

The first term is finite by Assumption 2.2 on T and τ , whereas

$$\mathbb{E}_{Q^t} \left[\left(\int_0^T e^{b\tilde{B}_r^t - \frac{b^2}{2}r} dr \right)^8 \right] \leq \mathbb{E}_{Q^t} \left[\int_0^T e^{8b\tilde{B}_r^t - 4b^2r} dr \right] = \int_0^T \mathbb{E}_{Q^t} \left[e^{8b\tilde{B}_r^t - 4b^2r} \right] dr < \infty.$$

Then (3.19) holds and we have the result. \square

We have therefore proved that, if we take $\alpha^{t,1}$, $\alpha^{t,2}$ and $\alpha^{t,3}$ as in (3.15)-(3.18), then (3.6), (3.7) and (3.3) are satisfied.

From now on we denote $Z_{t,s} := \frac{dQ^t}{dP} |_{\mathcal{F}_s}$ for all $s \geq t$, and $Z_{t,s} = 1$ for $s < t$.

Note that we have not yet used the hypothesis on μ and σ of Assumption 2.2 to derive (3.8). From now on we will need them to prove that $(Z_{t,s})_{s \in [t, T]}$ is a true martingale for each $t \in [0, T]$, i.e. that each Q^t , $t \in [0, T]$, in (3.8) belongs to $\mathcal{M}_{loc}(W)$.

Remark 3.4. *By Assumption 2.2, as proved in Proposition 2.4, we exclude that the integral $\int_0^\cdot M_s^2 ds$ can explode in finite time. This is a difference with respect to [32], where the bubble bursts (i.e. $\beta_t = 0$) at $\inf\{s \mid \int_0^s M_u^2 du = +\infty\}$.*

In our model, however, the bubble can be zero, and also negative, even if the liquidity is not zero: by (2.5) it can be seen that this can happen when the drift μ of the signed volume of market orders becomes negative. In this approach, therefore, whether or not the bubble is positive depends more on the attitude of the investors than on the liquidity. In Section 4 we propose an example to show how contagion between traders in financial networks can determine the value of μ .

From now on, we fix $t \in [0, T]$. We begin the analysis by noticing that, since $[B^1, N] \equiv [B^2, N] \equiv 0$,

$$\begin{aligned} Z_{t,s} &= \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 + \int_0^s \alpha_u^{t,3} d\tilde{N}_u \right) \\ &= \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 \right) \mathcal{E} \left(\int_0^s \alpha_u^{t,3} d\tilde{N}_s \right) \end{aligned}$$

for $s \in [0, T)$.

Moreover

$$\begin{aligned} \mathcal{E} \left(\int_0^s \alpha_u^{t,3} d\tilde{N}_u \right) &\leq \exp \left\{ \int_0^s \left[\alpha_u^{t,3} - \frac{1}{2} (\alpha_u^{t,3})^2 \right] dN_u - \int_0^s \alpha_u^{t,3} \pi_u du \right\} \\ &\quad \cdot \prod_{0 \leq u \leq s} (1 + \Delta(\alpha_u^{t,3} N_u)) \exp \left\{ \Delta(\alpha_u^{t,3} N_u) + \frac{1}{2} \Delta(\alpha_u^{t,3} N_u)^2 \right\} \\ &\leq 2 \exp \left\{ \frac{3}{2} + \int_0^s \left[|\alpha_u^{t,3}| + \frac{1}{2} |\alpha_u^{t,3}|^2 \right] dN_u + \int_0^s |\alpha_u^{t,3}| \pi_u du \right\} \\ &\leq 2e^{3+T\Pi}, \end{aligned}$$

since by (3.17) it holds $|\alpha_s^{t,3}| \leq 1$.

Then, taking $(\bar{Z}_{t,s})_{s \in [0, T)}$ with

$$\bar{Z}_{t,s} = \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 \right)$$

we have

$$Z_{t,s} \leq 2e^{3+T\Pi} \bar{Z}_{t,s}. \quad (3.20)$$

We give the following

Lemma 3.5. *Let X, Y be two positive stochastic processes such that $Y_t \leq X_t$ a.s. $\forall t \geq 0$, and let X be of class DL^1 . Then Y is of class DL as well.*

Proof. By Theorem 11 of Chapter I of [47] we have that a family of random variables $(U_\alpha)_{\alpha \in A}$ is uniformly integrable if and only if there exists a function G defined on $[0, \infty)$, positive, increasing and convex, such that $\lim_{x \rightarrow \infty} \frac{G(x)}{x} = +\infty$ and $\sup_\alpha \mathbb{E}[G \circ |U_\alpha|] < \infty$. Fix now $t \geq 0$, and call $J_t = \{\tau : \tau \leq t \text{ stopping time}\}$, $U_X^t = \{X_\tau : \tau \in J_t\}$ and $U_Y^t = \{Y_\tau : \tau \in J_t\}$. Since by hypothesis U_X^t is uniformly integrable, there exists a function G that satisfies the properties stated before. We have that

$$G(Y_\tau) \leq G(X_\tau), \quad a.s. \text{ for } \tau \in J_t,$$

and then that

$$\mathbb{E}[G(Y_\tau)] \leq \mathbb{E}[G(X_\tau)], \quad \tau \in J_t.$$

Thus

$$\sup_{\tau \in J_t} \mathbb{E}[G(Y_\tau)] \leq \sup_{\tau \in J_t} \mathbb{E}[G(X_\tau)] < \infty.$$

Therefore U_Y^t is uniformly integrable and Y is of class DL . \square

We have then the following

¹A stochastic process X is of class DL if, for each $t \geq 0$, $\{X_\tau : \tau \leq t \text{ stopping time}\}$ is uniformly integrable.

Proposition 3.6. $(Z_{t,s})_{s \in [0,T]}$ in (3.8) is a martingale if $(\bar{Z}_{t,s})_{s \in [0,T]}$ is a martingale.

Proof. Since a local martingale is a true martingale if and only if it is of class DL , see Proposition 1.7 of Chapter IV of [49], we have that if \bar{Z} is a true martingale then $2e^{3+T\Pi}\bar{Z}$, being a martingale as well, is of class DL . Thus, by Lemma 3.5 and by (3.20), Z is of class DL , and therefore by Proposition 1.7 of Chapter IV of [49] it is a true martingale. \square

To prove that \bar{Z} is a martingale we rely on some results provided by Mijatovic and Urusov [37] and by Wong and Heyde [58]. We first need some preliminaries.

Consider the state space $J = (l, r)$, $-\infty \leq l < r \leq \infty$ and a J -valued diffusion $Y = (Y_s)_{s \in [0,T]}$ on some filtered probability space, governed by the SDE

$$dY_s = \mu_Y(Y_s)ds + \sigma_Y(Y_s)dB_s, \quad 0 \leq s < T, \quad (3.21)$$

with $Y_0 = x_0 \in J$, W Brownian motion and deterministic functions $\mu_Y(\cdot)$ and $\sigma_Y(\cdot)$, that from now on we will simply denote by μ_Y and σ_Y , such that

$$\sigma_Y(x) \neq 0 \quad \forall x \in J \quad (3.22)$$

and

$$\frac{1}{\sigma_Y^2}, \frac{\mu_Y}{\sigma_Y^2} \in L_{loc}^1(J), \quad (3.23)$$

where $L_{loc}^1(J)$ denotes the class of locally integrable functions on J , i.e. the measurable functions $(J, \mathcal{B}(J)) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ that are integrable on compact subsets of J .

Consider the stochastic exponential

$$\mathcal{E} \left(\int_0^s f(Y_u)dB_u \right), \quad 0 \leq s < T, \quad (3.24)$$

with $f(\cdot)$ such that

$$\frac{f^2}{\sigma_Y^2} \in L_{loc}^1(J) \quad (3.25)$$

and the auxiliary J -valued diffusion \tilde{Y} governed by the SDE

$$d\tilde{Y}_s = \left(\mu_Y(\tilde{Y}_s) + f(\tilde{Y}_s)\sigma_Y(\tilde{Y}_s) \right) ds + \sigma_Y(\tilde{Y}_s)d\tilde{B}_s, \quad 0 \leq s < T, \quad (3.26)$$

where \tilde{B} is a Brownian motion on some probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$.

Put $\bar{J} = [l, r]$ and, fixing an arbitrary $c \in J$, define

$$\rho(x) := \exp \left\{ - \int_c^x \frac{2\mu_Y}{\sigma_Y^2}(y)dy \right\}, \quad x \in J, \quad (3.27)$$

$$\tilde{\rho}(x) := \rho(x) \exp \left\{ - \int_c^x \frac{2f}{\sigma_Y}(y)dy \right\}, \quad x \in J, \quad (3.28)$$

$$s(x) := \int_c^x \rho(y)dy, \quad x \in \bar{J}, \quad (3.29)$$

$$\tilde{s}(x) := \int_c^x \tilde{\rho}(y)dy, \quad x \in \bar{J}. \quad (3.30)$$

Denote $\rho = \rho(\cdot)$, $s = s(\cdot)$, $s(r) = \lim_{x \rightarrow r^-} s(x)$, $s(l) = \lim_{x \rightarrow l^+} s(x)$, and analogously for $\tilde{s}(\cdot)$ and $\tilde{\rho}(\cdot)$.

Recall that by Feller's test for explosions \tilde{Y} exits its state space with positive probability at the boundary point r if and only if

$$\tilde{s}(r) < \infty \quad \text{and} \quad \frac{\tilde{s}(r) - \tilde{s}}{\tilde{\rho}\sigma_Y^2} \in L_{loc}^1(r-), \quad (3.31)$$

where $L_{loc}^1(r-) := \{g|g : (J, \mathcal{B}(J)) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R})) \text{ such that } \int_x^r g(y)dy < \infty \text{ for some } x \in J\}$. Similarly, \tilde{Y} exits its state space with positive probability at the boundary point l if and only if

$$\tilde{s}(l) > -\infty \quad \text{and} \quad \frac{\tilde{s} - \tilde{s}(l)}{\tilde{\rho}\sigma_Y^2} \in L_{loc}^1(l+), \quad (3.32)$$

where $L_{loc}^1(l+) := \{g|g : (J, \mathcal{B}(J)) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R})) \text{ such that } \int_l^x g(y)dy < \infty \text{ for some } x \in J\}$. Moreover, the endpoint r of J is said to be *good* if

$$s(r) < \infty \quad \text{and} \quad \frac{(s(r) - s)f^2}{\rho\sigma_Y^2} \in L_{loc}^1(r-), \quad (3.33)$$

or equivalently (see [37]) if

$$\tilde{s}(r) < \infty \quad \text{and} \quad \frac{(\tilde{s}(r) - \tilde{s})f^2}{\tilde{\rho}\sigma_Y^2} \in L_{loc}^1(r-). \quad (3.34)$$

Similarly, the endpoint l of J is said to be *good* if

$$s(l) > -\infty \quad \text{and} \quad \frac{(s - s(l))f^2}{\rho\sigma_Y^2} \in L_{loc}^1(l+), \quad (3.35)$$

or equivalently if

$$\tilde{s}(l) > -\infty \quad \text{and} \quad \frac{(\tilde{s} - \tilde{s}(l))f^2}{\tilde{\rho}\sigma_Y^2} \in L_{loc}^1(l+). \quad (3.36)$$

We recall here Theorem 2.1 in [37].

Theorem 3.7. *Let the functions μ_Y , σ_Y , and f satisfy conditions (3.22), (3.23) and (3.25), and let Y be a solution of the SDE (3.21).*

Then the Doléans exponential given by (3.24) is a martingale for any $T < \infty$ if and only if both of the following requirements are satisfied:

- (a) *condition (3.31) does not hold or conditions (3.33)-(3.34) hold;*
- (b) *condition (3.32) does not hold or conditions (3.35)-(3.36) hold.*

We now obtain the following

Proposition 3.8. Let $S = (S_s)_{s \in [0, T]}$ be a geometric Brownian motion

$$dS_s = \mu_0 S_s ds + \sigma_0 S_s dB_s, \quad 0 \leq s < T, \quad (3.37)$$

where B is a Brownian motion, $\mu_0 \in \mathbb{R}$ and $\sigma_0 > 0$.

Then the process

$$\mathcal{E} \left(\int_0^s (S_u)^{-1} dB_u \right), \quad 0 \leq s < T,$$

is a martingale.

Proof. We show that the requirements of Theorem 3.7 hold for $Y = S$, with $\mu_Y(x) = \mu_0 x$, $\sigma_Y(x) = \sigma_0 x$ and $f(x) = x^{-1}$. Notice that μ_Y , σ_Y and f satisfy conditions (3.22), (3.23) and (3.25) with $J = (0, \infty)$. Then, taking $c = 1$ for the functions (3.27)-(3.30) and first assuming $\frac{2\mu_0}{\sigma_0^2} \neq 1$, we have

$$\rho(x) = \exp \left\{ - \int_1^x \frac{2\mu_Y(y)}{\sigma_Y^2(y)} dy \right\} = x^{-\frac{2\mu_0}{\sigma_0^2}}, \quad (3.38)$$

$$\tilde{\rho}(x) = \rho(x) \exp \left\{ - \int_1^x \frac{2f(y)}{\sigma_Y(y)} dy \right\} = x^{-\frac{2\mu_0}{\sigma_0^2}} \exp \left(\frac{2}{\sigma_0} \left(\frac{1}{x} - 1 \right) \right), \quad (3.39)$$

$$s(x) = \int_1^x \rho(y) dy = \frac{\sigma_0^2}{2\mu_0 - \sigma_0^2} (1 - x^{-\gamma_0}), \quad (3.40)$$

$$\tilde{s}(x) = \int_1^x \tilde{\rho}(y) dy = e^{-\frac{2}{\sigma_0}} \left(-\frac{2}{\sigma_0} \right)^{-\gamma_0} \left[\bar{\Gamma} \left(\gamma_0, -\frac{2}{x\sigma_0} \right) - \bar{\Gamma} \left(\gamma_0, -\frac{2}{\sigma_0} \right) \right], \quad (3.41)$$

with $\gamma_0 = \frac{2\mu_0}{\sigma_0^2} - 1$ and where $\bar{\Gamma}(a, z) = \int_z^\infty e^{-t} t^{a-1} dt$, $a \in \mathbb{R}^+$, $z \in \mathbb{R}$, is the incomplete Gamma function extended to all \mathbb{R} .

Notice that in (3.41) we have that

$$\begin{aligned} \tilde{s}(x) &= e^{-\frac{2}{\sigma_0}} \left(-\frac{2}{\sigma_0} \right)^{-\gamma_0} \left[\bar{\Gamma} \left(\gamma_0, -\frac{2}{x\sigma_0} \right) - \bar{\Gamma} \left(\gamma_0, -\frac{2}{\sigma_0} \right) \right] \\ &= e^{-\frac{2}{\sigma_0}} \left(\frac{2}{\sigma_0} \right)^{-\gamma_0} (-1)^{-\gamma_0} \int_{-\frac{2}{x\sigma_0}}^{-\frac{2}{\sigma_0}} e^{-t} (-1)^{\gamma_0-1} |t|^{\gamma_0-1} dt \\ &= -e^{-\frac{2}{\sigma_0}} \left(\frac{2}{\sigma_0} \right)^{-\gamma_0} \int_{-\frac{2}{x\sigma_0}}^{-\frac{2}{\sigma_0}} e^{-t} |t|^{\gamma_0-1} dt \in \mathbb{R}. \end{aligned} \quad (3.42)$$

We obtain that:

- in $l = 0$ we have

$$\tilde{s}(0) = -e^{-\frac{2}{\sigma_0}} \left(\frac{2}{\sigma_0} \right)^{-\gamma_0} \int_{-\infty}^{-\frac{2}{\sigma_0}} e^{-t} |t|^{\gamma_0-1} dt = -\infty,$$

thus condition (3.32) does not hold and the first requirement of (b) in Theorem 3.7 is fulfilled;

- if $\gamma_0 < 0$ we have

$$\tilde{s}(\infty) = e^{-\frac{2}{\sigma_0}} \left(\frac{2}{\sigma_0}\right)^{-\gamma_0} \int_{-\frac{2}{\sigma_0}}^0 e^{-t}|t|^{\gamma_0-1} dt = \infty$$

then condition (3.31) does not hold and the first requirement of (a) in Theorem 3.7 is fulfilled;

- if $\gamma_0 > 0$ then $s(\infty) = \frac{\sigma_0^2}{2\mu_0 - \sigma_0^2} = C < \infty$, and condition (3.33) holds since

$$\frac{s(r) - s}{\rho\sigma_0^2} = C \frac{x^{-\gamma_0} x^{\frac{2\mu_0}{\sigma_0^2}}}{x^4} = \frac{1}{x^3}.$$

Therefore the second requirement of (a) in Theorem 3.7 is fulfilled.

So we have that if $\gamma_0 \neq 0$ the requirements of Theorem 3.7 are satisfied, and thus Z is a martingale.

In the case $\gamma_0 = 0$, i.e. $\mu_0 = \frac{\sigma_0^2}{2}$, we have that the process $S = (S_u)_{u \in [0, T]}$ in (3.37) takes the form $S_u = e^{\sigma_0 B_u}$, $0 \leq u < T$. We can thus apply the results of Theorem 3.7 taking $J = (-\infty, \infty)$, $\mu_Y \equiv 0$, $\sigma_Y \equiv 1$, $f(x) = e^{-\sigma_0 x}$ and $c = 0$ in (3.27)-(3.30). We have

$$\begin{aligned} \rho(x) &= \exp \left\{ - \int_0^x \frac{2\mu_Y}{\sigma_Y^2}(y) dy \right\} = 1, \\ \tilde{\rho}(x) &= \rho(x) \exp \left\{ - \int_0^x \frac{2f}{\sigma_Y}(y) dy \right\} = \exp(2(e^{-\sigma_0 x} - 1)/\sigma_0), \\ s(x) &= \int_0^x \rho(y) dy = x \\ \tilde{s}(x) &= \int_0^x \tilde{\rho}(y) dy = \frac{1}{\sigma_0} e^{-\frac{2}{\sigma_0}} (Ei(2/\sigma_0) - Ei(2e^{-\sigma_0 x}/\sigma_0)), \end{aligned} \quad (3.43)$$

where $Ei(z) = - \int_{-z}^{\infty} \frac{e^{-u}}{u} du$ is the exponential integral function that satisfies $\lim_{z \rightarrow \infty} Ei(z) = \infty$ and $\lim_{z \rightarrow 0} Ei(z) = -\infty$. Therefore $\tilde{s}(\infty) = \infty$ and $\tilde{s}(-\infty) = -\infty$, then the first requirements of (a) and (b) of Theorem 3.7 are both satisfied and Z is a martingale. \square
Then we have immediately

Corollary 3.9. *Under Assumptions 2.2, the process*

$$\mathcal{E} \left(\int_{\tau}^s \frac{1}{W_u^F} dB_u^1 \right), \quad \tau \leq s < T, \quad (3.44)$$

is a martingale for every fixed $T < \infty$.

To prove that Corollary 3.9 also implies that $\mathcal{E}\left(\int_{\tau}^s \alpha_u^{t,1} dB_u^1\right)$ is a martingale, we extend the results of Wong and Heyde in [58].

To this purpose we consider a \mathbb{F} -progressively measurable d -dimensional process $H = (H_s)_{s \in [0, T]}$ of the form

$$H_s = \xi(B(\cdot), s)\zeta_s + \eta_s, \quad (3.45)$$

where $\xi \in C_0(\mathbb{R}^{d+1}, \mathbb{R}^d)$, B is a d -dimensional progressively measurable Brownian motion and ζ, η are d -dimensional stochastic processes independent of B . Here the product between ξ and ζ is intended componentwise.

Define

$$\tau_N^{M_H} = \inf \left(s \in [0, T) : M_H(t) := \int_0^t \|H_u\|^2 du \geq N \right),$$

with the convention that $\inf \emptyset = \infty$, and then

$$\tau^{M_H} = \lim_{N \rightarrow \infty} \tau_N^{M_H}. \quad (3.46)$$

Then we have the following

Proposition 3.10. *Let H be as in (3.45), and defined up to the explosion time τ^{M_H} in (3.46). Then there also exists a d -dimensional \mathbb{F} -progressively measurable process, $Y = (Y_s)_{s \in [0, T]}$ with $Y_s = \xi(W(\cdot) + \int_0^s Y_u du, s)\zeta_s + \eta_s$ defined up to the explosion time τ^{M_Y} with*

$$\tau^{M_Y} = \lim_{N \rightarrow \infty} \tau_N^{M_Y},$$

where

$$\tau_N^{M_Y} = \inf \left(s \geq 0 : M_Y(s) := \int_0^s \|Y_u\|^2 du \geq N \right) \wedge T,$$

such that the stochastic exponential $Z^H = (Z_s^H)_{s \in [0, T]}$ with $Z_s^H = \mathcal{E}\left(\int_0^s H_u dW_u\right)$ satisfies

$$P(\tau^{M_Y} > T) = \mathbb{E}[Z_T^H].$$

Hence Z^H is a (true) martingale if and only if $P(\tau^{M_Y} > T) = 1$.

Proof. Since the proof is a long but easy extension of the result in [58], we omit it here and refer to [36]. \square

Proposition 3.11. *In the setting of Section 2, the process*

$$\mathcal{E}\left(\int_0^s |\alpha_u^{t,1}| dB_u^1\right), \quad 0 \leq s < T,$$

with $\alpha^{t,1}$ in (3.18) is a martingale for each $t \in [0, T]$.

Proof. For $s < \tau$ we have

$$|\alpha_s^{t,1}| = \frac{a}{b} + \frac{2}{b}\pi_s\Lambda_s \frac{M_s}{M_s+1} \frac{1}{W_s^F+1} \leq \frac{a}{b} + \frac{2}{b}\Pi,$$

then $\mathcal{E}\left(\int_0^\cdot |\alpha_u^{t,1}| dB_u^1\right)$ is a martingale up to time τ since it satisfies Novikov condition since

$$\mathbb{E}\left[\exp\left(\int_0^\tau (\alpha_s^{t,1})^2 ds\right)\right] \leq \mathbb{E}[\exp(c^2\tau)]$$

with $c = \frac{a}{b} + \frac{2}{b}\Pi$.

Consider now $s \geq \tau$. We have that the process Y associated to $|\alpha_s^{t,1}|$ as in Proposition 3.10 satisfies

$$Y_s = \frac{2}{bW_s^F}(s + |\eta_{t,\tau}|) \exp\left(-b \int_{t \wedge \tau}^s Y_u du\right), \quad t \wedge \tau \leq s < T,$$

with $\eta_{t,\tau}$ in (3.16). On the other hand, we have

$$\tilde{Y}_s = \frac{1}{W_s^F} \exp\left(-b \int_{t \wedge \tau}^s \tilde{Y}_u du\right), \quad t \wedge \tau \leq s < T,$$

where \tilde{Y} is the process associated to $\frac{1}{W^F}$. By Corollary 3.9 and Proposition 3.10 it holds

$$\int_{t \wedge \tau}^T \tilde{Y}_s^2 < \infty. \quad (3.47)$$

We want to see that the integral of Y^2 does not explode as well.

We have that

$$\Delta_s = \frac{\tilde{Y}_s}{Y_s} = \frac{b}{s + |\eta_{t,\tau}|} \cdot \exp\left(b \int_{t \wedge \tau}^s (Y_u - \tilde{Y}_u) du\right), \quad t \wedge \tau \leq s < T. \quad (3.48)$$

Define the stopping time

$$\tau_1 = \inf\{s \in [t \wedge \tau, T) : \Delta_s \leq 1\} \wedge T$$

and notice that, since Y and \tilde{Y} are continuous, $\Delta_{\tau_1} = 1$.

Define

$$\tau_2 = \inf\{s \geq \tau_1 : \Delta_s \geq 1\} \wedge T,$$

If $\tau_1 = T$, we are done. Otherwise consider $s \in (\tau_1, \tau_2)$.

Since for $\tau_1 < s < \tau_2$ we have

$$\Delta_s = \frac{b}{s + |\eta_{t,\tau}|} \cdot \exp\left(b \int_{t \wedge \tau}^s (Y_u - \tilde{Y}_u) du\right) \geq \frac{b}{T + |\eta_{t,\tau}|} \exp\left(b \int_{t \wedge \tau}^{\tau_1} (Y_u - \tilde{Y}_u) du\right),$$

it follows

$$Y_s \leq \frac{\tilde{Y}_s(T + |\eta_{t,\tau}|)}{b} \exp\left(b \int_{t \wedge \tau}^{\tau_1} (\tilde{Y}_u - Y_u) du\right) \leq \frac{\tilde{Y}_s(T + |\eta_{t,\tau}|)}{b} \exp\left(b \int_{t \wedge \tau}^{\tau_1} \tilde{Y}_u du\right)$$

for $\tau_1 < s < \tau_2$, which implies, together with (3.47), that $M_Y(s) := \int_{t \wedge \tau}^s Y_s^2 ds$ does not explode before τ_2 .

But after τ_2 , up to $\tau_3 = \inf\{s \geq \tau_2 : \Delta_s \leq 1\} \wedge T$, Y is smaller than \tilde{Y} , hence $M_Y(s) \leq M_{\tilde{Y}_s}$ on $[\tau_2, \tau_3]$.

Repeating this argument up to T , we obtain that $\mathcal{E}\left(\int_0^s |\alpha_u^{t,1}| dB_u^1\right)$ is a martingale by Proposition 3.10. \square

We want now to prove that

$$\mathcal{E}\left(\int_0^s |\alpha_u^{t,2}| dB_u^2\right), \quad 0 \leq s < T, \quad (3.49)$$

with $\alpha^{t,2}$ in (3.15) is a martingale as well.

We start with the following

Proposition 3.12. *Let β be the bubble as in (2.5). Under Assumption 2.2, the Doléans exponential*

$$\mathcal{E}\left(\int_0^s \beta_u dB_u^2\right), \quad 0 \leq s < T,$$

is a martingale.

Proof. If we rewrite β in the form (3.45), we obtain that

$$\xi(B^2(\cdot), s) = \int_{\tau}^s \sigma_u \Lambda_u M_u e^{-k \int_u^s k \Lambda_r M_r dr} dB_u^2, \quad \tau \leq s < T,$$

i.e. the process Y associated to β in Proposition 3.10 is given by

$$\begin{aligned} Y_s &= \beta_{\tau} e^{\int_{\tau}^s (-k + \sigma_u) \Lambda_u M_u ds} + \int_{\tau}^s \mu_u \Lambda_u M_u e^{\int_u^s (-k + \sigma_r) \Lambda_r M_r dr} du \\ &\quad + \int_{\tau}^s \sigma_u \Lambda_u M_u e^{\int_u^s (-k + \sigma_r) \Lambda_r M_r dr} dB_u^2, \quad \tau \leq s < T. \end{aligned} \quad (3.50)$$

We first prove that $Y_s < \infty$ for each $s \in [\tau, T)$. We have $\int_u^s (-k + \sigma_r) \Lambda_r M_r dr < \infty$ a.s. for each $s \in [\tau, T)$ by the hypothesis on σ and Λ in Assumption 2.2 and by Proposition 2.4.

Thus by Theorem 2.4 of [38] and by the fact that T is bounded, we obtain

$$\int_{\tau}^T e^{\alpha \int_u^s (-k + \sigma_r) \Lambda_r M_r dr} du < \infty \quad (3.51)$$

for all $\alpha \in \mathbb{R}$, and then by the hypothesis on μ in Assumption 2.2, and again by Proposition 2.4, we have

$$\int_{\tau}^s \mu_u \Lambda_u M_u e^{\int_u^s (-k + \sigma_r) \Lambda_r M_r dr} du < \infty, \quad \tau \leq s < T.$$

By (3.51) and by Assumption 2.2 it follows that the stochastic integral in (3.50) does not explode before T , so we have that $Y_s < \infty$ for each $s \in [\tau, T)$.

We prove that this implies $\int_{t \wedge \tau}^T Y_s^2 ds < \infty$. By the expression of Y in (3.50) we have

$$\int_{t \wedge \tau}^T Y_s^2 ds = \int_{t \wedge \tau}^T Y_s^2 \frac{1}{M_s^2 \Lambda_s^2 \sigma_s^2} d[Y, Y]_s$$

(by the Kunita-Watanabe inequality)

$$\leq \left(\int_{t \wedge \tau}^T Y_s^4 d[Y, Y]_s \right)^{1/2} \left(\int_{t \wedge \tau}^T \frac{1}{M_s^4 \Lambda_s^4 \sigma_s^4} d[Y, Y]_s \right)^{1/2}$$

(by the occupation time formula)

$$= \left(\int_{-\infty}^{\infty} a^4 L_T^a da \right)^{1/2} \left(\int_{t \wedge \tau}^T \frac{1}{M_s^2 \Lambda_s^2 \sigma_s^2} ds \right)^{1/2} < \infty : \quad (3.52)$$

the first integral is finite because the local time L_T^a has bounded support in $(-\infty, \infty)$, since Y does not explode before T , and the second one is finite by Assumption 2.2 and Proposition 2.4. Then the result follows by Proposition 3.10. \square

Proposition 3.13. *Under Assumption 2.2 the process*

$$\mathcal{E} \left(\int_0^s |\alpha_u^{t,2}| dB_u^2 \right), \quad 0 \leq s < T,$$

with $\alpha^{t,2}$ in (3.15) is a martingale for each $t \in [0, T)$.

Proof. We have that

$$|\alpha_s^{t,2}| \leq \frac{1}{\sigma_s} \left(\frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k|\beta_s| \right), \quad \tau \wedge t \leq s < T.$$

Let \tilde{Y} be the process associated to $\frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k|\beta|$ in Proposition 3.10, and \bar{Y} the one associated to $k|\beta|$.

We have

$$\begin{aligned} \tilde{Y}_s &= \frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k|\beta_s| + k \int_{\tau}^s \sigma_u \Lambda_u M_u \tilde{Y}_u e^{-k \int_u^s \Lambda_r M_r dr} du \\ &= \frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k|\beta_s| + k \int_{\tau}^s \sigma_u \Lambda_u M_u \bar{Y}_u e^{-k \int_u^s \Lambda_r M_r dr} du \\ &\quad + k \int_{\tau}^s \sigma_u \Lambda_u M_u (\tilde{Y}_u - \bar{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du \\ &= \frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + \bar{Y}_s + k \int_{\tau}^s \sigma_u \Lambda_u M_u (\tilde{Y}_u - \bar{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du, \quad \tau \leq s < T, \end{aligned}$$

and consequently, for $\bar{D}_s := \tilde{Y}_s - \bar{Y}_s$,

$$d\bar{D}_s = d\left(\frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s\right) + k\Lambda_s M_s \left[(\sigma_s - 1)\bar{D}_s + \frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s \right] ds, \quad \tau \leq s < T,$$

so that we can write

$$\bar{D}_s = \frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k \int_{\tau}^s \left(\frac{\eta_{t,\tau} + T}{\lambda} + k\mu_u M_u \right) \sigma_u \Lambda_u e^{k \int_u^s \Lambda_r M_r (\sigma_r - 1) dr} du \quad \tau \leq s < T.$$

By Assumption 2.2 and by Proposition 2.4, with the same argument as in the proof of Proposition 3.12, we have that

$$\int_{t \wedge \tau}^T \bar{D}_s^2 ds = \int_{t \wedge \tau}^T |\tilde{Y}_s - \bar{Y}_s|^2 ds < \infty.$$

Then, since by Proposition 3.12 we have $\int_{t \wedge \tau}^T |\bar{Y}_s|^2 ds < \infty$, we obtain

$$\int_{t \wedge \tau}^T |\tilde{Y}_s|^2 ds < \infty. \quad (3.53)$$

Now call Y the process associated to $R^{t,2}$ in Proposition 3.10.

It holds

$$\begin{aligned} Y_s &= \frac{1}{\sigma_s} \left(\frac{\eta_{t,\tau} + T}{\lambda M_s} + k\mu_s + k|\beta_s| + k \int_{\tau}^s \Lambda_u M_u \tilde{Y}_u e^{-k \int_u^s \Lambda_r M_r dr} du \right) \\ &\quad + \frac{1}{\sigma_s} k \int_{\tau}^s \Lambda_u M_u (Y_u - \tilde{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du \\ &= \frac{1}{\sigma_s} \left(\tilde{Y}_s + k \int_{\tau}^s \Lambda_u M_u (Y_u - \tilde{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du \right), \quad \tau \leq s < T. \end{aligned}$$

Then we have

$$\sigma_s Y_s - \tilde{Y}_s = \Psi_s + k \int_{\tau}^s \Lambda_u M_u (\sigma_u Y_u - \tilde{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du, \quad \tau \leq s < T,$$

where $(\Psi_s)_{s \in [\tau, T]}$ is given by

$$\Psi_s = k \int_{\tau}^s \Lambda_u M_u (\tilde{Y}_u - \sigma_u \tilde{Y}_u) e^{-k \int_u^s \Lambda_r M_r dr} du, \quad \tau \leq s < T. \quad (3.54)$$

It follows that $D_s = \sigma_s Y_s - \tilde{Y}_s$ satisfies

$$dD_s = d\Psi_s + k\Lambda_s M_s \Psi_s ds, \quad \tau \leq s < T,$$

and so that it takes the form

$$D_s = \Psi_s + k \int_{\tau}^s \Lambda_u M_u \Psi_u du, \quad \tau \leq s < T.$$

Since by Assumption 2.2 the process Ψ in (3.54) does not explode before T , $D_s = \sigma_s Y_s - \tilde{Y}_s < \infty$ a.s. for each $s \in [0, T)$.

Thus, with the same argument as in the proof of Proposition 3.12 it can be proved that

$$\int_{t \wedge \tau}^T |\sigma_s Y_s - \tilde{Y}_s|^2 ds < \infty.$$

By (3.53) we then have

$$\int_{t \wedge \tau}^T |\sigma_s Y_s|^2 ds < \infty.$$

Then by the integrability hypothesis on $\frac{1}{\sigma^4}$ in (ii) of Assumption 2.2 it holds

$$\int_{t \wedge \tau}^T |Y_s|^2 ds < \infty.$$

The result then follows by Proposition 3.10 and by the fact that if Y^α is the process associated to $|\alpha^{t,2}|$ it can easily be seen that $Y_s^\alpha \leq Y_s$ a.s. for each $s \in [\tau, T)$. \square

Proposition 3.14. *Consider $(Z_{t,s}^1)_{s \in [0, T)}$ and $(Z_{t,s}^2)_{s \in [0, T)}$, with*

$$Z_{t,s}^1 = \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 \right) \tag{3.55}$$

and

$$Z_{t,s}^2 = \mathcal{E} \left(\int_0^s \alpha_u^{t,2} dB_u^2 \right), \tag{3.56}$$

where $\alpha^{t,1}$ and $\alpha^{t,2}$ are as in (3.18) and (3.15), and suppose that Assumption 2.2 holds. Then $(Z_{t,s}^1)_{s \in [0, T)}$ and $(Z_{t,s}^2)_{s \in [0, T)}$ are true martingales.

The proof follows by Proposition 3.11, by Proposition 3.13 and by the following

Lemma 3.15. *Consider $H_s = \int_0^s Y_u dB_u$ and $\bar{H}_s = \int_0^s |Y_u| dB_u$, $s \geq 0$, where Y is a stochastic process such that the stochastic integral is well defined. Then $\mathcal{E}(H)$ is a martingale if and only if $\mathcal{E}(\bar{H})$ is a martingale.*

Proof. Theorem 4.1 in [13] states that, for a general continuous local martingale H , $\mathcal{E}(H)$ is a martingale if and only if

$$\lim_{n \rightarrow \infty} Q_s(\{A_s < n\}) = 1 \text{ for all } s \geq 0,$$

where $A_s = [H, H]_s$ and $dQ_s = \mathcal{E}(H_{T_s}) dP$, and $T_s := \inf\{u \geq 0 : A_u > s\}$. Since $[H, H]_s = \int_0^s Y_u^2 du = \int_0^s |Y_u|^2 du = [\bar{H}, \bar{H}]_s$, this property holds for H if and only if it holds for \bar{H} . Hence we have the result. \square

We are now ready to state the main result of the Section:

Theorem 3.16. *Under Assumption 2.2, Q^t defined in (3.8) belongs to $\mathcal{M}_{loc}(W)$ for each $t \in [0, T)$.*

Proof The proof follows by the fact that taking $\alpha^{t,1}$ and $\alpha^{t,2}$ as in (3.18) and (3.15), with $\mu_t, \sigma_t, M, \Lambda$ and π satisfying Assumption 2.2, then $(\bar{Z}_{t,s})_{s \in [0, T)}$ with

$$\bar{Z}_{t,s} = \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 \right)$$

is a martingale with respect to time s .

This follows immediately from Proposition 3.14: $(Z_{t,s}^1)_{s \in [0, T)}$ in (3.55) and $(Z_{t,s}^2)_{s \in [0, T)}$ in (3.56) are martingales, so by Proposition 3.10 we know that $H^1 = \alpha^{t,1}$ and $H^2 = \alpha^{t,2}$ are such that the associated processes Y^1 and Y^2 defined in Proposition 3.10 do not explode before T . Taking now $H = (H^1, H^2)$, the associated process $Y = (Y^1, Y^2)$ does not explode before T as well, and this concludes the proof. \square

Remark 3.17. *Theorem 3.16 shows that our constructive model can be included in the more fundamental view of the martingale theory of bubble of [29] and [30]. To this purpose we need to admit the possibility of shifting pricing views over time as suggested in [11]. However we emphasize that our definition of bubble and the models proposed in Section 2 and further investigated in Section 4 are independent of any choice of $Q \in \mathcal{M}_{loc}(W)$. This can be seen as an advantage of this framework since the definition of Q -bubble could arise some criticisms (see [25]).*

Note that Theorem 3.16 also implies that $\mathcal{M}_{loc}(W) \neq \emptyset$, hence that our market model is arbitrage-free on $[0, T)$.

4 Liquidity induced bubbles in a network

As an illustration of the previous results, we focus on a particular example. We note however that the results of this section are of independent interest since we provide one of the few contributions on mathematical modeling of bubbles in a network. For further results on this topic, we also refer to [7], where it is shown how bubbles can have an impact on the structure of a banking network, and to [14], where the authors describe the passage from a well-connected network with high global confidence to a poorly connected network with low global confidence, producing a boom and bust cycle. Our approach is however quite different: we consider an *information* network of N investors who may be influenced by the trading activity of their neighborhoods. In particular we assume that the number N of traders in the network is big enough to guarantee that our hypothesis on the linearity of the supply curve holds. Investors may place a buy market order on the bubbly asset because they imitate neighbors in the network that have successfully bought the asset as well, eventually leading to some self-exciting herding effect.

We refer to some literature about information networks (see among others Ozsoylev and Walden [43], Ozsoylev et al. [44], Walden [55]) where investors share information with

neighbors so that, as in [44], two traders linked together buy or sell the same stock at a similar point in time.

We model the trading contagion mechanism between agents taking place from time τ via the evolution dynamics of the signed volume of market orders. Our analysis is based on some epidemiological studies, which describe how diseases spread in social networks, or how computer viruses spread from computer to computer. In particular, we here focus on the SIS model, studied for example by Pastor-Satorras and Vespignani (see [45] and [46]) to analyze virus diffusion in a population. We reinterpret virus diffusion as trading contagion and consider as a first step in our model building process the following stochastic version of the SIS model for the contagion evolution of the fraction $(\rho_t^k)_{\tau \leq t \leq T}$ of traders of degree $0 \leq k \leq N$ (i.e. traders with information channels to k other traders) that has bought the asset before or at time t :

$$d\rho_t^k = (-\delta\rho_t^k + \lambda k m_t (1 - \rho_t^k)) dt + \bar{\sigma}_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha dB_t^2, \quad \tau \leq t \leq T, \quad 0 < \rho_\tau^k < 1. \quad (4.1)$$

Here m_t is the probability that an individual at the end of an edge has bought the asset before or at time t , λ is the rate of buying contagion, δ is the rate of selling, $\bar{\sigma}^k = (\bar{\sigma}_t^k)_{\tau \leq t \leq T}$, $k = 1, \dots, N$, are progressively measurable processes, which we assume bounded from above and away from zero, and $\alpha \geq 1$. Then the evolution (4.1) guarantees that $0 \leq \rho^k \leq 1$, and we further assume that the parameters are chosen such that even $0 < \rho^k < 1$, which can be shown for example for $\alpha > 1$ and $\frac{1}{z} \lambda p_k k^2 - \delta > 0$ by using Proposition 2.4 of [38].

To determine the probability m_t , we observe that by Bayes rule, and since for any given node v it holds

$$P(\text{meet } v | \text{deg}(v) = k) = \frac{k}{\sum_j j q_j}$$

where q_j is the number of nodes with degree j , we have that

$$P(\text{deg}(v) = k | \text{meet } v) = \frac{P(\text{meet } v | \text{deg}(v) = k) P(\text{deg}(v) = k)}{P(\text{meet } v)} = \frac{k}{\frac{1}{N} \sum_j j q_j} p_k = \frac{k p_k}{z},$$

where $z := \frac{1}{N} \sum_j j q_j$ is the average degree and $p_k = P(\text{deg}(v) = k) = q_k/N$. Therefore we have

$$m_t = \sum_k P(\text{deg}(v) = k | \text{meet } v) \rho_t^k = \frac{1}{z} \sum_k k p_k \rho_t^k, \quad \tau \leq t < T. \quad (4.2)$$

Given the contagion evolution of the fraction ρ^k , we model the average signed volume of market orders of an agent of degree k by $X_t^k = \theta_t^k \rho_t^k$, where the size of market orders $(\theta_t^k)_{\tau \leq t \leq T}$ of a trader of degree k that buys the asset is given by a positive continuous process with dynamics

$$d\theta_t^k = \mu_t^k dt + \sigma_t^k dB_t^2, \quad \tau \leq t < T, \quad 0 < \theta_\tau^k, \quad (4.3)$$

where for all $k = 1, \dots, N$, $(\mu_t^k)_{\tau \leq t \leq T}$ is an adapted continuous process, and $(\sigma_t^k)_{\tau \leq t \leq T}$ is a positive adapted continuous process.² Since we have $d[\rho^k, \theta^k]_t = \bar{\sigma}_t^k \sigma_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha dt$, by

²Note that the following analysis still holds under different integrability and measurability conditions on $\bar{\sigma}$ and σ^k, μ^k .

Itô's formula it holds

$$\begin{aligned} dX_t^k &= \theta_t^k d\rho_t^k + \rho_t^k d\theta_t^k + d[\rho^k, \theta^k]_t \\ &= (-\delta X_t^k + \lambda k m_t (\theta_t^k - X_t^k) + \rho_t^k \mu_t^k + \bar{\sigma}_t^k \sigma_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha) dt \\ &\quad + (\theta_t^k \bar{\sigma}_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha + \rho_t^k \sigma_t^k) dB_t^2. \end{aligned} \quad (4.4)$$

Finally, we obtain that the signed volume of total market orders is given by $X_t = \sum_{k=0}^N q_k X_t^k$, where q_k is the number of investors of degree k . From (4.4) we thus obtain

$$dX_t = (-\delta X_t + \lambda m_t (\theta_t - n_t) + \eta_t) dt + \bar{\Sigma}_t dB_t^2, \quad (4.5)$$

with

$$n_t = \sum_k k q_k X_t^k, \quad \theta_t = \sum_k k q_k \theta_t^k, \quad \eta_t = \sum_k k q_k (\rho_t^k \mu_t^k + \bar{\sigma}_t^k \sigma_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha) \quad (4.6)$$

and

$$\bar{\Sigma}_t = \sum_k q_k (\bar{\sigma}_t^k \theta_t^k (\rho_t^k)^\alpha (1 - \rho_t^k)^\alpha + \rho_t^k \sigma_t^k) \quad (4.7)$$

We are thus in the framework³ of Section 2, with

$$\mu_t = -\delta X_t + \lambda m_t (\theta_t - n_t) + \eta_t \quad (4.8)$$

and $\sigma_t = \bar{\Sigma}_t$, leading to the following SDE for the bubble β :

$$d\beta_t = \Lambda_t M_t [-k\beta_t + 2(-\delta X_t + \lambda m_t (\theta_t - n_t) + \eta_t)] dt + 2\Lambda_t M_t \bar{\Sigma}_t dB_t^2 \quad (4.9)$$

for $\tau \leq t < T$, with explicit solution

$$\begin{aligned} \beta_t &= \beta_\tau e^{-k \int_\tau^t \Lambda_s M_s ds} + \int_\tau^t (-\delta X_s + \lambda m_s (\theta_s - n_s) + \eta_s) \Lambda_s M_s e^{-k \int_s^t \Lambda_u M_u du} ds + \\ &\quad + \int_\tau^t \bar{\Sigma}_s \Lambda_s M_s e^{-k \int_s^t \Lambda_u M_u du} dB_s^2, \quad \tau \leq t < T. \end{aligned} \quad (4.10)$$

Remark 4.1. Setting $\mu^j \equiv \bar{\sigma}^j \equiv \sigma^j \equiv 0$ for all $0 \leq j \leq N$ in (4.1) and (4.3) respectively, we can identify the driving deterministic contagion evolution for the signed volume of market orders as implied by the SIS network model approach:

$$dX_t = (-\delta X_t + \lambda m_t (\theta_t - n_t)) dt. \quad (4.11)$$

³The assumption that θ^k is driven by the same Brownian motion of ρ^k allows to show the existence of the flow by using directly the results of Section 2, but it can be easily relaxed, letting θ^k depend also on an additional Brownian motion B^θ independent of B^2 , as we do in Section 4.1.

Remark 4.2. *In the next Subsection 4.1 we consider two different types of network topologies in order to see how the characteristics of the network influence the dynamics of the expected fraction of buyers through n_t . In the first one we have a connectivity distribution which is very peaked at the average value z and decaying exponentially fast for $k \gg z$ and $k \ll z$. Examples of this kind of networks are random graph models [22] and the small-world model of Watts and Strogatz [57]. In the second one the degree distribution is more right skewed, following for example a power law, as in the Barabási and Albert preferential attachment model [6]. From (4.11) and (4.6) we can see that the expected contagion between buyers will spread faster in the second kind of network, since the distribution puts more weight on the nodes with higher degree, resulting in a bigger value of n_t in (4.6).*

We conclude the introduction of the model by showing a sufficient condition under which the above bubble specification can be represented by a flow of local martingale measures as analyzed in the general framework of the previous sections, i.e. that there exists a flow $Q^t \in \mathcal{M}_{loc}(W)$ with Radon-Nykodim derivative process

$$Z_{t,s} = \frac{dQ^t}{dP} \Big|_{\mathcal{F}_s} = \mathcal{E} \left(\int_0^s \alpha_u^{t,1} dB_u^1 + \int_0^s \alpha_u^{t,2} dB_u^2 + \int_0^s \alpha_u^{t,3} d\tilde{N}_u \right), \quad s \in [0, T], \quad (4.12)$$

such that

$$W_t^F = \mathbb{E}_{Q^t}[W_T^F | \mathcal{F}_t], \quad 0 \leq t \leq T.$$

Taking $\alpha^{t,1}$, $\alpha^{t,2}$ and $\alpha^{t,3}$ in (3.18), (3.15) and (3.17) respectively we only need to show that that Z in (4.12) is in fact a martingale.

Proposition 4.3. *Assume that there exists a $\bar{k} \in 1, \dots, N$ such that $\theta_t^{\bar{k}} > \epsilon$ a.s. for all $t \in [\tau, T]$, where $\epsilon > 0$. Then for each $t \in [0, T]$, $(Z_{t,u})_{u \in [0, T]}$ is a (P, \mathcal{F}) -martingale.*

Proof. We show that μ and Σ in (4.8) and (4.7) satisfy Assumption 2.2.

We have $\int_\tau^T \mu_s^2 ds < \infty$ since $m, \bar{\sigma}^j$ are bounded and $\sigma^j, \mu^j, X, \theta, n$ are continuous processes for $j \in 1, \dots, N$. Analogously one can show $\int_\tau^T \bar{\Sigma}_s^2 ds < \infty$.

Finally by using that $\sigma^{\bar{k}}, \rho^{\bar{k}} \geq 0$ and that $\theta^{\bar{k}}, \bar{\sigma}^{\bar{k}}$ are bounded away from zero, it is easy to see that

$$\int_\tau^T \frac{1}{\bar{\Sigma}_s^4} ds \leq \frac{C}{q_{\bar{k}}^4} \int_\tau^T \frac{1}{(\rho_s^{\bar{k}})^{4\alpha} (1 - \rho_s^{\bar{k}})^{4\alpha}} ds \quad (4.13)$$

for some constant C . We can show that the integral on the right side of (4.13) is finite by applying Theorem 2.6 of [38]. \square

4.1 Analysis of the model

We now comment on our model and specify how the evolution of the bubble described in (4.9) depends on the involved parameters as well as on the structure of the network.

The evolution of the bubble is characterized by two different phases: in the first one the

bubble builds up, since the quick increase of the signed volume of market orders X dominates in equation (2.2). However, after a while the processes ρ^k in equation (4.1) tend towards an equilibrium in which the drift of ρ^k vanishes. When this drift's component (and also the contribution of η in (4.5)) is sufficiently small, the mean reverting term of equation (2.2) starts to dominate, leading to the burst of the bubble and to the second phase, i.e. the decrease of the bubble towards zero.

In the ascending phase, assuming first for illustration purposes the process $(\theta_t)_{t \geq \tau}$ to be constantly equal to $\theta > 0$ and $\bar{\sigma}^j = 0$ for all $0 \leq j \leq N$, the essential force of the bubble is given by the deterministic contagion mechanism (4.11) driving the signed volume of market orders X in (4.5). The contagion accelerates to a maximum and then slows down. In this way X evolves along an ‘‘S’’ shape as shown in Figure 1 growing towards an equilibrium/maximum that is increasing in the volume term θ and the contagion rate λ and decreasing in the recovery rate δ . Further, the speed at which X grows towards the maximum is increasing in λ and decreasing in δ . However, since both the length and the maximum of observed speculation bubbles are highly uncertain, we randomize this mechanism by letting θ be a stochastic process. The impact of a random volume term θ will be to modify the ‘‘S’’ pattern by allowing the bubble to slow down or pick up in a random way until it reaches a random maximum. In the bursting phase, the dynamics of the bubble will be dominated by the mean reverting factor k , which drives the bubble down.

We now focus on the impact of the choice of the underlying network on the dynamics of the bubble. We compare two different cases, an Erdős-Rényi network with Poisson degree distribution

$$p_j = \frac{e^{-\tilde{\lambda}} \tilde{\lambda}^j}{j!}, \quad j \in \mathbb{N}, \quad \tilde{\lambda} \in \mathbb{R},$$

and a scale-free network with a power law distribution

$$p_j \sim j^{-\gamma}, \quad 2 < \gamma < 3, \quad j \in \mathbb{N}. \quad (4.14)$$

The Erdős-Rényi network has a degree distribution which is very peaked around the mean degree z , whereas the scale-free one, that is well-known to better represent real world information networks (see [44]), has a much larger right tale, which allows for a more heterogeneous degree distribution with some nodes being very connected and others less (core-periphery structure).

For simplicity, we consider the following deterministic specifications: we set $\bar{\sigma}^j = 0$ for all $0 \leq j \leq N$ and assume the processes $(M_t)_{t \geq \tau}$, $(\Lambda_t)_{t \geq \tau}$ and $(\theta_t)_{t \geq \tau}$ to be constantly equal to $M = \Lambda = \theta = 1$. Further, we choose $\tau = 0$.

We take two different values of γ in (4.14), i.e. $\gamma_1 = 2.2$ and $\gamma_2 = 2.5$, obtaining therefore a more connected network (with $z = z_1 \sim 3.2$) and a less connected one (with $z = z_2 \sim 1.9$). We consider as well two Erdős-Rényi networks with $z = z_1 \sim 3.2$ and $z = z_2 \sim 1.9$, respectively. We take the distribution p_j up to a maximum degree that corresponds to a network with 5000 nodes, see 3.3.2 of [40].

In Figure 1 we illustrate the trajectories of X for the four different networks taking $\lambda = 1$, $\delta = 1$. One can notice that both the mean degree and the degree heterogeneity play a key




Figure 1: Deterministic fraction of buyers for different networks, with $\lambda = 1$, $\delta = 1$.

role in the evolution of X : in particular, both of them are positively correlated with the speed of the increase. It can also be seen that in the Erdős-Rényi network, i.e. in the less right skewed one, as well as in the less connected networks, the fraction reaches its equilibrium later in time.

We then focus on the behaviour of the bubble and consider a mean reversion level $k = 0.4$ in (4.10). In Figure 2 and Figure 3 we show the maximum reached by the bubble as a function of λ and δ respectively, whereas in Figure 4 and Figure 5 we plot the time needed to reach the maximum, again as a function of λ and δ respectively.

Figure 5 shows that the time to the maximum is decreasing in δ in the scale-free network and increasing in δ in the Erdős-Rényi one, i.e. the two networks give rise to different behaviors. It can be seen that for small λ and big δ the maximum is higher in the case of the scale-free network, whereas the opposite holds for big λ and small δ . On the other hand, the time needed by the bubble to attain the maximum is always higher in the case of the Erdős-Rényi network.

In our analysis up to this point, we have taken the process $\theta = (\theta_t)_{t \geq \tau}$ to be constant. We now show the influence of the process θ on the dynamics of the bubble assuming that it satisfy the SDE

$$d\theta_t = \sigma^\theta \theta_t dB_t^3, \quad \tau \leq t < T,$$

where $\sigma^\theta = 0.4$, taking $\delta = 0.2$, $\lambda = 0.4$, $\Lambda = 0.5$, $k = 1$, $\bar{\sigma}^j = 0.1$ for all $0 \leq j \leq N$, $\tau = 0$, $T = 7$, $M = 1$, $\theta_0 = 3$. See for example Figure 7 and Figure 6 for the case of a scale-free network with mean degree $z = 3.2$.

The influence of the process θ on the bubble is apparent. If θ has an increase from its initial value, the bubble bursts relatively late, see the yellow dynamics: in this sense, the growth of θ can postpone the burst of the bubble. The other trajectories evolve similarly to each other up to the point where the corresponding processes θ differ. In the blue case, θ decreases and the bubble bursts soon. For the red dynamics, θ increases, making the bubble growing more.




Figure 2: Maximum value of the bubble as a function of λ with $\delta = 1$, $k = 0.4$.




Figure 3: Maximum value of the bubble as a function of δ with $\lambda = 1$, $k = 0.4$.




Figure 4: Time to the maximum as a function of λ with $\delta = 1$, $k = 0.4$.




Figure 5: Time to the maximum as a function of δ with $\lambda = 1$, $k = 0.4$.




Figure 6: Example of trajectories of θ with $\mu^\theta = 0$, $\sigma^\theta = 0.2$.




Figure 7: Corresponding trajectories of the bubble in a scale-free network with mean degree $z = 3.2$.

We conclude the section illustrating the impact of the structure of the network by showing three trajectories of the bubble in Figure 8 for the the scale-free case and in Figure 9 for the Erdős-Rényi one. We choose $\delta = 0.2$, $\lambda = 0.3$, $\Lambda = 0.5$, $k = 1$, $\bar{\sigma}^j = 0.2$ for all $0 \leq j \leq N$, $\tau = 0$, $T = 3$, $M = 1$, $\theta_0 = 3$ and $\sigma^\theta = 0.2$. We can see that the bubble builds up faster in the scale-free network, but at the same time the trajectories have a steeper decrease, and therefore the effect of the burst of the bubble is more dramatic. On the other hand, Figure 7 shows that a quick decrease of θ can also lead to a quick burst, and then to an hard landing.

4.2 Model testing on real data

In this subsection we test some features of our model on real data. Since we were not able to find tick by tick data for the signed volume of market orders of well-known bubbles of the past such as for example for the *dot com bubble*, we consider the asset prices Alphabet Inc (NASDAQ:GOOG) and Amazon.com Inc (NASDAQ:AMZN). For these stocks we could obtain tick by tick data for the signed volume of their market orders starting from the first months of 2016. These companies, as it can be seen also by the prices reported in Figure 10 and in Figure 11, have experienced in the last years a boom, which has brought many financial analysts to propose the presence of a new tech bubble, after the *dot com mania* of the late 1990s (see for example [9], [42], [52], [53]).

We consider the realized signed volume of market orders since 2016. As shown in Figures 12 and 13, the signed volume tends to increase over time, for both Alphabet and Amazon. This behavior indicates the tendency of traders to invest in these companies, contributing to the increase of the price in line with our model.

Our aim is to investigate whether typical trading behaviour in a bubble environment is




Figure 8: Example of trajectories for a bubble in a scale-free network.




Figure 9: Example of trajectories for a bubble in an Erdős-Rényi network.




Figure 10: Price in USD of Alphabet Inc, June 2016 - October 2017.




Figure 11: Price in USD of Amazon.com Inc, July 2016 - October 2017.

captured in our model. In particular, since we deal with a relatively small time window of a potential bubble, we calibrate the coefficients of the deterministic component \bar{X} for X in (4.11), underlying the signed volume of market orders on the observed data for Amazon and Google by employing a quadratic regression. In doing so, we further assume $\bar{\sigma}^j = 0$ for $j \in 1, \dots, N$, the process $(\theta_t)_{t \geq \tau}$ to be constant and that all the nodes of the network have the same degree $d = 3$, i.e. that the degree distribution of our network is a Dirac delta centered in $d = 3$. In this way, \bar{X} has dynamics

$$d\bar{X}_t = (-a\bar{X}_t^2 + b\bar{X}_t)dt, \quad t \geq 0, \quad (4.15)$$

where $a = \frac{3\lambda}{\theta}$ and $b = 3\lambda - \delta$.

Further, for every asset we compute the mean relative squared error of the prediction, where the relative squared error at point t_i is defined as $(\bar{X}_{t_i} - X_{t_i}^{obs})^2 / (X_{t_i}^{obs})^2$. Here \bar{X} is the signed volume predicted by our model and X^{obs} is the observed signed volume. Moreover, we also compute the 99% and 95% confidence intervals for the estimates of the two parameters a and b in (4.15).

The mean relative squared errors are shown in Table 1.

Alphabet Inc	Amazon.com
0.0511	0.0291

Table 1: Mean relative squared error of the predicted signed volume of market orders versus the observed data for Alphabet Inc and Amazon.com Inc.

The confidence intervals of the parameters, together with the selected values, are shown in Tables 2 and 3, for Alphabet Inc and Amazon.com Inc respectively.

	99% c.i., l. endp.	95% c.i., l. endp.	Parameter	95% c.i., r. endp.	99% c.i., r. endp.
a	1.3013	1.3015	1.3016	1.3017	1.3019
b	10.8271	10.8381	10.8727	10.9074	10.9184

Table 2: Confidence intervals for parameters a and b in equation (4.15), for Alphabet Inc.

	99% c.i., l. endp.	95% c.i., l. endp.	Parameter	95% c.i., r. endp.	99% c.i., r. endp.
a	0.3557	0.3559	0.3560	0.3561	0.3563
b	6.6935	6.7113	6.7676	6.8239	6.8417

Table 3: Confidence intervals for parameters a and b in equation (4.15), for Amazon.com Inc.

In Figures 12 and 13, we illustrate how our model changes when the parameters a and b are equal to the endpoints of the 99% confidence intervals. In particular, we show the trajectories




Figure 12: Realized signed volume of market orders and deterministic signed volume given by eq. (4.11), Alphabet Inc, June 2016 - October 2017.

of the predicted signed volume of market orders when a is equal to the right endpoint of the confidence intervals and b to the left one and viceversa, i.e. when a is equal to the left endpoint and b to the right one.

In Figure 12 and Figure 13 we can observe the “S” behavior discussed in Section 4.1. We remark that since we perform a local analysis by considering a specific short time window with constant θ , this behaviour cannot be directly interpreted as indication for a decreasing phase of the bubble. In the next time window the signed volume may start to grow steeply again, due to the impact of a stochastic θ . In this case the curve describing the evolution of the signed volume would also grow for a longer time, distorting the “S” shape as illustrated in Figure 6 and in Figure 7.

We can conclude that the analysis shows the flexibility of our model and its capacity of

1. describing both the increasing and the descending phase of the bubble;
2. capturing the impact of signed volume market orders on bubbles’ formation and burst;
3. taking into account the underlying network structure in the contagion process of a bubble’s evolution;
4. describing typical features of a bubble’s behavior like steep increase and hard landing.




Figure 13: Realized signed volume of market orders and deterministic signed volume given by eq. (4.11), Amazon.com Inc, June 2016 - October 2017.

Acknowledgments

The authors would like to thank two anonymous referees and the associated editor for a very careful reading and useful comments which have contributed to improve the quality of the paper.

References

- [1] D. Abreu and M.K. Brunnermeier. Bubbles and crashes. *Econometrica*, 71(1):173–204, 2003.
- [2] D. Acemoglu, V.M. Carvalho, A. Ozdaglar, and A. Tahbaz-Salehi. The Network Origins of Aggregate Fluctuations. *Econometrica*, 80(5):1977–2016, 2012.
- [3] F. Allen and D. Gale. Bubbles and crisis. *The Economic Journal*, 110:236–255, 2000.
- [4] F. Allen and D. Gale. Financial Contagion. *The Journal of Political Economy*, 108(1): 1–33, 2000.
- [5] H. Amini, R. Cont, and A. Minca. Resilience to contagion in financial networks. *Mathematical Finance*, 26(2):329–365, 2016.

- [6] A.-L. Barabási and R. Albert. Emergence of scaling in random networks. *Science*, 286: 509–512, 1999.
- [7] P. Battiston. Rational bubbles in closed economies. 2015.
- [8] P. Bayer, K. Mangum, and J.W. Roberts. Speculative fever: Investor contagion in the housing bubble. Technical report, National Bureau of Economic Research, 2016.
- [9] J. Bercovici. Yes, it’s a Tech Bubble. *Inc.*, 2017. URL <https://www.inc.com/magazine/201509/jeff-bercovici/are-we-in-a-tech-bubble.html>.
- [10] F. Biagini and S. Nedelcu. The formation of financial bubbles in defaultable markets. *SIAM Journal of Financial Mathematics*, 6(1):530–558, 2015.
- [11] F. Biagini, H. Föllmer, and S. Nedelcu. Shifting martingale measures and the slow birth of a bubble. *Finance and Stochastics*, 18(2):297–326, 2014.
- [12] M. Blais and P. Protter. An analysis of the supply curve for liquidity risk through book data. *International Journal of Theoretical and Applied Finance*, 13(6):821–838, 2010.
- [13] S. Blei and H.-J. Engelbert. On exponential local martingales associated with strong Markov continuous local martingales. *Stochastic Process. Appl.*, 119(9):2859–2880, 2009.
- [14] J.P. Bouchard, D. Challet, and J. da Gama Batista. Sudden Trust Collapse in Networked Societies. *Eur Phys*, 88(3):1–11, 2015.
- [15] U. Cetin, R. Jarrow, P. Protter, and M. Warachka. Pricing Options in an Extended Black Scholes Economy with Illiquidity: Theory and Empirical Evidence. *The Review of Financial Studies*, 19(2):493–529, 2006.
- [16] Y. Choi and R. Douady. Financial Crisis and Contagion: A Dynamical Systems Approach. 2011. URL <http://ssrn.com/abstract=1733706>.
- [17] Y. Choi and R. Douady. Chaos and Bifurcation in 2007-08 Financial Crisis. *Management Science*, 2011.
- [18] R. Cont, A. Moussa, and E.B. Santos. Networks structure and systemic risk in banking systems. Technical report, Handbook of Systemic Risk, Cambridge University Press, 2013.
- [19] A.M.G. Cox and D.G. Hobson. Local martingales, bubbles and option prices. *Finance Stochastics*, 9(4):477–492, 2005.
- [20] J.B. DeLong, A. Shleifer, L. Summers, and R. Waldmann. Noise trader risk in financial markets. *Journal of Political Economy*, 98(4):703–738, 1990.
- [21] P.E. Earl, T-C Peng, and J. Potts. Decision-rule cascades and the dynamics of speculative bubbles. *Journal of Economic Psychology*, 28:351–364, 2007.

- [22] P. Erdős and A. Rényi. On the evolution of random graphs. *Publication of the Mathematical Institute of the Hungarian Academy of Science*, 17-61, 1960.
- [23] H. Föllmer. Equilibria in financial markets with heterogeneous agents: A probabilistic perspective. *Journal of Mathematical Economics*, 41(1-2):123–155, 2005.
- [24] P. Gai and S. Kapadia. Contagion in financial networks. *Proceedings of the Royal Society*, 466:2401–2423, 2010.
- [25] P. Guasoni and M. Rasonyi. Fragility of arbitrage and bubbles in local martingale diffusion models. *Finance and Stochastics*, 19(2):215–231, 2015.
- [26] J.M. Harrison and D.M. Kreps. Speculative investor behavior in a stock market with heterogeneous expectations. *The Quarterly Journal of Economics*, 92(2):323–336, 1978.
- [27] R. Jarrow and P. Protter. Forward and futures prices with bubbles. *International Journal of Theoretical and Applied Finance*, 12(7):901–924, 2009.
- [28] R. Jarrow and P. Protter. Foreign currency bubbles. *Review of Derivatives Research*, 14(1):67–83, 2011.
- [29] R. Jarrow, P. Protter, and K. Shimbo. Asset price bubbles in complete markets. *Advances in Mathematical Finance*, In Honor of Dilip B. Madan:105–130, 2007.
- [30] R. Jarrow, P. Protter, and K. Shimbo. Asset price bubbles in incomplete markets. *Mathematical Finance*, 20(2):145–185, 2010.
- [31] R. Jarrow, Y. Kchia, and P. Protter. How to detect an asset bubble. *SIAM Journal on Financial Mathematics*, 2:839–865, 2011.
- [32] R. Jarrow, P. Protter, and A. Roch. A Liquidity Based Model for Asset Price. *Quantitative Finance*, 12(1):1339–1349, 2012.
- [33] T. Kaizoji. Speculative bubbles and crashes in stock markets: an interacting-agent model of speculative activity. *Physica A*, 287:493–506, 2000.
- [34] M. Loewenstein and G.A. Willard. Rational equilibrium asset-pricing bubbles in continuous trading models. *Journal of Economic Theory*, 91(1):17–58, 2000.
- [35] T. Lux. Herd behaviour, bubbles and crashes. *The Economic Journal*, 105(431):881–896, 1995.
- [36] A. Mazzon. Financial asset bubbles in networks. *PhD thesis*, in preparation.
- [37] A. Mijatovic and M. Urusov. On the martingale property of certain local martingales. *Probability Theory and Related Fields*, 152(1-2):1–30, 2012.

- [38] A. Mijatovic and M. Urusov. Convergence of integral functionals of one-dimensional diffusions. *Electronic Communications in Probability*, 17(61):1–13, 2012.
- [39] E.M. Miller. Risk, uncertainty, and divergence of opinion. *The Journal of Finance*, 32(4):1151–1168, 1977.
- [40] M.E.J. Newman. The structure and function of complex networks. *SIAM review*, 45(2):167–256, 2003.
- [41] M.E.J. Newman, S.H. Strogatz, and D.J. Watts. Random graphs with arbitrary degree distributions and their applications. *Physical review E*, 64(2):026118, 2001.
- [42] A. Ozimek. There obviously is a tech bubble, but hopefully that doesn't matter. *Forbes*, 2017. URL <https://www.forbes.com/sites/modeledbehavior/2017/07/16/there-obviously-is-a-tech-bubble-but-hopefully-that-doesnt-matter/#737270d9d426>.
- [43] H. Ozsoylev and J. Walden. Asset Pricing in Large Information Networks. *Journal of Economic Theory*, 146:2252–2280, 2011.
- [44] H. Ozsoylev, J. Walden, D. Yavuz, and R. Bildick. Investor Networks in the stock Market. *The Review of Financial Studies*, 27(5):1323–1366, 2014.
- [45] R. Pastor-Satorras and A. Vespignani. Epidemic dynamics and endemic states in complex networks. *Phys. Rev. E.*, 63(066117), 2001.
- [46] R. Pastor-Satorras and A. Vespignani. Epidemic spreading in scale-free networks. *Phys. Rev. Lett.*, 86:3200–3203, 2001.
- [47] P. Protter. *Stochastic integration and differential equations, Second Edition*. Springer-Verlag, Berlin, 2005.
- [48] P. Protter. *A mathematical theory of financial bubbles*, volume 2081 of Lecture Notes in Mathematics of V. Henderson and R. Sincar editors, Paris-Princeton Lectures on Mathematical Finance. Springer, 2013.
- [49] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion, Third Edition*. Springer-Verlag, New York, 1999.
- [50] J. Scheinkman and W. Xiong. Overconfidence and speculative bubbles. *Journal of political economy*, 111(6):1183–1219, 2003.
- [51] J. Scheinkman and W. Xiong. Speculation, trading and bubbles. Economic Theory Center Research 050, Princeton University, 2013.
- [52] R. Seria. The tech bubble: how close is it to bursting? *Telegraph*, 2017. URL <http://www.telegraph.co.uk/technology/2017/06/22/tech-bubble-close-bursting/>.

- [53] R. Sharma. When will the tech bubble burst? *New York Times*, 2017. URL <https://www.nytimes.com/2017/08/05/opinion/sunday/when-will-the-tech-bubble-burst.html>.
- [54] J. Tirole. On the possibility of speculation under rational expectations. *Econometrica*, 53(6):1163–1182, 1982.
- [55] J. Walden. Trading, profits, and volatility in a dynamic information network model. 2014. URL <https://dx.doi.org/10.2139/ssrn.2561055.html>.
- [56] D. Watts. A simple model of global cascades on random networks. *Proc. Natl Acad. Sci. USA*, 99:5766–5771, 2002.
- [57] D.J. Watts and S.H. Strogatz. Collective dynamics of 'small-world' networks. *Nature*, 393:440–442, 1998.
- [58] B. Wong and C. Heyde. On the martingale property of stochastic exponentials. *Journal of Applied Probability*, 41(3):654–664, 2004.
- [59] W. Xiong. Bubbles, crises, and heterogeneous beliefs. Technical report, Princeton University, Princeton, NJ, 2012.
- [60] S. Zhuk. Speculative bubbles, information flow and real investment. *Information Flow and Real Investment*, 2013.