

THE CARDIAC BIDOMAIN MODEL AND HOMOGENIZATION

E. GRANDELIUS AND K. H. KARLSEN

ABSTRACT. We provide a rather simple proof of a homogenization result for the bidomain model of cardiac electrophysiology. Departing from a microscopic cellular model, we apply the theory of two-scale convergence to derive the bidomain model. To allow for some relevant nonlinear membrane models, we make essential use of the boundary unfolding operator. There are several complications preventing the application of standard homogenization results, including the degenerate temporal structure of the bidomain equations and a nonlinear dynamic boundary condition on an oscillating surface.

CONTENTS

1. Introduction	1
2. Preliminaries	5
2.1. Some functional spaces and tools	5
2.2. Two-scale convergence	7
2.3. Unfolding operators	10
3. Microscopic bidomain model	12
3.1. Assumptions on the data	12
3.2. Weak solutions	14
3.3. Existence of solution and a priori estimates	15
4. The homogenization result	21
4.1. Extracting two-scale limits	23
4.2. The nonlinear terms and strong convergence	24
4.3. Concluding the proof of Theorem 4.3	28
References	29

1. INTRODUCTION

The bidomain model [39, 13, 38] is widely used as a quantitative description of the electric activity in cardiac tissue. The relevant unknowns are the intracellular (u_i) and extracellular (u_e) potentials, along with the so-called transmembrane potential ($v := u_i - u_e$). In this model, the intra- and extracellular spaces are considered as two separate homogeneous domains superimposed on the cardiac domain. The two domains are separated by the cell membrane creating a discontinuity surface for the cardiac potential. Conduction of electrical signals in cardiac tissue relies on the flow of ions through channels in the cell membrane. In the bidomain model, the celebrated Hodgkin-Huxley [23] framework is used to dynamically couple the intra- and extracellular potentials through voltage gated ionic channels.

Date: November 19, 2018.

2010 Mathematics Subject Classification. Primary: 35K57, 35B27; Secondary: 35K65, 92C30.

Key words and phrases. Bidomain equations, cardiac electric field, reaction-diffusion system, degenerate, weak solution, homogenization, two-scale convergence, unfolding operator.

This work was supported by the Research Council of Norway (project 250674/F20).

The bidomain model can be viewed as a PDE system consisting of two degenerate reaction-diffusion equations involving the unknowns u_i, u_e, v and two conductivity tensors σ_i, σ_e . These equations are supplemented by a nonlinear ODE system for the dynamics of the ion channels. The bidomain model is often derived heuristically by interpreting σ_i, σ_e as some sort of “average” conductivities, applying Ohm’s electrical conduction law and the continuity equation (conservation of electrical charge) to the intracellular and extracellular domains [13, 38].

Starting from a more accurate microscopic (cell-level) model of cardiac tissue, with the heterogeneity of the underlying cellular geometry represented in great detail, it is possible to heuristically derive the bidomain model (tissue-level) using the multiple scales method of homogenization. This derivation was first carried out in [31]. It should be noted that the microscopic model is in general too complex to allow for full organ simulations, although there have been some very recent efforts in that direction [40]. The complexity of cell-level models, which themselves can be heuristically derived from the Poisson-Nernst-Planck equations [35], motivates the search for simpler homogenized (macroscopic) models. The work [31] assumes, as we do herein, that cardiac tissue can be viewed as a uniformly oriented periodic assembly of cells (see also [14, 22]). There have been some attempts to remove this assumption. We refer to [26, 25, 36] for extensions to somewhat more realistic tissue geometries.

Despite the widespread use of the bidomain model, there are few mathematical rigorous derivations of the model from a microscopic description of cardiac tissue. From a mathematical point of view, rigorous homogenization is often linked to the study of the asymptotic behavior (convergence) of solutions to PDEs with oscillating coefficients. In the literature several approaches have been developed to handle this type of problem, like Tartar’s method of oscillating test functions, Γ -convergence, two-scale convergence, and the unfolding method. We refer to [12] for an accessible introduction to the mathematics of homogenization and for an overview of the different homogenization methods.

We are aware of two earlier works [4, 34] containing rigorous homogenization results for the bidomain model (but see [15, 16, 43] for examples of elliptic and parabolic equations on “two-component” domains). With a fairly advanced proof involving Γ -convergence, the De Giorgi “minimizing movement” approach, time-discretization, variational problems, and two limit procedures, the homogenization result in [34] covers the generalized FitzHugh-Nagumo ionic model [17]. The proof of the result in [4] is more basic in the sense that it employs only two-scale convergence arguments, but it handles only a restricted class of ionic models.

We mention that there are several complications preventing the application of standard homogenization results (for elliptic/parabolic equations) to the bidomain equations, including its degenerate structure (seen at the tissue-level), resulting from differing anisotropies of the intra- and extracellular spaces, and the highly nonlinear, oscillating dynamic boundary condition (seen at the cell-level).

The main contribution of our paper is to provide a simple homogenization proof that can handle some relevant nonlinear membrane models (the generalized FitzHugh-Nagumo model), relying only on basic two-scale convergence techniques.

We now explain our contribution in more detail. The point of departure is the following microscopic model [13, 14, 22, 41] for the electric activity in cardiac tissue:

$$(1.1) \quad \begin{aligned} -\operatorname{div}(\sigma_i^\varepsilon \nabla u_i^\varepsilon) &= s_i^\varepsilon && \text{in } (0, T) \times \Omega_i^\varepsilon, \\ -\operatorname{div}(\sigma_e^\varepsilon \nabla u_e^\varepsilon) &= s_e^\varepsilon && \text{in } (0, T) \times \Omega_e^\varepsilon, \\ \varepsilon(\partial_t v^\varepsilon + I(v^\varepsilon, w^\varepsilon)) &= -\nu \cdot \sigma_i^\varepsilon \nabla u_i^\varepsilon && \text{on } (0, T) \times \Gamma^\varepsilon, \\ \varepsilon(\partial_t v^\varepsilon + I(v^\varepsilon, w^\varepsilon)) &= -\nu \cdot \sigma_e^\varepsilon \nabla u_e^\varepsilon && \text{on } (0, T) \times \Gamma^\varepsilon, \\ \partial_t w^\varepsilon &= H(v^\varepsilon, w^\varepsilon) && \text{on } (0, T) \times \Gamma^\varepsilon, \end{aligned}$$

where ν denotes the unit normal pointing out of Ω_i^ε (and into Ω_e^ε). Cardiac tissue consists of an assembly of elongated cylindrical-shaped cells coupled together (end-to-end and side-to-side) to provide intercellular communication. The entire cardiac domain $\Omega \subset \mathbb{R}^3$ is viewed as a “two-component” domain and split into two ε -periodic open sets $\Omega_i^\varepsilon, \Omega_e^\varepsilon$ corresponding to the intra- and extracellular spaces. The sets $\Omega_i^\varepsilon, \Omega_e^\varepsilon$, which are assumed to be disjoint and connected, are separated by an ε -periodic surface Γ^ε representing the cell membrane, so that $\Omega = \Omega_i^\varepsilon \cup \Omega_e^\varepsilon \cup \Gamma^\varepsilon$. The main geometrical assumption is that the intra- and extracellular domains are ε -dilations of some reference cells $Y_i, Y_e \subset Y := [0, 1]^3$, periodically repeated over \mathbb{R}^3 . Although our results are valid for general Lipschitz domains, for simplicity of presentation, we assume that the entire cardiac domain Ω is the open cube

$$(1.2) \quad \Omega = (0, 1) \times (0, 1) \times (0, 1).$$

In (1.1), σ_j^ε is the conductivity tensor and s_j^ε is the stimulation current, relative to Ω_j^ε for $j = i, e$. The functions $s_i^\varepsilon, s_e^\varepsilon$ are assumed to be at least bounded in L^2 , independently of ε . As usual in homogenization theory, the conductivity tensors $\sigma_i^\varepsilon, \sigma_e^\varepsilon$ are assumed to have the form

$$\sigma_j^\varepsilon(x) = \sigma_j \left(x, \frac{x}{\varepsilon} \right), \quad j = i, e,$$

where $\sigma_j = \sigma_j(x, y)$ satisfies the usual conditions of uniform ellipticity and periodicity (in y). Despite the fact that the inhomogeneities of the domains impose ε -oscillations in the conductivity tensors (via gap junctions), the main source of inhomogeneity in the microscopic model is not the conductivities σ_i^ε and σ_e^ε , but the domains Ω_i^ε and Ω_e^ε themselves. We allow for inhomogeneous and oscillating conductivities for the sake of generality.

We denote by u_j^ε the electric potential in Ω_j^ε ($j = i, e$). On Γ^ε , $v^\varepsilon := u_i^\varepsilon - u_e^\varepsilon$ is the transmembrane potential and $I(v^\varepsilon, w^\varepsilon)$ is the ionic current depending on v^ε and a gating variable w^ε . The left-hand side of the third and fourth equations in (1.1) describes the current across the membrane as having a capacitive component, depending on the time derivative of the transmembrane potential, and a nonlinear ionic component I corresponding to the chosen membrane model. In this article we consider the generalized FitzHugh-Nagumo model [17]. We choose to focus on this membrane model for definiteness, but our arguments can be adapted to many other models satisfying reasonable technical assumptions [7, 8, 13, 38, 41, 42].

For each fixed $\varepsilon > 0$, the functions $\sigma_j^\varepsilon, s_j^\varepsilon, I, H$ in (1.1) are given and we wish to solve for $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$. To this end, we must augment the system (1.1) with initial conditions for $v^\varepsilon, w^\varepsilon$ and Neumann-type boundary conditions for $u_i^\varepsilon, u_e^\varepsilon$ (ensuring no current flow out of the heart):

$$(1.3) \quad \begin{aligned} v^\varepsilon|_{t=0} &= v_0^\varepsilon && \text{in } \Omega, \quad w^\varepsilon|_{t=0} = w_0^\varepsilon && \text{in } \Omega, \\ n \cdot \sigma_j \nabla u_j^\varepsilon &= 0 && \text{on } (0, T) \times (\partial\Omega \cap \partial\Omega_j^\varepsilon), \quad j = i, e, \end{aligned}$$

where n is the outward unit normal to Ω . It is proved in [14, 41] that the microscopic bidomain model (1.1), (1.3) possesses a unique weak solution. This solution satisfies

a series of a priori estimates. For us it is essential to know how these estimates depend on the parameter ε . We will therefore outline a proof of these estimates.

The dimensionless number ε is a small positive number representing the ratio of the microscopic and macroscopic scales, that is, considering Ω as fixed, it is proportional to the cell diameter. The goal of homogenization is to investigate the limit of a sequence of solutions $\{(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)\}_{\varepsilon>0}$ to (1.1), (1.3). By the multiple scales method [12, 14, 22], the electric potentials $u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon$ and the state variable w^ε exhibit the following asymptotic expansions in powers of the parameter ε :

$$\begin{aligned} u_j^\varepsilon(t, x, y) &= u_j(t, x, y) + \varepsilon u_j^{(1)}(t, x, y) + \varepsilon^2 u_j^{(2)}(t, x, y) + \dots \quad (j = i, e), \\ v^\varepsilon(t, x, y) &= v(t, x, y) + \varepsilon v^{(1)}(t, x, y) + \varepsilon^2 v^{(2)}(t, x, y) + \dots, \\ w^\varepsilon(t, x, y) &= w(t, x, y) + \varepsilon w^{(1)}(t, x, y) + \varepsilon^2 w^{(2)}(t, x, y) + \dots, \end{aligned}$$

where $y = x/\varepsilon$ denotes the microscopic variable, and each term in the expansions is a function of both the slow (macroscopic) variable x and the fast (microscopic) variable y , periodic in y . Substituting the above expansions into (1.1), and equating all terms of the same orders in powers of ε , we obtain after some routine arguments that the zero order terms u_i, u_e, v, w are independent of the fast variable y and satisfy the (macroscopic) bidomain model [14, 22, 34]

$$(1.4) \quad \begin{cases} |\Gamma| \partial_t v - \operatorname{div}(M_i \nabla u_i) + |\Gamma| I(v, w) = |Y_i| s_i, & \text{in } (0, T) \times \Omega, \\ |\Gamma| \partial_t v + \operatorname{div}(M_e \nabla u_e) + |\Gamma| I(v, w) = -|Y_e| s_e, & \text{in } (0, T) \times \Omega, \\ \partial_t w = H(v, w), & \text{in } (0, T) \times \Omega, \end{cases}$$

where the homogenized conductivity tensors $M_i(x), M_e(x)$ are given by

$$(1.5) \quad M_j(x) = \int_{Y_j} \sigma_j(x, y) (I + \nabla_y \chi_j(x, y)) dy, \quad j = i, e,$$

and the y -periodic (vector-valued) function $\chi_j = \chi_j(x, y)$ solves the cell problem

$$(1.6) \quad \begin{cases} -\operatorname{div}_y(\sigma_j \nabla_y \chi_j) = -\operatorname{div}_y \sigma_j, & \text{in } \Omega \times Y_j, \\ \nu \cdot \sigma_j \nabla_y \chi_j = \nu \cdot \sigma_j, & \text{on } \Omega \times \Gamma. \end{cases}$$

Note that the effective potentials u_i, u_e in (1.4) are defined at every point of Ω , while in the microscopic model they live on disjoint sets $\Omega_i^\varepsilon, \Omega_e^\varepsilon$. In (1.4), (1.5), (1.6) the sets Y_i, Y_e are the intra and extracellular spaces within the reference unit cell Y , separated by the cell membrane Γ (see Section 2 for details). It is worth noting that the bidomain model is often stated in terms of the ‘‘geometric’’ parameter $\chi = \frac{|\Gamma|}{|Y|} = |\Gamma|$ representing the surface-to-volume ratio of the cardiac cells.

Regarding the existence and uniqueness of properly defined solutions, standard theory for parabolic-elliptic systems does not apply naturally to the bidomain model (1.4). A number of works [6, 7, 8, 14, 42] have recently provided well-posedness results for (1.4), applying differing solution concepts and technical frameworks.

As alluded to earlier, we will provide a rigorous derivation of the homogenized system (1.4), (1.5), (1.6) based on the theory of two-scale convergence (see [33] and [1, 2, 28]). This result is not covered by standard parabolic homogenization theory. A complication is the nonlinear dynamic boundary condition (posed on an underlying oscillating surface), which makes it difficult to pass to the limit in (1.1) as $\varepsilon \rightarrow 0$. The aim is to prove that a sequence $\{u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon\}_{\varepsilon>0}$ of solutions to the microscopic problem two-scale converges to the solution of the bidomain model (1.4). However, two-scale convergence is not ‘‘strong enough’’ to justify passing to the limit in the nonlinear boundary condition. To handle this difficulty we use the boundary unfolding operator [11], establishing strong convergence of $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ in $L^2((0, T) \times \Omega \times \Gamma)$, where $\mathcal{T}_\varepsilon^b$ denotes the boundary unfolding operator. The

boundary unfolding operator makes our proof flexible enough to handle a range of membrane models, exemplified by the generalized FitzHugh-Nagumo model.

Unfolding operators, presented and carefully analyzed in [10, 11], facilitate elementary proofs of classical homogenization results on fixed as well as perforated domains/surfaces. An unfolding operator \mathcal{T}_ε maps a function $v(x)$ defined on an oscillating domain/surface to a higher dimensional function $\mathcal{T}_\varepsilon(v)(x, y)$ on a fixed domain, to which one can apply standard convergence theorems in fixed L^p spaces. Reflecting the “two-component” nature of the cardiac domain, it makes sense to use two unfolding operators $\mathcal{T}_\varepsilon^{i,e}$, linked to the intra- and extracellular domains $\Omega_{i,e}^\varepsilon$. In this paper, however, we mainly unfold functions defined on the cell membrane, utilizing the boundary unfolding operator $\mathcal{T}_\varepsilon^b$.

For somewhat similar unfolding of “two-component” domains separated by a periodic boundary, see [15, 16, 43]. For other relevant works that combine two-scale convergence and unfolding methods, we refer to [19, 18, 20, 29, 32]. Among these, our work borrows ideas mostly from [19, 18, 32].

The remaining part of the paper is organized as follows: In Section 2, we collect relevant functional spaces and analysis results. Moreover, we gather definitions and tools linked to two-scale convergence and unfolding operators. In Section 3, we define precisely what is meant by a weak solution of the microscopic problem (1.1), state a well-posedness result, and establish several “ ε -independent” a priori estimates. The main homogenization result is stated and proved in Section 4.

2. PRELIMINARIES

2.1. Some functional spaces and tools. For a general review of integer and fractional order Sobolev spaces (on Lipschitz domains) and relevant analysis tools, see [9, Chaps. 2 & 3] and [30, Chap. 3]. For relevant background material on mathematical homogenization, we refer to [12].

Let $\Omega \subset \mathbb{R}^3$ be a bounded open set with Lipschitz boundary. We denote by $C_0^\infty(\Omega)$ the (infinitely) smooth functions with compact support in Ω . The space of smooth Y -periodic functions is denoted by $C_{\text{per}}^\infty(Y)$. The closure of this space under the norm $\|\nabla(\cdot)\|_{L^2(Y)}$ is denoted by $H_{\text{per}}^1(Y)$. We write H^s for the L^2 -based Sobolev spaces $W^{s,2}$ ($s \in (0, 1]$).

We make use of Sobolev spaces on surfaces, as defined for example in [27, p. 34] and [30, p. 96]. Specifically, we use the (Hilbert) space $H^{1/2}(\Gamma)$, for a two-dimensional Lipschitz surface $\Gamma \subset \Omega$, equipped with the norm

$$\|u\|_{H^{1/2}(\Gamma)}^2 = \|u\|_{L^2(\Gamma)}^2 + |u|_{H_0^{1/2}(\Gamma)}^2,$$

where

$$|u|_{H_0^{1/2}}^2 = \int_\Gamma \int_\Gamma \frac{|u(x) - u(x')|^2}{|x - x'|^3} dS(x) dS(x'),$$

and dS is the two-dimensional surface measure. We define the dual space of $H^{1/2}(\Gamma)$ as $H^{-1/2}(\Gamma) := (H^{1/2}(\Gamma))^*$, equipped with the norm of dual spaces

$$\|u\|_{H^{-1/2}(\Gamma)} := \sup_{\substack{\phi \in H^{1/2}(\Gamma) \\ \|\phi\|_{H^{1/2}(\Gamma)}=1}} \langle u, \phi \rangle_{H^{-1/2}(\Gamma), H^{1/2}(\Gamma)}.$$

The following trace inequality holds:

$$(2.1) \quad \|u|_\Gamma\|_{H^{1/2}(\Gamma)} \leq C \|u\|_{H^1(\Omega)}, \quad u \in H^1(\Omega).$$

Any function in $H^{1/2}(\Gamma)$ can be characterized as the trace of a function in $H^1(\Omega)$. The trace map has a continuous right inverse $\mathcal{I} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$, satisfying

$$(2.2) \quad \|\mathcal{I}(u)\|_{H^1(\Omega)} \leq C \|u\|_{H^{1/2}(\Gamma)}, \quad \forall u \in H^{1/2}(\Gamma),$$

where the constant C depends only on Γ . We need the Sobolev inequality

$$(2.3) \quad \|u\|_{L^4(\Gamma)} \leq C \|u\|_{H^{1/2}(\Gamma)}.$$

Indeed, $H^{1/2}(\Gamma)$ is continuously embedded in $L^p(\Gamma)$ for $p \in [1, 4]$. This embedding is compact for $p \in [1, 4)$. In particular, $H^{1/2}(\Gamma)$ is compactly embedded in $L^2(\Gamma)$.

Let X be a separable Banach space X and $p \in [1, \infty]$. We make routinely use of Lebesgue-Bochner spaces such as $L^p(\Omega; X)$ and $L^p(0, T; X)$. We also use the spaces of continuous functions from Ω to X and $(0, T)$ to X , denoted by $C(\Omega; X)$ and $C(0, T; X)$, respectively, and the similar spaces with C replaced by C^p or C_0^p . If X is a Banach space, then X/\mathbb{R} denotes the space consisting of classes of functions in X that are equal up to an additive constant.

Recall that $H^{1/2}(\Gamma)$ is a Hilbert space embedded in a continuous and dense way in $L^2(\Gamma)$. The (Lions-Magenes) integration-by-parts formula holds for functions u_1, u_2 that belong to the Banach space

$$\mathcal{V}_{\Gamma, T} = \left\{ u \in L^2(0, T; H^{1/2}(\Gamma)) \cap L^4((0, T) \times \Gamma) \mid \partial_t u \in L^2(0, T; H^{-1/2}(\Gamma)) + L^{4/3}((0, T) \times \Gamma) \right\},$$

equipped with the norm

$$\|u\|_{\mathcal{V}_{\Gamma, T}} = \|u\|_{L^2(0, T; H^{1/2}(\Gamma)) \cap L^4((0, T) \times \Gamma)} + \|\partial_t u\|_{L^2(0, T; H^{-1/2}(\Gamma)) + L^{4/3}((0, T) \times \Gamma)},$$

where $\|u\|_{X_1 \cap X_2} = \max(\|u\|_{X_1}, \|u\|_{X_2})$, $\|u\|_{X_1 + X_2} = \inf_{u=u_1+u_2} (\|u_1\|_{X_1} + \|u_2\|_{X_2})$. Indeed, for $u_1, u_2 \in \mathcal{V}_{\Gamma, T}$, $[0, T] \ni t \mapsto (u_1(t), u_2(t))_{L^2(\Gamma)}$ is continuous and

$$(2.4) \quad \int_{t_1}^{t_2} \langle \partial_t u_1, u_2 \rangle dt + \int_{t_1}^{t_2} \langle \partial_t u_2, u_1 \rangle dt = (u_1(t_2), u_2(t_2))_{L^2(\Gamma)} - (u_1(t_1), u_2(t_1))_{L^2(\Gamma)},$$

for all $t_1, t_2 \in [0, T]$, $t_1 < t_2$, where $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $H^{-1/2}(\Gamma) + L^{4/3}(\Gamma)$ and $H^{1/2}(\Gamma) \cap L^4(\Gamma)$. For a proof of (2.4) that can be adapted to our situation, see e.g. [9, p. 99].

Taking $u_1 = u_2 = u \in \mathcal{V}_{\Gamma, T}$, we obtain the chain rule

$$(2.5) \quad \int_{t_1}^{t_2} \langle \partial_t u, u \rangle dt = \frac{1}{2} \|u(t_2)\|_{L^2(\Gamma)}^2 - \frac{1}{2} \|u(t_1)\|_{L^2(\Gamma)}^2.$$

Adapting standard arguments (see e.g. [9, p. 101]), the embedding

$$(2.6) \quad \mathcal{V}_{\Gamma, T} \hookrightarrow C(0, T; L^2(\Gamma))$$

is continuous. Indeed, this result follows from the continuity of the squared norm $t \mapsto \|u(t)\|_{L^2(\Gamma)}^2$ (see above) and the weak continuity of u in $L^2(\Gamma)$. The latter results from an easily obtained bound on u in $L^\infty(0, T; L^2(\Gamma))$ and the continuity of u in $H^{-1/2}(\Gamma)$, both facts being deducible from (2.4).

Let us dwell a bit further on the time continuity of functions in $\mathcal{V}_{\Gamma, T}$. By (2.3), $H^{1/2} \subset L^4(\Gamma)$ and so $L^{4/3}(\Gamma) \subset H^{-1/2}(\Gamma)$. Therefore,

$$L^2(0, T; H^{-1/2}(\Gamma)) + L^{4/3}((0, T) \times \Gamma) \subset L^{4/3}(0, T; H^{-1/2}(\Gamma)).$$

With $u_1 = u \in \mathcal{V}_{\Gamma, T}$ and $u_2 = \phi \in H^{1/2}(\Gamma)$ in (2.4), it follows that

$$\begin{aligned} (u(t_2) - u(t_1), \phi)_{L^2(\Gamma)} &= \int_{t_1}^{t_2} \langle \partial_t u, \phi \rangle dt \\ &\leq \|\phi\|_{H^{1/2}(\Gamma)} \int_{t_1}^{t_2} \|\partial_t u\|_{H^{-1/2}(\Gamma)} dt \\ &\leq \|\phi\|_{H^{1/2}(\Gamma)} \|\partial_t u\|_{L^{4/3}(t_1, t_2; H^{-1/2}(\Gamma))} (t_2 - t_1)^{1/4}. \end{aligned}$$

Fix a small shift $\Delta_t > 0$. Specifying $t_1 = t \in (0, T - \Delta_t)$, $t_2 = t + \Delta_t$, and $\phi = u(t + \Delta_t, \cdot) - u(t, \cdot)$ gives

$$\begin{aligned} \|u(t + \Delta_t, \cdot) - u(t, \cdot)\|_{L^2(\Gamma)}^2 &\leq \|u(t + \Delta_t, \cdot) - u(t, \cdot)\|_{H^{1/2}(\Gamma)} \|\partial_t u\|_{L^{4/3}(0, T; H^{-1/2}(\Gamma))} \Delta_t^{1/4}. \end{aligned}$$

Integrating this inequality over $t \in (0, T - \Delta_t)$, accompanied by a few elementary manipulations, results in the temporal translation estimate

$$\begin{aligned} \|u(\cdot + \Delta_t, \cdot) - u(\cdot, \cdot)\|_{L^2(0, T - \Delta_t; L^2(\Gamma))} &\leq C_T \|u\|_{L^2(0, T; H^{1/2}(\Gamma))}^{1/2} \|\partial_t u\|_{L^{4/3}(0, T; H^{-1/2}(\Gamma))}^{1/2} \Delta_t^{1/8}, \quad u \in \mathcal{V}_{\Gamma, T}, \end{aligned}$$

where $C_T = 2^{1/2} T^{1/4}$. A similar estimate holds for negative Δ_t .

There is a compact embedding of $\mathcal{V}_{\Gamma, T}$ in $L^2(0, T; L^2(\Gamma))$. As pointed out above, $\mathcal{V}_{\Gamma, T}$ is a subset of $\{u \in L^2(0, T; H^{1/2}(\Gamma)) : \partial_t u \in L^{4/3}(0, T; H^{-1/2}(\Gamma))\}$, which is compactly embedded in $L^2(0, T; L^2(\Gamma))$ by the Aubin-Lions theorem.

We need a generalization of this result due to Simon [37]. Given two Banach spaces $X_1 \subset X_0$, with X_1 compactly embedded in X_0 , let \mathcal{K} be a collection of functions in $L^p(0, T; X_0)$, $p \in [1, \infty]$. The work [37] supplies several results ensuring the compactness of \mathcal{K} in $L^p(0, T; X_0)$ (in $C([0, T]; X_0)$ if $p = \infty$). For example, we can assume that \mathcal{K} is bounded in $L^1_{\text{loc}}(0, T; X_1)$ and

$$\|u(\cdot + \Delta_t) - u\|_{L^p(0, T - \Delta_t; X_0)} \rightarrow 0 \quad \text{as } \Delta_t \rightarrow 0, \text{ uniformly for } u \in \mathcal{K},$$

cf. [37, Theorem 3]. We apply this result with $p = 2$, $X_1 = H^{1/2}(\Gamma^\varepsilon)$, $X_0 = L^2(\Gamma^\varepsilon)$. Another result involves a third Banach space X_{-1} (e.g. $X_{-1} = H^{-1/2}(\Gamma^\varepsilon)$), such that $X_1 \subset X_0 \subset X_{-1}$ and X_1 is compactly embedded in X_0 . Compactness of \mathcal{K} in $L^p(0, T; X_0)$ follows if the set \mathcal{K} is bounded in $L^p(0, T; X_1)$ and, as $\Delta_t \rightarrow 0$, $\|u(\cdot + \Delta_t) - u\|_{L^p(0, T - \Delta_t; X_{-1})} \rightarrow 0$, uniformly for $u \in \mathcal{K}$ [37, Theorem 5].

2.2. Two-scale convergence. Recall that $\Omega \subset \mathbb{R}^3$ denotes the entire (connected, bounded, open) cardiac domain, assumed to be of the form (1.2). The assumption (1.2) simplifies the presentation. With mild modifications of the upcoming proofs, the results remain valid for general domains with Lipschitz boundary. Let Y be a reference unit cell in \mathbb{R}^3 , which we fix to be the unit cube $Y := [0, 1]^3$.

Let Y_i and Y_e be the (disjoint, connected, open) intra and extracellular spaces within Y , separated by the cell membrane Γ :

$$\overline{Y_i} \cup \overline{Y_e} = Y, \quad \Gamma = \partial Y_i \setminus \partial Y.$$

Denote by K^ε the set of $k \in \mathbb{Z}^3$ for which $\bigcup_{k \in K^\varepsilon} \varepsilon(k + Y) = \overline{\Omega}$. We define the intracellular domain Ω_i^ε , the extracellular domain Ω_e^ε , and the cell membrane Γ^ε as

$$(2.7) \quad \Omega_j^\varepsilon = \bigcup_{k \in K^\varepsilon} \varepsilon(k + Y_j), \quad j = i, e, \quad \Gamma^\varepsilon = \bigcup_{k \in K^\varepsilon} \varepsilon(k + \Gamma).$$

Both sets $\Omega_i^\varepsilon, \Omega_e^\varepsilon$ are connected Lipschitz domains, see Figure 1.1. Note however that it is impossible to have both Ω_i^ε and Ω_e^ε connected in a two-dimensional picture.




FIGURE 1. The rescaled sets Ω_i^ε , Ω_e^ε , Γ^ε (left) and the unit cell Y (right).

To derive estimates for the microscopic model, we employ the following trace inequality for ε -periodic hypersurfaces:

$$(2.8) \quad \varepsilon \|u|_{\Gamma^\varepsilon}\|_{L^2(\Gamma^\varepsilon)}^2 \leq C \left(\|u\|_{L^2(\Omega_j^\varepsilon)}^2 + \varepsilon^2 \|\nabla u\|_{L^2(\Omega_j^\varepsilon)}^2 \right), \quad u \in H^1(\Omega_j^\varepsilon), \quad j = i, e,$$

for some constant C independent of ε , cf. [24, Lemma 3] or [29, Lemma 4.2].

We need a uniform Poincaré inequality for perforated domains [11].

Lemma 2.1. *There exists a constant C , independent of $\varepsilon > 0$, such that*

$$(2.9) \quad \left\| u - \frac{1}{|\Omega_j^\varepsilon|} \int_{\Omega_j^\varepsilon} u \, dx \right\|_{L^2(\Omega_j^\varepsilon)} \leq C \|\nabla u\|_{L^2(\Omega_j^\varepsilon)},$$

for all $u \in H^1(\Omega_j^\varepsilon)$, $j = i, e$.

Estimate (2.9) holds under mild regularity assumptions on the perforated domains; a Lipschitz boundary is more than sufficient (but connectedness is essential).

Recall that a sequence $\{u^\varepsilon\}_{\varepsilon>0} \subset L^2((0, T) \times \Omega)$ two-scale converges to u in $L^2((0, T) \times \Omega; L^2_{\text{per}}(Y))$ if

$$(2.10) \quad \int_0^T \int_{\Omega} u^\varepsilon(t, x) \varphi\left(t, x, \frac{x}{\varepsilon}\right) dx dt \xrightarrow{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega} \int_Y u(t, x, y) \varphi(t, x, y) dy dx dt,$$

for all $\varphi \in C([0, T] \times \bar{\Omega}; C_{\text{per}}(Y))$. We express this symbolically as

$$u^\varepsilon \xrightarrow{2} u.$$

By density properties, the convergence (2.10) also holds for test functions φ from $L^2((0, T) \times \Omega; C_{\text{per}}(Y))$ [12, p. 176].

The two-scale compactness theorem [1, 12] is of fundamental importance.

Theorem 2.2 (two-scale compactness). *Let $\{u^\varepsilon\}_{\varepsilon>0}$ be a bounded sequence in $L^2((0, T) \times \Omega)$, $\|u^\varepsilon\|_{L^2((0, T) \times \Omega)} \leq C \forall \varepsilon > 0$. Then there exist a subsequence $\varepsilon_n \rightarrow 0$ and a function $u \in L^2((0, T) \times \Omega; L^2(Y))$ such that u^{ε_n} two-scale converges to u as $n \rightarrow \infty$.*

Consider a sequence $\{u_j^\varepsilon\}_{\varepsilon>0}$ of functions defined on the perforated domain $(0, T) \times \Omega_j^\varepsilon$, $j = i, e$. We write $\widetilde{u}_j^\varepsilon$ for the zero-extension of u_j^ε to $(0, T) \times \Omega$:

$$\widetilde{u}_j^\varepsilon(t, x) := \begin{cases} u_j^\varepsilon(t, x) & \text{if } (t, x) \in (0, T) \times \Omega_j^\varepsilon, \\ 0 & \text{if } (t, x) \in (0, T) \times (\Omega \setminus \Omega_j^\varepsilon). \end{cases}$$

By Theorem 2.2, $\{\widetilde{u}_j^\varepsilon\}_{\varepsilon>0}$ has a two-scale convergent subsequence, provided we know that $\|u_j^\varepsilon\|_{L^2((0, T) \times \Omega_j^\varepsilon)} \leq C \forall \varepsilon > 0$. However, this is not true in general for the

gradient of $\widetilde{u}_j^\varepsilon$, even if $\|u_j^\varepsilon\|_{L^2(0,T;H^1(\Omega_j^\varepsilon))} \leq C$, since the extension by zero creates a discontinuity across Γ^ε . Instead the following statement holds true for the gradient:

Lemma 2.3. *Fix $j \in \{i, e\}$ and suppose $u^\varepsilon = u_j^\varepsilon$ satisfies $\|u^\varepsilon\|_{L^2(0,T;H^1(\Omega_j^\varepsilon))} \leq C$ $\forall \varepsilon > 0$. Then there exist a subsequence $\varepsilon_n \rightarrow 0$ and functions $u \in L^2(0,T;H^1(\Omega))$, $u_1 \in L^2((0,T) \times \Omega; H_{\text{per}}^1(Y_j))$ such that as $n \rightarrow \infty$,*

$$\begin{aligned} \widetilde{u}^{\varepsilon_n} &\xrightarrow{2} \mathbb{1}_{Y_j}(y)u(t,x) \text{ in } L^2((0,T) \times \Omega; L_{\text{per}}^2(Y)), \\ \widetilde{\nabla u}^{\varepsilon_n} &\xrightarrow{2} \mathbb{1}_{Y_j}(y)(\nabla_x u(t,x) + \nabla_y u_1(t,x,y)) \text{ in } L^2((0,T) \times \Omega; L_{\text{per}}^2(Y)). \end{aligned}$$

Here, $\mathbb{1}_{Y_j}(y)$ denotes the characteristic function of Y_j ,

$$\mathbb{1}_j(y) = \begin{cases} 1 & \text{if } y \in Y_j, \\ 0 & \text{if } y \notin Y_j. \end{cases}$$

For a proof of this lemma in the time independent case, see [1, Theorem 2.9]. The extension to time dependent functions is straightforward.

There is an extension of two-scale convergence to periodic surfaces [2]. Recall that a periodic surface Γ^ε is given by

$$\Gamma^\varepsilon := \left\{ x \in \Omega \mid \frac{x}{\varepsilon} \in k + \Gamma \text{ for some } k \in \mathbb{Z}^3 \right\},$$

where $\Gamma \subset Y$ is a surface in the unit cell. Since $|\Gamma^\varepsilon| \sim \varepsilon^{-1}$, it is necessary to introduce a normalizing factor in the definition of two-scale convergence on surfaces.

A sequence $\{v^\varepsilon\}_{\varepsilon>0}$ of functions in $L^2((0,T) \times \Gamma^\varepsilon)$ *two-scale converges* to v in $L^2((0,T) \times \Omega; L^2(\Gamma))$, written

$$v^\varepsilon \xrightarrow{2-S} v,$$

if, for all $\varphi \in C([0,T] \times \overline{\Omega}; C_{\text{per}}(\Gamma))$,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} v^\varepsilon(t,x) \varphi\left(t,x,\frac{x}{\varepsilon}\right) dS(x) dt \\ = \int_0^T \int_\Omega \int_\Gamma v(t,x,y) \varphi(t,x,y) dS(y) dx dt. \end{aligned}$$

As with (2.10), this convergence continues to hold for test functions φ that belong to $L^2((0,T) \times \Omega; C_{\text{per}}(\Gamma))$.

There is a version of Theorem 2.2 for functions on periodic surfaces [2].

Theorem 2.4 (two-scale compactness on surfaces). *Suppose $\{v^\varepsilon\}_{\varepsilon>0}$ is a sequence of functions in $L^2((0,T) \times \Gamma_\varepsilon)$ satisfying*

$$(2.11) \quad \varepsilon \int_0^T \int_{\Gamma^\varepsilon} |v^\varepsilon|^2 dS dt \leq C,$$

for some function C that is independent of $\varepsilon > 0$. Then there exist a subsequence $\varepsilon_n \rightarrow 0$ and a function $v \in L^2((0,T) \times \Omega; L^2(\Gamma))$ such that as $n \rightarrow \infty$,

$$v^{\varepsilon_n} \xrightarrow{2-S} v.$$

One can characterize the two-scale limit of traces of bounded sequences in $L^2(0,T;H^1(\Omega^\varepsilon))$ as the trace of the two-scale limit [2].

Lemma 2.5. *Fix $j \in \{i, e\}$. Suppose $u^\varepsilon = u_j^\varepsilon$ satisfies $\|u^\varepsilon\|_{L^2(0,T;H^1(\Omega_j^\varepsilon))} \leq C$ $\forall \varepsilon > 0$ and, cf. Lemma 2.3, $\widetilde{u}^\varepsilon \xrightarrow{2} \mathbb{1}_{Y_j}(y)u(t,x)$ in $L^2((0,T) \times \Omega; L^2(Y))$. Let*

$$g^\varepsilon := u^\varepsilon|_{\Gamma^\varepsilon} \in L^2((0,T) \times \Gamma^\varepsilon)$$

be the trace of u^ε on $\Gamma^\varepsilon = \partial\Omega_j^\varepsilon \setminus \partial\Omega$. Then, up to a subsequence,

$$g^\varepsilon \xrightarrow{2-S} g := \mathbb{1}_\Gamma(y)u(t, x).$$

Remark 2.6. In view of Lemma 2.5, we have (in the sense of measures)

$$\varepsilon u^\varepsilon|_{\Gamma^\varepsilon} dS dt \xrightarrow{*} |\Gamma| u dx dt, \quad \widetilde{u}^\varepsilon dx dt \xrightarrow{*} |Y_j| u dx dt.$$

2.3. Unfolding operators. An alternative approach to studying convergence on oscillating surfaces Γ^ε is provided by the boundary unfolding operator [11]. For any $x \in \mathbb{R}^3$, we have the decomposition $x = [x] + \{x\}$, where $[x] \in \mathbb{Z}^3$ and $\{x\} \in [0, 1]^3$ denotes the integer and fractional parts of x , respectively. For later use, note the following simple properties, which hold for any $x, \bar{x} \in \mathbb{R}^3$, $n \in \mathbb{Z}^3$:

$$[x + n] = [x] + n, \quad \{x + n\} = \{x\}, \quad [x + \bar{x}] \leq [x] + [\bar{x}] + (1, 1, 1).$$

Applying the above decomposition to x/ε gives

$$x = \varepsilon ([x/\varepsilon] + \{x/\varepsilon\}),$$

where $[x/\varepsilon] \in \mathbb{Z}^3$, $\{x/\varepsilon\} \in [0, 1]^3$.

The *boundary unfolding operator* $\mathcal{T}_\varepsilon^b$ is defined by

$$(2.12) \quad \begin{aligned} \mathcal{T}_\varepsilon^b : L^2(0, T; L^2(\Gamma^\varepsilon)) &\rightarrow L^2(0, T; L^2(\Omega \times \Gamma)), \\ \mathcal{T}_\varepsilon^b(v)(t, x, y) &= v\left(t, \varepsilon \left[\frac{x}{\varepsilon} \right] + \varepsilon y\right), \quad (t, x, y) \in (0, T) \times \Omega \times \Gamma. \end{aligned}$$

The advantage of the unfolding operator is that we can formulate questions of convergence in a fixed space $L^2(0, T; L^2(\Omega \times \Gamma))$. All definitions and results in this section are formulated in L^2 spaces. Everything remains the same, however, if we replace L^2 by L^p for any $p \in [1, \infty)$. We refer to [11] for the definition of the boundary unfolding operator and proofs of the properties listed next.

The boundary unfolding operator $\mathcal{T}_\varepsilon^b$ is bounded, linear, and satisfies

$$(2.13) \quad \mathcal{T}_\varepsilon^b(v_1 v_2) = \mathcal{T}_\varepsilon^b(v_1) \mathcal{T}_\varepsilon^b(v_2), \quad v_1, v_2 \in L^2(0, T; L^2(\Gamma^\varepsilon)).$$

For any Y -periodic function $\psi \in L^2(\Gamma)$, set $\psi_\varepsilon(x) := \psi(x/\varepsilon)$. Then

$$\mathcal{T}_\varepsilon^b(\psi_\varepsilon)(x, y) = \psi(y), \quad x \in \Omega, y \in \Gamma.$$

For $v \in L^2(0, T; L^2(\Gamma^\varepsilon))$, we have the integration formula

$$(2.14) \quad \varepsilon \int_{\Gamma^\varepsilon} v(t, x) dS(x) = \int_\Omega \int_\Gamma \mathcal{T}_\varepsilon^b(v)(t, x, y) dS(y) dx,$$

for a.e. $t \in (0, T)$, thereby converting an integral over the oscillating set Γ^ε to an integral over the fixed set $\Omega \times \Gamma$. For $v \in L^2(0, T; L^2(\Gamma^\varepsilon))$,

$$(2.15) \quad \|\mathcal{T}_\varepsilon^b(v)\|_{L^2(\Omega \times \Gamma)} = \varepsilon^{1/2} \|v\|_{L^2(\Gamma^\varepsilon)},$$

for a.e. $t \in (0, T)$. For any $v \in L^2(0, T; L^2(\Gamma^\varepsilon))$,

$$(2.16) \quad \mathcal{T}_\varepsilon^b(v) \xrightarrow{\varepsilon \downarrow 0} v \quad \text{in } L^2(\Omega \times \Gamma),$$

for a.e. $t \in (0, T)$, and also in $L^2(0, T; L^2(\Omega \times \Gamma))$. Suppose $\{v^\varepsilon\}_{\varepsilon > 0}$ is a sequence of functions in $L^2((0, T) \times \Gamma^\varepsilon)$ satisfying (2.11). Then

$$(2.17) \quad v^\varepsilon \xrightarrow{2-S} v \iff \mathcal{T}_\varepsilon^b(v^\varepsilon) \rightharpoonup v \quad \text{in } L^2((0, T) \times \Omega \times \Gamma).$$

We need also the *unfolding operators* linked to the domains $\Omega_i^\varepsilon, \Omega_e^\varepsilon$ [11]:

$$\begin{aligned} \mathcal{T}_\varepsilon^j : L^2((0, T) \times \Omega_j^\varepsilon) &\rightarrow L^2(0, T; L^2(\Omega \times Y_j)), \quad j = i, e, \\ \mathcal{T}_\varepsilon^j(u)(t, x, y) &= u\left(t, \varepsilon \left[\frac{x}{\varepsilon} \right] + \varepsilon y\right), \quad (t, x, y) \in (0, T) \times \Omega \times Y_j. \end{aligned}$$

The unfolding operator $\mathcal{T}_\varepsilon^j$ maps functions defined on the oscillating set $(0, T) \times \Omega_j^\varepsilon$ into functions defined on the fixed domain $(0, T) \times \Omega \times Y_j$. The operator $\mathcal{T}_\varepsilon^j$ is bounded, linear and satisfies

$$\mathcal{T}_\varepsilon^j(uv) = \mathcal{T}_\varepsilon^j(u)\mathcal{T}_\varepsilon^j(v), \quad u, v \in L^2((0, T) \times \Omega_j^\varepsilon).$$

For any Y -periodic function $\psi \in L^2(Y_j)$, set $\psi_\varepsilon(x) := \psi(x/\varepsilon)$. Then

$$\mathcal{T}_\varepsilon^j(\psi_\varepsilon)(x, y) = \psi(y), \quad x \in \Omega, y \in Y_j.$$

For $u \in L^2(0, T; L^2(\Omega_j^\varepsilon))$, we have the integration formula

$$\int_{\Omega_j^\varepsilon} u(t, x) dx dt = \int_{\Omega} \int_{Y_j} \mathcal{T}_\varepsilon^j(u)(t, x, y) dy dx dt,$$

for a.e. $t \in (0, T)$. The integration formula implies

$$(2.18) \quad \|\mathcal{T}_\varepsilon^j(u)\|_{L^2(\Omega \times Y_j)} = \|u\|_{L^2(\Omega_j^\varepsilon)},$$

for a.e. $t \in (0, T)$. Let $u \in L^2(0, T; H^1(\Omega_j^\varepsilon))$. Then

$$\nabla_y \mathcal{T}_\varepsilon^j(u) = \varepsilon \mathcal{T}_\varepsilon^j(\nabla u), \quad \text{a.e. in } (0, T) \times \Omega \times Y_j,$$

and hence $\mathcal{T}_\varepsilon^j(u) \in L^2(\Omega; H^1(Y_j))$, for a.e. $t \in (0, T)$:

$$(2.19) \quad \|\nabla_y \mathcal{T}_\varepsilon^j(u)\|_{L^2(\Omega \times Y_j)} = \varepsilon \|\nabla u\|_{L^2(\Omega_j^\varepsilon)}.$$

The unfolding operators $\mathcal{T}_\varepsilon^b$ and $\mathcal{T}_\varepsilon^j$ are related in the following sense:

$$(2.20) \quad \mathcal{T}_\varepsilon^b(u|_{\Gamma^\varepsilon}) = \mathcal{T}_\varepsilon^j(u)|_\Gamma, \quad u \in L^2(0, T; H^1(\Omega_j^\varepsilon)), \quad j = i, e,$$

for a.e. $t \in (0, T)$. Combining (2.20), (2.18), (2.19), and the trace inequality (2.1) in $H^1(\Omega_j^\varepsilon)$, we obtain

$$(2.21) \quad \begin{aligned} & \|\mathcal{T}_\varepsilon^b(u|_{\Gamma^\varepsilon})\|_{L^2(\Omega; H^{1/2}(\Gamma))}^2 \\ & \leq C \left(\|\mathcal{T}_\varepsilon^j(u)\|_{L^2(\Omega; L^2(Y_j))}^2 + \|\nabla_y \mathcal{T}_\varepsilon^j(u)\|_{L^2(\Omega; L^2(Y_j))}^2 \right) \\ & = C \left(\|u\|_{L^2(\Omega_j^\varepsilon)}^2 + \varepsilon^2 \|\nabla u\|_{L^2(\Omega_j^\varepsilon)}^2 \right), \quad j = i, e, \end{aligned}$$

for a.e. $t \in (0, T)$, where the constant C is independent of ε and t . Whenever it is convenient, we will write $\mathcal{T}_\varepsilon^b(u)$ instead of $\mathcal{T}_\varepsilon^b(u|_{\Gamma^\varepsilon})$.

Next, we consider the local average (mean in the cells) operator

$$\mathcal{M}_\varepsilon^j(u)(t, x) = \int_{Y_j} \mathcal{T}_\varepsilon^j(u)(t, x, y) dy, \quad (t, x) \in (0, T) \times \Omega, \quad j = i, e,$$

and the piecewise linear interpolation operator [11, Definition 2.5]

$$(2.22) \quad \begin{aligned} & Q_\varepsilon^j : L^2(0, T; H^1(\Omega_j^\varepsilon)) \rightarrow L^2(0, T; H^1(\Omega)), \quad j = i, e, \\ & Q_\varepsilon^j(u) \text{ is the } Q_1\text{-interpolation (in } x\text{) of } \mathcal{M}_\varepsilon^j(u). \end{aligned}$$

Given the Lipschitz regularity of Y_j , the interpolation operator Q_ε^j satisfies the following estimates [11, Propositions 2.7 and 2.8]:

$$(2.23) \quad \begin{aligned} & \|Q_\varepsilon^j(u) - u\|_{L^2((0, T) \times \Omega_j^\varepsilon)} \leq C\varepsilon \|\nabla u\|_{L^2((0, T) \times \Omega_j^\varepsilon)}, \\ & \|\nabla Q_\varepsilon^j(u)\|_{L^2((0, T) \times \Omega_j^\varepsilon)} \leq C \|\nabla u\|_{L^2((0, T) \times \Omega_j^\varepsilon)}, \end{aligned}$$

where C is a constant that is independent of ε .

3. MICROSCOPIC BIDOMAIN MODEL

In this section we present a relevant notion of (weak) solution for the microscopic problem (1.1), (1.3), along with an accompanying existence theorem. We also derive some “ ε -independent” a priori estimates, which are used later to extract two-scale convergent subsequences.

3.1. Assumptions on the data. We impose the following set of assumptions on the “membrane” functions I, H :

- Generalized FitzHugh-Nagumo model: For $v, w \in \mathbb{R}$,

$$\begin{aligned} I(v, w) &= I_1(v) + I_2(v)w, & H(v, w) &= h(v) + c_{H,1}w, \\ \text{(GFHN)} \quad & \text{where } I_1, I_2, h \in C^1(\mathbb{R}), c_{H,1} \in \mathbb{R}, \text{ and} \\ & |I_1(v)| \leq c_{I,1} (1 + |v|^3), & I_1(v)v &\geq c_I |v|^4 - c_{I,2} |v|^2, \\ & I_2(v) = c_{I,3} + c_{I,4}v, & |h(v)| &\leq c_{H,2} (1 + |v|^2), \end{aligned}$$

for some constants $c_I > 0$ and $c_{I,1}, c_{I,2}, c_{I,3}, c_{I,4}, c_{H,2} \geq 0$.

The classical FitzHugh-Nagumo model corresponds to

$$(3.1) \quad I(v, w) = v(v - a)(v - 1) + w, \quad H(v, w) = \varepsilon(kv - w),$$

where $a \in (0, 1)$ and $k, \varepsilon > 0$ are constants.

Repeated applications of Cauchy’s inequality yields

$$(3.2) \quad vI(v, w) - wH(v, w) \geq \gamma |v|^4 - \beta (|v|^2 + |w|^2),$$

for some constants $\gamma > 0$ and $\beta \geq 0$. This inequality will be used to bound the transmembrane potential in the L^4 norm.

Consider a quadratic matrix A , which always can be written as the sum of its symmetric part $\frac{1}{2}(A + A^\top)$ and its skew-symmetric part $\frac{1}{2}(A - A^\top)$. Recall that in a quadratic form $z \mapsto Az \cdot z$ the skew-symmetric part does not contribute. Therefore, letting λ_{\min} and λ_{\max} denote respectively the minimum and maximum eigenvalues of the symmetric part of A , we have

$$\lambda_{\min} |z|^2 \leq Az \cdot z \leq \lambda_{\max} |z|^2, \quad \forall z.$$

For $\mu > 0$, consider the function $F^\mu : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$F^\mu(z) = \begin{pmatrix} \mu I(z) \\ -H(z) \end{pmatrix}.$$

Denote by $\lambda_{\min}^\mu(z)$, $\lambda_{\max}^\mu(z)$ the minimum, maximum eigenvalues of the symmetric part of the matrix $\nabla F^\mu(z)$. To ensure that weak solutions are unique, we need an additional assumption on I, H expressed via F^μ [8, p. 479]: $\exists \mu, \lambda > 0$ such that

$$(3.3) \quad \lambda_{\min}^\mu(z) \geq \lambda, \quad \forall z \in \mathbb{R}^2.$$

One can verify that the FitzHugh-Nagumo model (3.1) obeys (3.3) (with $\mu = \varepsilon k$).

A consequence of (3.3) is that $\nabla F^\mu(\bar{z})z \cdot z \geq \lambda |z|^2$ for all $\bar{z}, z \in \mathbb{R}^2$. Therefore, writing

$$F^\mu(z_2) - F^\mu(z_1) = \int_0^1 \nabla F^\mu(\theta z_2 + (1 - \theta)z_1)(z_2 - z_1) d\theta,$$

it follows that

$$(F^\mu(z_2) - F^\mu(z_1)) \cdot (z_2 - z_1) \geq -\lambda |z_2 - z_1|^2.$$

More explicitly, assumption (3.3) implies the following “dissipative structure” on a suitable linear combination of I and H :

$$\begin{aligned} & \mu (I(v_2, w_2) - I(v_1, w_2)) (v_2 - v_1) - (H(v_2, w_2) - H(v_1, w_1)) (w_2 - w_1) \\ & \geq -\lambda (|v_2 - v_1|^2 + |w_2 - w_1|^2), \quad \forall v_1, v_2, w_1, w_2 \in \mathbb{R}. \end{aligned}$$

This inequality implies the L^2 stability (and thus uniqueness) of weak solutions.

Remark 3.1. *There are many membrane models of cardiac cells [13, 38]. We utilize the FitzHugh-Nagumo model [17], which is a simplification of the Hodgkin-Huxley model of voltage-gated ion channels. It is possible to treat other membrane models by blending the arguments used herein with those found in [7, 8, 13, 38, 41, 42].*

As a natural assumption for homogenization, we assume that the ε -dependence of the conductivities σ_j^ε ($j = i, e$), the applied currents s_j^ε ($j = i, e$), and the initial data $v_0^\varepsilon, w_0^\varepsilon$ decouples into a “fast” and a “slow” variable:

$$(3.4) \quad \begin{aligned} \sigma_j^\varepsilon(x) &= \sigma_j\left(x, \frac{x}{\varepsilon}\right), & s_j^\varepsilon(x) &= s_j\left(x, \frac{x}{\varepsilon}\right), \\ v_0^\varepsilon(x) &= v_0\left(x, \frac{x}{\varepsilon}\right), & w_0^\varepsilon(x) &= w_0\left(x, \frac{x}{\varepsilon}\right), \end{aligned}$$

for some fixed functions $\sigma_j(x, y), s_j(x, y), v_0(x, y), w_0(x, y)$ that are Y -periodic in the second argument.

The conductivity tensors are assumed to be bounded and continuous,

$$(3.5) \quad \sigma_j(x, y) \in L^\infty(\Omega \times Y_j) \cap C(\Omega; C_{\text{per}}(Y_j)),$$

and satisfy the usual ellipticity condition, i.e., there exists $\alpha > 0$ such that

$$(3.6) \quad \eta \cdot \sigma_j(x, y) \eta \geq \alpha |\eta|^2, \quad \forall \eta \in \mathbb{R}^3, \forall (x, y) \in \Omega \times Y_j,$$

for $j = i, e$. Finally, we assume that each σ_j is symmetric: $\sigma_j^T = \sigma_j$.

The regularity assumption (3.5) implies that σ_j is an admissible test function for two-scale convergence [1], which means that

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega_j^\varepsilon} \sigma_j^\varepsilon(x) \varphi^\varepsilon(x) dx = \int_{\Omega \times Y_j} \sigma(x, y) \varphi(x, y) dx dy, \quad j = i, e,$$

for every two-scale convergent sequence $\{\varphi^\varepsilon\}_{\varepsilon > 0}$, $\varphi^\varepsilon \xrightarrow{2} \varphi$. This convergence still holds if the second part of (3.5) is replaced by $\sigma_j \in L^2(\Omega; C_{\text{per}}(Y_j))$.

For the stimulation currents we assume the compatibility condition

$$(3.7) \quad \sum_{j=i,e} \int_{\Omega_j^\varepsilon} s_j^\varepsilon dx = 0,$$

and the boundedness in L^2 :

$$(3.8) \quad \|s_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \leq C, \quad j = i, e,$$

which, in view of (3.4), is guaranteed if we take [12, p. 174]

$$(3.9) \quad s_j \in L^2(\Omega; C_{\text{per}}(Y_j)).$$

Similarly, we assume that

$$(3.10) \quad v_0, w_0 \in L^2(\Omega; C_{\text{per}}(\Gamma)).$$

Throughout this paper we denote by C a generic constant, not depending on the parameter ε . The actual value of C may change from one line to the next.

3.2. Weak solutions. Testing (1.1) against appropriate functions $\varphi_i, \varphi_e, \varphi_w$ we obtain the weak formulation of the microscopic bidomain model (1.1), (1.3), cf. [14, 41] for details. We note that the terms involving the boundary $\partial\Omega$ vanish due to the Neumann boundary condition (1.3).

Definition 3.2 (weak formulation of microscopic system). *A weak solution to (1.1), (1.3) is a collection $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$ of functions satisfying the following conditions:*

i (algebraic relation).

$$(3.11) \quad v^\varepsilon = u_i^\varepsilon|_{\Gamma^\varepsilon} - u_e^\varepsilon|_{\Gamma^\varepsilon} \quad \text{a.e. on } (0, T) \times \Gamma^\varepsilon.$$

ii (regularity).

$$\begin{aligned} u_j^\varepsilon &\in L^2(0, T; H^1(\Omega_j^\varepsilon)), \quad j = i, e, \\ \int_{\Omega^\varepsilon} u_e^\varepsilon(t, x) dx &= 0, \quad t \in (0, T), \\ v^\varepsilon &\in L^2(0, T; H^{1/2}(\Gamma^\varepsilon)) \cap L^4((0, T) \times \Gamma^\varepsilon), \\ \partial_t v &\in L^2(0, T; H^{-1/2}(\Gamma^\varepsilon)) + L^{4/3}((0, T) \times \Gamma^\varepsilon), \\ w^\varepsilon &\in H^1(0, T; L^2(\Gamma^\varepsilon)). \end{aligned}$$

iii (initial conditions).

$$(3.12) \quad v^\varepsilon(0) = v_0^\varepsilon, \quad w^\varepsilon(0) = w_0^\varepsilon.$$

iv (differential equations).

$$(3.13) \quad \begin{aligned} \varepsilon \int_0^T \langle \partial_t v^\varepsilon, \varphi_i \rangle dt + \int_0^T \int_{\Omega_i^\varepsilon} \sigma_i^\varepsilon \nabla u_i^\varepsilon \cdot \nabla \varphi_i dx dt \\ + \varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi_i dS dt = \int_0^T \int_{\Omega_i^\varepsilon} s_i^\varepsilon \varphi_i dx dt, \end{aligned}$$

$$(3.14) \quad \begin{aligned} \varepsilon \int_0^T \langle \partial_t v^\varepsilon, \varphi_e \rangle dt - \int_0^T \int_{\Omega_e^\varepsilon} \sigma_e^\varepsilon \nabla u_e^\varepsilon \cdot \nabla \varphi_e dx dt \\ + \varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi_e dS dt = - \int_0^T \int_{\Omega_e^\varepsilon} s_e^\varepsilon \varphi_e dx dt, \end{aligned}$$

$$(3.15) \quad \int_{\Gamma^\varepsilon} \partial_t w^\varepsilon \varphi_w dx = \int_{\Gamma^\varepsilon} H(v^\varepsilon, w^\varepsilon) \varphi_w dS,$$

for all $\varphi_j \in L^2(0, T; H^1(\Omega_j^\varepsilon))$ with $\varphi_j \in L^4((0, T) \times \Gamma^\varepsilon)$ ($j = i, e$), and for all $\varphi_w \in L^2(0, T; L^2(\Gamma^\varepsilon))$.

Remark 3.3. In (3.13), (3.14) we use $\langle \cdot, \cdot \rangle$ to denote the duality pairing between $H^{-1/2}(\Gamma^\varepsilon) + L^{4/3}(\Gamma^\varepsilon)$ and $H^{1/2}(\Gamma^\varepsilon) \cap L^4(\Gamma^\varepsilon)$. For a motivation of the regularity conditions in Definition 3.2, see Remark 3.9 below.

By the embedding (2.6), $v^\varepsilon, w^\varepsilon \in C(0, T; L^2(\Gamma^\varepsilon))$ and therefore the pointwise evaluations $v(0), w(0)$ in (3.12) are well defined. The time derivative $\partial_t v^\varepsilon$ is a distribution belonging to $L^2(0, T; H^{-1/2}(\Gamma^\varepsilon)) + L^{4/3}((0, T) \times \Gamma^\varepsilon)$ with initial values v_0^ε , so that the integration-by-parts formula (2.4) holds. Consequently, we may replace

$$\varepsilon \int_0^T \langle \partial_t v^\varepsilon, \varphi_j \rangle dt, \quad j = i, e,$$

in (3.13) and (3.14) by

$$(3.16) \quad - \varepsilon \int_0^T \int_{\Gamma^\varepsilon} v^\varepsilon \partial_t \varphi_j dS(x) dt - \varepsilon \int_{\Gamma^\varepsilon} v_0^\varepsilon \varphi_j(0) dS(x) dt,$$

for all test functions $\varphi_j \in C_0^\infty([0, T] \times \Omega_j^\varepsilon)$, or for all $\varphi_j \in L^2(0, T; H^1(\Omega_j^\varepsilon))$ such that $\partial_t \varphi_j \in L^2((0, T) \times \Gamma^\varepsilon)$, $\varphi_j \in L^4((0, T) \times \Gamma^\varepsilon)$, and $\varphi_j(T) = 0$. Later, when passing to the limit $\varepsilon \rightarrow 0$ in (3.13) and (3.14), we make use of the form (3.16).

Remark 3.4. Consider a weak solution $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$ according to Definition 3.2. Thanks to (2.1), the trace of $\varphi_j \in L^2(0, T; H^1(\Omega_j^\varepsilon))$ belongs to $L^2(0, T; H^{1/2}(\Gamma^\varepsilon))$. By the Sobolev inequality (2.3), the trace of φ_j belongs also to $L^2(0, T; L^4(\Gamma^\varepsilon))$. In Definition 3.2 we ask additionally that the trace of φ_j belongs to $L^4((0, T) \times \Gamma^\varepsilon)$ to ensure that the surface terms in (3.13), (3.14) are well-defined

Indeed, $\int_0^T \langle \partial_t v^\varepsilon, \varphi_j \rangle dt$ is well-defined for such φ_j . Moreover,

$$J := \left| \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi_j dS dt \right| \leq \|I(v^\varepsilon, w^\varepsilon)\|_{L^{4/3}((0, T) \times \Gamma^\varepsilon)} \|\varphi_j\|_{L^4((0, T) \times \Gamma^\varepsilon)}.$$

For the membrane model (**GFHN**), the growth condition on I implies

$$(3.17) \quad |I(v, w)|^{\frac{4}{3}} \leq C_I \left(1 + |v|^4 + |w|^2 \right),$$

and therefore $\|I(v^\varepsilon, w^\varepsilon)\|_{L^{4/3}((0, T) \times \Gamma^\varepsilon)}^{4/3} < \infty$. Consequently, $J < \infty$.

We actually have a more precise bound. As $H^{1/2}(\Gamma^\varepsilon) \subset L^4(\Gamma^\varepsilon)$, we have $H^{-1/2}(\Gamma^\varepsilon) \subset L^{4/3}(\Gamma^\varepsilon)$ and

$$\|I(v^\varepsilon, w^\varepsilon)\|_{H^{-1/2}(\Gamma^\varepsilon)}^{4/3} \leq \tilde{C} \|I(v^\varepsilon, w^\varepsilon)\|_{L^{4/3}(\Gamma^\varepsilon)}^{4/3} \leq C \left(1 + \|v^\varepsilon\|_{L^4(\Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right).$$

Integrating this over $t \in (0, T)$ yields $I(v^\varepsilon, w^\varepsilon) \in L^{4/3}(0, T; H^{-1/2}(\Gamma^\varepsilon))$.

The integral on the right-hand side of (3.15) can be treated similarly, since $|H(v, w)|^2 \leq C(|v|^4 + |w|^2)$. The remaining integrals are trivially well defined.

3.3. Existence of solution and a priori estimates. Existence and uniqueness results for certain classes of membrane models have been established in [14, 41]. These works employ the variable $v = u_i - u_e$ to convert (1.1) into a non-degenerate ‘‘abstract’’ parabolic equation. The authors in [14] then appeal to the theory of variational inequalities, whereas in [41] the Schauder fixed point theorem is applied to conclude the existence of a solution.

The following theorem can be proved by adapting arguments found in [14, 41] (see also [21]), or those utilized in [8, 6, 5] for the (macroscopic) bidomain model.

Theorem 3.5 (existence of weak solution for microscopic system). *Fix any $\varepsilon > 0$, and suppose (**GFHN**), (1.2), (3.4), (3.5), (3.6), (3.7), (3.9), and (3.10) hold. Then the microscopic bidomain model (1.1), (1.3) possesses a weak solution $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$ (in the sense of Definition 3.2). This weak solution satisfies the a priori estimates collected in Lemma 3.8 below.*

Remark 3.6. *The unknowns $u_i^\varepsilon, u_e^\varepsilon$ are determined uniquely up to a constant. We can fix this constant by imposing the normalization condition $\int u_e^\varepsilon dx = 0$, as is done in Definition 3.2. Weak solutions are unique (and L^2 stable) provided we impose the structural condition (3.3) on the membrane functions I, H .*

The next lemma, which is utilized below to derive some a priori estimates, is a consequence of the uniform Poincaré inequality (2.9) and the trace inequality for ε -periodic surfaces (2.8). A similar result is used in [34].

Lemma 3.7. *Let $u_j \in H^1(\Omega_j^\varepsilon)$, $j = i, e$, $\int_{\Omega_j^\varepsilon} u_e dx = 0$, and set $v := u_i|_{\Gamma^\varepsilon} - u_e|_{\Gamma^\varepsilon}$. There is a positive constants C , independent of ε , such that*

$$\|u_i\|_{L^2(\Omega_i^\varepsilon)}^2 \leq C \left(\varepsilon \|v\|_{L^2(\Gamma^\varepsilon)}^2 + \sum_{j=i,e} \int_{\Omega_j^\varepsilon} |\nabla u_j|^2 dx \right).$$

Proof. First, since $\int u_e = 0$,

$$(3.18) \quad \|u_e\|_{L^2(\Omega_\varepsilon^\varepsilon)}^2 \stackrel{(2.9)}{\leq} C_1 \int_{\Omega_\varepsilon^\varepsilon} |\nabla u_e|^2 dx.$$

To estimate the L^2 norm of u_i , write $u_i = \bar{u}_i + \tilde{u}_i$, where $\bar{u}_i := \frac{1}{|\Omega_i^\varepsilon|} \int_{\Omega_i^\varepsilon} u_i dx$ is constant in Ω_i^ε and $\tilde{u}_i := u_i - \bar{u}_i$ has zero mean in Ω_i^ε . Clearly,

$$\|u_i\|_{L^2(\Omega_i^\varepsilon)}^2 = \|\bar{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 + \|\tilde{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2.$$

In view of the Poincaré inequality (2.9),

$$(3.19) \quad \|\tilde{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 \leq \tilde{C}_1 \int_{\Omega_i^\varepsilon} |\nabla \tilde{u}_i|^2 dx = \tilde{C}_1 \int_{\Omega_i^\varepsilon} |\nabla u_i|^2 dx.$$

Let us bound $\|\bar{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 = \frac{|\Omega_i^\varepsilon|}{|\Gamma^\varepsilon|} \|\bar{u}_i\|_{L^2(\Gamma^\varepsilon)}^2$. Since $|\Gamma^\varepsilon| = |K^\varepsilon| \int_{\varepsilon\Gamma} dS = \varepsilon^{-3} \varepsilon^2 \int_\Gamma dS$ (recall (2.7)), $\varepsilon |\Gamma^\varepsilon| \geq c > 0$. Because of this and $|\Omega_i^\varepsilon| \leq |\Omega|$,

$$\|\bar{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 \leq \bar{C}_1 \varepsilon \|\bar{u}_i\|_{L^2(\Gamma^\varepsilon)}^2.$$

Noting that

$$|\bar{u}_i|^2 \leq \bar{C}_2 \left(|u_i - u_e|^2 + |\tilde{u}_i|^2 + |u_e|^2 \right),$$

we obtain

$$\begin{aligned} \|\bar{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 &\leq \bar{C}_3 \left(\varepsilon \|v\|_{L^2(\Gamma^\varepsilon)}^2 + \varepsilon \|\tilde{u}_i\|_{L^2(\Gamma^\varepsilon)}^2 + \varepsilon \|u_e\|_{L^2(\Gamma^\varepsilon)}^2 \right) \\ &\stackrel{(2.8)}{\leq} \bar{C}_3 \varepsilon \|v\|_{L^2(\Gamma^\varepsilon)}^2 \\ &\quad + \bar{C}_4 \left(\|\tilde{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 + \varepsilon^2 \|\nabla \tilde{u}_i\|_{L^2(\Omega_i^\varepsilon)}^2 \right) \\ &\quad + \bar{C}_4 \left(\|u_e\|_{L^2(\Omega_\varepsilon^\varepsilon)}^2 + \varepsilon^2 \|\nabla u_e\|_{L^2(\Omega_\varepsilon^\varepsilon)}^2 \right) \\ &\leq \bar{C}_3 \varepsilon \|v\|_{L^2(\Gamma^\varepsilon)}^2 + \bar{C}_5 \|\nabla u_i\|_{L^2(\Omega_i^\varepsilon)}^2 + \bar{C}_5 \|\nabla u_e\|_{L^2(\Omega_\varepsilon^\varepsilon)}^2, \end{aligned}$$

where the final inequality is a result of (3.18) and (3.19). \square

For the sake of the upcoming homogenization result, we now list some precise (ε -independent) a priori estimates.

Lemma 3.8 (basic estimates for microscopic system). *Referring Theorem 3.5, the weak solution $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$ of (1.1), (1.3) satisfies the a priori estimates*

$$(3.20) \quad \begin{aligned} (a) \quad &\|\nabla u_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \leq C, \quad j = i, e, \\ (b) \quad &\|u_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \leq C, \quad j = i, e, \\ (c) \quad &\varepsilon^{1/2} \|v^\varepsilon\|_{L^\infty(0,T; L^2(\Gamma^\varepsilon))} \leq C, \\ (d) \quad &\varepsilon^{1/4} \|v^\varepsilon\|_{L^4((0,T) \times \Gamma^\varepsilon)} \leq C, \\ (e) \quad &\varepsilon^{1/2} \|w^\varepsilon\|_{L^\infty(0,T; L^2(\Gamma^\varepsilon))} \leq C, \\ (f) \quad &\varepsilon^{1/2} \partial_t w^\varepsilon \in L^2(0, T; L^2(\Gamma^\varepsilon)), \end{aligned}$$

where C is a positive constant independent of ε .

Proof. We only outline a proof of these (mostly standard) estimates.

Specifying $(\varphi_i, \varphi_e, \varphi_w) = (u_i^\varepsilon, -u_e^\varepsilon, w^\varepsilon)$ as test functions in (3.13), (3.14), and (3.15), adding the resulting equations, applying the chain rule (2.5), and using

(3.6), we obtain

$$(3.21) \quad \begin{aligned} & \frac{\varepsilon}{2} \frac{d}{dt} \left(\|v^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 + \|w^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right) + \alpha \sum_{j=i,e} \int_{\Omega_j^\varepsilon} |\nabla u_j^\varepsilon|^2 dx, \\ & + \varepsilon \int_{\Gamma^\varepsilon} (I(v^\varepsilon, w^\varepsilon)v^\varepsilon - H(v^\varepsilon, w^\varepsilon)w^\varepsilon) dS \leq \sum_{j=i,e} \int_{\Omega_j^\varepsilon} s_j^\varepsilon u_j^\varepsilon dx. \end{aligned}$$

Thanks to (3.2),

$$(3.22) \quad \begin{aligned} & \int_{\Gamma^\varepsilon} (I(v^\varepsilon, w^\varepsilon)v^\varepsilon - H(v^\varepsilon, w^\varepsilon)w^\varepsilon) dS \\ & \geq \gamma \|v^\varepsilon\|_{L^4(\Gamma^\varepsilon)}^4 - \beta \left(\|v^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 + \|w^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right), \end{aligned}$$

for some ε -independent constants $\gamma > 0$ and $\beta \geq 0$.

Using Cauchy's inequality ("with δ "), the source term can be bounded as

$$(3.23) \quad \begin{aligned} \left| \sum_{j=i,e} \int_{\Omega_j^\varepsilon} s_j^\varepsilon u_j^\varepsilon dx \right| & \leq \sum_{j=i,e} \left(\|u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)} \|s_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)} \right) \\ & \leq \delta \sum_{j=i,e} \|u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2 + C_\delta \sum_{j=i,e} \|s_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2, \end{aligned}$$

with $\delta > 0$ small and C_δ independent of ε . Lemma 3.7 and (3.18) ensure that

$$(3.24) \quad \sum_{j=i,e} \|u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2 \leq C_1 \left(\varepsilon \|v^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 + \sum_{j=i,e} \int_{\Omega_j^\varepsilon} |\nabla u_j^\varepsilon|^2 dx \right).$$

Insert (3.22), (3.23), (3.24) into (3.21), use (3.6), and choose δ (appropriately) small. Integrating the result over the time interval $(0, \tau)$, for $\tau \in (0, T)$, yields

$$\begin{aligned} & \varepsilon \left(\|v_\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 + \|w_\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 \right) + \frac{\alpha}{2} \sum_{j=i,e} \int_0^\tau \|\nabla u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2 dt \\ & + \gamma \varepsilon \int_0^\tau \|v^\varepsilon(t)\|_{L^4(\Omega)}^4 dt \leq \varepsilon \left(\|v_\varepsilon(0)\|_{L^2(\Gamma^\varepsilon)}^2 + \|w_\varepsilon(0)\|_{L^2(\Gamma^\varepsilon)}^2 \right) \\ & + C_2 \varepsilon \int_0^\tau \left(\|v_\varepsilon(t)\|_{L^2(\Gamma^\varepsilon)}^2 + \|w_\varepsilon(t)\|_{L^2(\Gamma^\varepsilon)}^2 \right) dt + C_\delta \sum_{j=i,e} \|s_j^\varepsilon\|_{L^2((0,\tau) \times \Omega_j^\varepsilon)}^2, \end{aligned}$$

for some positive constant C_2 independent of ε . Applying Gröwall's inequality, recalling (3.10) and (3.8), we obtain estimates (a), (c), (d), (e) in (3.20). Estimate (b) follows from (3.24) and (a), (c). Finally, note that **(GFHN)** implies the bound

$$|\partial_t w^\varepsilon|^2 \leq C \left(1 + |v^\varepsilon|^4 + |v^\varepsilon|^2 \right).$$

Estimate (f) follows by integrating this over $(0, T) \times \Gamma^\varepsilon$ and using (d), (e). \square

Remark 3.9. *Let us motivate the regularity requirements in Definition 3.2 that are not covered by Lemma 3.8. First, due to (3.11), (2.1) and (3.20),*

$$\|v^\varepsilon\|_{L^2(0,T;H^{1/2}(\Gamma^\varepsilon))} \leq \tilde{C}_\varepsilon \sum_{j=i,e} \|u_j^\varepsilon\|_{L^2(0,T;H^1(\Omega_j^\varepsilon))} \leq C_\varepsilon,$$

where the constants $\tilde{C}_\varepsilon, C_\varepsilon$ may depend on ε (via (2.1) with Γ replaced by Γ^ε).

Next, we claim that

$$\varepsilon \|\partial_t v^\varepsilon\|_{L^2(0,T;H^{-1/2}(\Gamma^\varepsilon)) + L^4((0,T) \times \Gamma^\varepsilon)} \leq C_\varepsilon,$$

for some constant that may depend on ε . To see this use a version of (3.13) or (3.14) (with time-independent test functions) to write

$$\varepsilon \langle \partial_t v^\varepsilon, \psi \rangle_{H^{-1/2}(\Gamma^\varepsilon), H^{1/2}(\Gamma^\varepsilon)} = J_1(\psi) + J_2(\psi) + J_3(\psi),$$

for a.e. $t \in (0, T)$ and for any $\psi \in H^{1/2}(\Gamma^\varepsilon)$ with $\|\psi\|_{H^{1/2}(\Gamma^\varepsilon)} = 1$, where

$$\begin{aligned} |J_1(\psi)| &= \left| \int_{\Omega_j^\varepsilon} \sigma_j^\varepsilon \nabla u_j^\varepsilon \cdot \nabla \mathcal{I}_j(\psi) dx \right|, & |J_2(\psi)| &= \left| \int_{\Omega_j^\varepsilon} s_j^\varepsilon \mathcal{I}_j(\psi) dx \right|, \\ |J_3(\psi)| &= \left| \varepsilon \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \psi dS \right|, \end{aligned}$$

and $\mathcal{I}_j(\cdot)$ is the right inverse of the trace operator relative to Ω_j^ε , for $j = i$ or $j = e$. Clearly, using the Cauchy-Schwarz inequality and (2.2),

$$\begin{aligned} |J_1(\psi)| &\leq \tilde{C}_1 \|\nabla u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)} \|\psi\|_{H^{1/2}(\Gamma^\varepsilon)}, & |J_2(\psi)| &\leq \tilde{C}_2 \|s_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)} \|\psi\|_{H^{1/2}(\Gamma^\varepsilon)}, \\ \text{and } |J_3(\psi)| &\leq \varepsilon \|I(v^\varepsilon, w^\varepsilon)\|_{H^{-1/2}(\Gamma^\varepsilon)} \|\psi\|_{H^{1/2}(\Gamma^\varepsilon)}, \end{aligned}$$

where the constants \tilde{C}_1, \tilde{C}_2 may depend on ε (via the inverse trace inequality (2.2) with Γ replaced by Γ^ε). As a result, for a.e. $t \in (0, T)$,

$$\|J_1\|_{H^{-1/2}(\Gamma^\varepsilon)}^2 \leq \tilde{C}_1^2 \|\nabla u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2, \quad \|J_2\|_{H^{-1/2}(\Gamma^\varepsilon)}^2 \leq \tilde{C}_2^2 \|s_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2.$$

Integrating this over $t \in (0, T)$ and using (3.20)-(a), (3.8) it follows that

$$\|J_1\|_{L^2(0, T; H^{-1/2}(\Gamma^\varepsilon))} \leq C_1, \quad \|J_2\|_{L^2(0, T; H^{-1/2}(\Gamma^\varepsilon))} \leq C_2,$$

for some constants C_1, C_2 (that may depend on ε).

As in Remark 3.4,

$$\|I(v^\varepsilon, w^\varepsilon)\|_{H^{-1/2}(\Gamma^\varepsilon)} \leq C_I \left(1 + \|v^\varepsilon\|_{L^4(\Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right)^{3/4},$$

and hence, for a.e. $t \in (0, T)$,

$$\|J_3\|_{H^{-1/2}(\Gamma^\varepsilon)}^{4/3} \leq \tilde{C}_3 \varepsilon^{4/3} \left(1 + \|v^\varepsilon\|_{L^4(\Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right).$$

Integrating over $t \in (0, T)$ and using estimates (3.20)-(d, e), we conclude

$$\|J_3\|_{L^{4/3}(0, T; H^{-1/2}(\Gamma^\varepsilon))} \leq C_3,$$

for some constant C_3 (independent of ε).

For the upcoming convergence analysis, we need a temporal translation estimate for the membrane potential v^ε .

Lemma 3.10 (temporal translation estimate in L^2). *Referring back to Theorem 3.5, let $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)$ be the weak solution of (1.1), (1.3). Then v^ε satisfies the following L^2 temporal translation estimate for some ε -independent constant $C > 0$, for any sufficiently small $\Delta_t > 0$:*

$$(3.25) \quad \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} |v^\varepsilon(t + \Delta_t, x) - v^\varepsilon(t, x)|^2 dx dt \leq C \Delta_t.$$

Proof. The translated functions $(u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon)(t + \Delta_t)$ constitute a weak solution of (1.1) on $(0, T - \Delta_t)$ with initial data $(v^\varepsilon, w^\varepsilon)(\Delta_t)$ and stimulation currents $(s_i^\varepsilon, s_e^\varepsilon)(t + \Delta_t)$. We subtract the equations (linked to Ω_i^ε and Ω_e^ε) for the original weak solution from the equations satisfied by the translated one, and add the resulting equations. The result is

$$\begin{aligned}
& - \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} (v^\varepsilon(t+\Delta_t) - v^\varepsilon(t)) \partial_t(\varphi_i - \varphi_e) dS dt \\
& + \varepsilon \int_{\Gamma^\varepsilon} (v^\varepsilon(T) - v^\varepsilon(T-\Delta_t)) (\varphi_i(T-\Delta_t) - \varphi_e(T-\Delta_t)) dS \\
& - \varepsilon \int_{\Gamma^\varepsilon} (v^\varepsilon(\Delta_t) - v_0^\varepsilon) (\varphi_i(0) - \varphi_e(0)) dS \\
& + \sum_{j=i,e} \int_0^{T-\Delta_t} \int_{\Omega_j^\varepsilon} \sigma_j^\varepsilon \nabla (u_j^\varepsilon(t+\Delta_t) - u_j^\varepsilon(t)) \cdot \nabla \varphi_j dx dt \\
& + \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} (I(v^\varepsilon(t+\Delta_t), w^\varepsilon(t+\Delta_t)) - I(v^\varepsilon(t), w^\varepsilon(t))) (\varphi_i - \varphi_e) dS dt \\
& - \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} (H(v^\varepsilon(t+\Delta_t), w^\varepsilon(t+\Delta_t)) - H(v^\varepsilon(t), w^\varepsilon(t))) \varphi_w dS dt \\
& = \sum_{j=i,e} \int_0^{T-\Delta_t} \int_{\Omega_j^\varepsilon} (s_j^\varepsilon(t+\Delta_t) - s_j^\varepsilon(t)) \varphi_j dx dt.
\end{aligned}$$

Specifying the test functions $\varphi_i, \varphi_e, \varphi_w$ as

$$\varphi_j(t) = - \int_t^{t+\Delta_t} u_j^\varepsilon(s) ds, \quad j = i, e, \quad \varphi_w(t) = - \int_t^{t+\Delta_t} w_j^\varepsilon(s) ds,$$

we obtain

$$\begin{aligned}
& \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} |v^\varepsilon(t+\Delta_t) - v^\varepsilon(t)|^2 dS dt \\
& + \varepsilon \int_{\Gamma^\varepsilon} (v^\varepsilon(T) - v^\varepsilon(T-\Delta_t)) \left(- \int_{T-\Delta_t}^T v(s) ds \right) dS \\
& - \varepsilon \int_{\Gamma^\varepsilon} (v^\varepsilon(\Delta_t) - v_0^\varepsilon) \left(- \int_0^{\Delta_t} v(s) ds \right) dS \\
& + \sum_{j=i,e} \int_0^{T-\Delta_t} \int_{\Omega_j^\varepsilon} \sigma_j^\varepsilon \nabla (u_j^\varepsilon(t+\Delta_t) - u_j^\varepsilon(t)) \\
& \quad \cdot \nabla \left(- \int_t^{t+\Delta_t} u_j^\varepsilon(s) ds \right) dx dt \\
& + \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} (I(v^\varepsilon(t+\Delta_t), w^\varepsilon(t+\Delta_t)) - I(v^\varepsilon(t), w^\varepsilon(t))) \\
& \quad \times \left(- \int_t^{t+\Delta_t} v^\varepsilon(s) ds \right) dS dt \\
& + \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} (H(v^\varepsilon(t+\Delta_t), w^\varepsilon(t+\Delta_t)) - H(v^\varepsilon(t), w^\varepsilon(t))) \\
& \quad \times \left(\int_t^{t+\Delta_t} w^\varepsilon(s) ds \right) dS dt \\
& = \sum_{j=i,e} \int_0^{T-\Delta_t} \int_{\Omega_j^\varepsilon} s_j^\varepsilon \left(- \int_t^{t+\Delta_t} u_j^\varepsilon(s) ds \right) dx dt.
\end{aligned}$$

Let us write this equation as

$$\varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} |v^\varepsilon(t+\Delta_t) - v^\varepsilon(t)|^2 dS dt + J_1 + J_2 + J_3 + J_4 + J_5 = J_6.$$

By the Cauchy-Schwarz and Minkowski integral inequalities,

$$\begin{aligned}
|J_1| &\leq \varepsilon \|v^\varepsilon(T) - v^\varepsilon(T - \Delta_t)\|_{L^2(\Gamma^\varepsilon)} \left\| \int_{T-\Delta_t}^T v^\varepsilon(s) ds \right\|_{L^2(\Gamma^\varepsilon)} \\
&\leq 2\varepsilon \|v^\varepsilon\|_{L^\infty(0,T;L^2(\Gamma^\varepsilon))} \int_{T-\Delta_t}^T \|v^\varepsilon\|_{L^2(\Gamma^\varepsilon)} ds \\
&\leq 2\varepsilon \|v^\varepsilon\|_{L^\infty(0,T;L^2(\Gamma^\varepsilon))}^2 \Delta_t \stackrel{(3.20)-(c)}{\leq} C_1 \Delta_t.
\end{aligned}$$

Similarly, $|J_2| \leq C_2 \Delta_t$.

We need the following facts involving a real-valued function $f \in L^p$ ($p > 1$):

$$\begin{aligned}
(3.26) \quad &\int_t^{t+\Delta_t} f(s) ds = (\mathbb{1}_{[0,\Delta_t]} \star f)(t), \\
&\|\mathbb{1}_{[0,\Delta_t]} \star f\|_{L^p} \leq \|\mathbb{1}_{[0,\Delta_t]}\|_{L^1} \|f\|_{L^p} = \Delta_t \|f\|_{L^p},
\end{aligned}$$

where the second line follows from Young's convolution inequality.

By (3.5) and the Cauchy-Schwarz inequality,

$$|J_3| \leq \sum_{j=i,e} 2 \|\sigma_j^\varepsilon\|_{L^\infty} \|\nabla u_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \left\| \int_t^{t+\Delta_t} \nabla u_j^\varepsilon(s) ds \right\|_{L^2((0,T-\Delta_t) \times \Omega_j^\varepsilon)}.$$

As a result of Minkowski integral inequality,

$$\begin{aligned}
&\left\| \int_t^{t+\Delta_t} \nabla u_j^\varepsilon(s) ds \right\|_{L^2((0,T-\Delta_t) \times \Omega_j^\varepsilon)}^2 \\
&= \int_0^{T-\Delta_t} \int_{\Omega_j^\varepsilon} \left| \int_t^{t+\Delta_t} \nabla u_j^\varepsilon(s) ds \right|^2 dx dt \\
&\leq \int_0^{T-\Delta_t} \left(\int_t^{t+\Delta_t} \left(\int_{\Omega_j^\varepsilon} |\nabla u_j^\varepsilon(s)|^2 dx \right)^{1/2} ds \right)^2 dt \\
&= \int_0^{T-\Delta_t} \left(\int_t^{t+\Delta_t} \|\nabla u_j^\varepsilon(s)\|_{L^2(\Omega_j^\varepsilon)} ds \right)^2 dt \\
&\stackrel{(3.26)}{\leq} \Delta_t^2 \int_0^{T-\Delta_t} \|\nabla u_j^\varepsilon\|_{L^2(\Omega_j^\varepsilon)}^2 dt \leq \Delta_t^2 \|\nabla u_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)}^2.
\end{aligned}$$

Therefore,

$$|J_3| \leq \sum_{j=i,e} 2 \|\sigma_j^\varepsilon\|_{L^\infty} \|\nabla u_j^\varepsilon(t)\|_{L^2((0,T) \times \Omega_j^\varepsilon)}^2 \Delta_t \stackrel{(3.20)-(a)}{\leq} C_3 \Delta_t.$$

An application of Hölder's inequality yields

$$\begin{aligned}
|J_4| &\leq 2\varepsilon \|I^\varepsilon\|_{L^{4/3}((0,T) \times \Gamma^\varepsilon)} \left\| \int_t^{t+\Delta_t} v^\varepsilon(s) ds \right\|_{L^4((0,T-\Delta_t) \times \Gamma^\varepsilon)} \\
&\stackrel{(3.17)}{\leq} C_4 \varepsilon \left(\|v^\varepsilon\|_{L^4((0,T) \times \Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2((0,T) \times \Gamma^\varepsilon)}^2 \right)^{3/4} \Delta_t \|v^\varepsilon\|_{L^4((0,T) \times \Gamma^\varepsilon)} \\
&\leq C_5 \Delta_t \varepsilon \left(\|v^\varepsilon\|_{L^4((0,T) \times \Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2((0,T) \times \Gamma^\varepsilon)}^2 \right) \stackrel{(3.20)-(d,e)}{\leq} C_6 \Delta_t,
\end{aligned}$$

where we have repeated the argument for J_1 involving (3.26), with the implication that $\left\| \int_t^{t+\Delta_t} v^\varepsilon(s) ds \right\|_{L^4_{t,x}}$ is bounded in terms of $\Delta_t \|v^\varepsilon\|_{L^4_{t,x}}$. Moreover, we have used the basic inequality $ab \leq \frac{3}{4}a^{4/3} + \frac{1}{4}b^4$ for positive numbers a, b .

Similarly,

$$\begin{aligned} |J_5| &\stackrel{(\mathbf{GFHN})}{\leq} C_7 \Delta_t \varepsilon \left(\|v^\varepsilon\|_{L^4((0,T) \times \Gamma^\varepsilon)}^4 + \|w^\varepsilon\|_{L^2((0,T) \times \Gamma^\varepsilon)}^2 \right)^{1/2} \|v^\varepsilon\|_{L^2((0,T) \times \Gamma^\varepsilon)} \\ &\stackrel{(3.20)-(d,e)}{\leq} C_8 \Delta_t, \end{aligned}$$

and

$$|J_6| \leq 2\Delta_t \sum_{j=i,e} \|s_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \|u_j^\varepsilon\|_{L^2((0,T) \times \Omega_j^\varepsilon)} \stackrel{(3.8), (3.20)-(b)}{\leq} C_9 \Delta_t.$$

Summarizing our findings, we conclude that (3.25) holds. \square

Remark 3.11. *Due to the degenerate structure of the microscopic system (1.1), temporal estimates are not available for the intra- and extracellular potentials $u_i^\varepsilon, u_e^\varepsilon$.*

4. THE HOMOGENIZATION RESULT

This section contains the main result of the paper. We start by recalling the weak formulation of the macroscopic bidomain system (1.4), which is augmented with the following initial and boundary conditions:

$$(4.1) \quad \begin{aligned} v|_{t=0} &= v_0 \quad \text{in } \Omega, & w|_{t=0} &= w_0, \quad \text{in } \Omega. \\ n \cdot \sigma \nabla u_j &= 0 \quad \text{on } (0, T) \times \partial\Omega, \quad j = i, e. \end{aligned}$$

Definition 4.1 (weak formulation of bidomain system). *A weak solution to (1.4), (4.1) is a collection (u_i, u_e, v, w) of functions satisfying the following conditions:*

i (algebraic relation).

$$v = u_i - u_e \quad \text{a.e. in } (0, T) \times \Omega.$$

ii (regularity).

$$\begin{aligned} u_j &\in L^2(0, T; H^1(\Omega)), \quad j = i, e, \\ \int_{\Omega} u_e^\varepsilon(t, x) dx &= 0, \quad t \in (0, T), \\ v &\in L^2(0, T; H^1(\Omega)) \cap L^4((0, T) \times \Omega), \\ \partial_t v &\in L^2(0, T; H^{-1}(\Omega)) + L^{4/3}((0, T) \times \Omega), \\ w &\in H^1(0, T; L^2(\Omega)). \end{aligned}$$

iii (initial conditions).

$$v(0) = v_0, \quad w(0) = w_0.$$

iv (differential equations).

$$\begin{aligned} |\Gamma| \int_0^T \langle \partial_t v, \varphi_i \rangle dt + \int_0^T \int_{\Omega} M_i \nabla u_i \cdot \nabla \varphi_i dx dt \\ + |\Gamma| \int_0^T \int_{\Omega} I(v, w) \varphi_i dx dt = |Y_i| \int_0^T \int_{\Omega} s_i \varphi_i dx dt, \end{aligned}$$

$$\begin{aligned}
& |\Gamma| \int_0^T \langle \partial_t v, \varphi_e \rangle dt - \int_0^T \int_{\Omega} M_e \nabla u_e \cdot \nabla \varphi_e dx dt \\
& + |\Gamma| \int_0^T \int_{\Omega} I(v, w) \varphi_e dx dt = -|Y_e| \int_0^T \int_{\Omega} s_e \varphi_e dx dt, \\
(4.2) \quad & \int_0^T \int_{\Omega} \partial_t w \varphi_w dx dt = \int_0^T \int_{\Omega} H(v, w) \varphi_w dx dt,
\end{aligned}$$

for all test functions $\varphi_i, \varphi_e \in L^2(0, T; H^1(\Omega)) \cap L^4((0, T) \times \Omega)$, $\varphi_w \in L^2(0, T; L^2(\Omega))$. We denote by $\langle \cdot, \cdot \rangle$ the duality pairing between $L^2(0, T; H^{-1}(\Omega)) + L^{4/3}((0, T) \times \Omega)$ and $L^2(0, T; H^1(\Omega)) \cap L^4((0, T) \times \Omega)$.

The macroscopic bidomain system is well studied for a variety of cellular models [5, 6, 7, 8, 14, 42]. For the following result, see [5, 6, 7].

Theorem 4.2 (well-posedness of bidomain system). *Suppose (1.2) holds, I and H satisfy the conditions in (GFHN) and (3.3) (for uniqueness), $M_i(x)$ and $M_e(x)$ are bounded, positive definite matrices, $s_i, s_e \in L^2((0, T) \times \Omega)$, and $v_0, w_0 \in L^2(\Omega)$. Then there exists a unique weak solution to the bidomain system (1.4), (4.1).*

We are now in a position to state the main result, which should be compared to Theorem 1.3 in [34].

Theorem 4.3 (convergence to the bidomain system). *Suppose conditions (GFHN), (1.2), (3.3), (3.4), (3.5), (3.6), (3.7), (3.9), and (3.10) hold. Let ε take values in a sequence tending to zero (e.g. $\varepsilon^{-1} \in \mathbb{N}$). Then the sequence $\{u_i^\varepsilon, u_e^\varepsilon, v^\varepsilon, w^\varepsilon\}_{\varepsilon > 0}$ of weak solutions to the microscopic system (1.1), (1.3) two-scale converges (in the sense of (4.4) below) to the weak solution (u_i, u_e, v, w) of the macroscopic bidomain system (1.4), (4.1). Moreover, $\{v^\varepsilon\}_{\varepsilon > 0}$ converges strongly in the sense that*

$$(4.3) \quad \varepsilon^{1/2} \|v^\varepsilon - v\|_{L^2(\Gamma^\varepsilon)} \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Remark 4.4. *The strong convergence (4.3) is a corrector-type result. In the current setting it allows us to pass to the limit in the nonlinear ionic terms. By employing standard techniques [12] one can also show that the energies*

$$\left\{ \int_0^T \int_{\Omega_j^\varepsilon} \sigma_j^\varepsilon \nabla u_i^\varepsilon \cdot \nabla u_j^\varepsilon dx dt \right\}_{\varepsilon > 0}$$

converge to the homogenized energy

$$\int_0^T \int_{\Omega} M_j \nabla u_i \cdot \nabla u_j dx dt,$$

where M_j is defined in (1.5), $j = i, e$.

The rest of this section is devoted to the proof of Theorem 4.3. Homogenization of the linear terms in (1.1) is handled with standard techniques, cf. Subsection 4.1.

Passing to the limit in the nonlinear terms I, H is more challenging. Although the intracellular/extracellular functions u_i^ε (defined on Ω_i^ε) and u_e^ε (defined on Ω_e^ε) do not converge strongly, some kind of strong compactness is expected for the ε -scaled version of the transmembrane potential $v^\varepsilon = u_i^\varepsilon|_{\Gamma^\varepsilon} - u_e^\varepsilon|_{\Gamma^\varepsilon}$ (defined on Γ^ε), since we control both the temporal (3.25) and spatial (fractional) derivatives (3.20).

Wild oscillations of the underlying domain do however pose difficulties. For this reason, we use the boundary unfolding operator $\mathcal{T}_\varepsilon^b$, cf. (2.12), to transform the problem of convergence on the oscillating set Γ^ε to the fixed set $\Omega \times \Gamma$. Roughly speaking, (3.20) is used to conclude that $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ is uniformly bounded in $L_t^2 L_x^2 H_y^{1/2}$.

In addition, in view of (3.25), $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ possesses an ε -uniform temporal translation estimate with respect to the $L^2_{t,x,y}$ norm. As a result, the Simon compactness result (cf. Subsection 2.1) implies that $\{(t, y) \mapsto \mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, y)\}_{\varepsilon>0}$ is precompact in $L^2((0, T) \times \Gamma)$, for fixed x . Next we demonstrate that $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ is equicontinuous in x (with values in $L^2((0, T) \times \Gamma)$). Applying the Simon-type compactness criterion found in [19] (cf. Theorem 4.6 below), it follows that $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ converges along a subsequence. Owing to the uniqueness of solutions to the bidomain system (4.1), the entire sequence $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ converges (not just a subsequence). We refer to Subsection 4.2 for details. For inspirational works deriving macroscopic models by combining two-scale and unfolding techniques, we refer to [19, 18, 20, 29, 32].

4.1. Extracting two-scale limits. Recall that $\tilde{\cdot}$ denotes the extension to Ω by zero, and that $\mathbb{1}_{Y_j}$ is the indicator function of Y_j ($j = i, e$). Using the a priori estimates provided by Lemma 3.8, we can apply Lemma 2.3, Theorem 2.4, and Lemma 2.5 to extract subsequences (not relabelled) such that

$$(4.4) \quad \begin{aligned} \widetilde{u_j^\varepsilon} &\xrightarrow{2} \mathbb{1}_{Y_j}(y) u_j(t, x), \quad j = i, e \\ \widetilde{\nabla u_j^\varepsilon} &\xrightarrow{2} \mathbb{1}_{Y_j}(y) (\nabla_x u_j(t, x) + \nabla_y u_j^1(t, x, y)), \quad j = i, e, \\ v^\varepsilon &\xrightarrow{2-S} v = u_i - u_e, \\ w^\varepsilon &\xrightarrow{2-S} w, \quad \partial_t w^\varepsilon \xrightarrow{2-S} \partial_t w, \\ I(v^\varepsilon, w^\varepsilon) &\xrightarrow{2-S} \bar{I}, \quad H(v^\varepsilon, w^\varepsilon) \xrightarrow{2-S} \bar{H}, \end{aligned}$$

for some limits $u_i, u_e \in L^2(0, T, H^1(\Omega))$, $u_i^1, u_e^1 \in L^2((0, T) \times \Omega; H^1_{\text{per}}(Y))$, and $w \in L^2((0, T) \times \Omega \times \Gamma)$. Here we identify $v = u_i - u_e$ as an element in $L^2(0, T, H^1(\Omega))$. It is easily verified that the two-scale limit u_e satisfies $\int_\Omega u_e dx = 0$.

Nonlinear functions are not continuous with respect to weak convergence, which prevents us from immediately making the identifications

$$\bar{I} = I(v, w), \quad \bar{H} = H(v, w).$$

Using the two-scale convergences in (4.4), Remark 3.3, the choice of test function

$$\begin{aligned} \varphi(t, x) + \varepsilon \varphi_1\left(t, x, \frac{x}{\varepsilon}\right), \quad \text{with} \\ \varphi \in C_0^\infty((0, T) \times \Omega), \quad \varphi_1 \in C_0^\infty((0, T) \times \Omega; C^\infty_{\text{per}}(Y)), \end{aligned}$$

in (3.13), (3.14), and (3.15), standard manipulations [12, 2] will reveal that the two-scale limit (u_i, u_e, v, w) satisfies the following equations:

$$(4.5) \quad \begin{aligned} |\Gamma| \int_0^T \int_\Omega \langle \partial_t v, \varphi \rangle dx dt + \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi dS dt \\ + \int_0^T \int_\Omega M_i \nabla u_i \cdot \nabla \varphi dx dt = |Y_i| \int_0^T \int_\Omega s_i \varphi dx dt, \end{aligned}$$

$$(4.6) \quad \begin{aligned} |\Gamma| \int_0^T \int_\Omega \langle \partial_t v, \varphi \rangle dx dt + \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi dS dt \\ + \int_0^T \int_\Omega M_e \nabla u_e \cdot \nabla \varphi dx dt = |Y_e| \int_0^T \int_\Omega s_e \varphi dx dt, \end{aligned}$$

and

$$(4.7) \quad |\Gamma| \int_0^T \int_\Omega \partial_t w \varphi dx dt = \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} H(v^\varepsilon, w^\varepsilon) \varphi dS dt,$$

for all $\varphi \in C_0^\infty((0, T) \times \Omega)$.

In (4.5) and (4.6), M_i and M_e are the homogenized conductivity tensors (1.5). Let us briefly recall how one arrives at the homogenized conductivities. Setting $\varphi \equiv 0$ and considering

$$\Phi(t, x, y) := \sigma_j(x, y) \nabla_y \varphi_1(t, x, y)$$

as a test function for two-scale convergence, we have by (4.4) that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega_\varepsilon} \Phi \left(t, x, \frac{x}{\varepsilon} \right) \cdot \nabla u_j^\varepsilon dx dt \\ = \int_0^T \int_{\Omega} \int_{Y_j} \sigma_j [\nabla_x u_j + \nabla_y u_j^1] \cdot \nabla_y \varphi_1 dx dy dt. \end{aligned}$$

Note that the oscillating φ_1 term is suppressed in the limit of the weak formulation (3.2) by the ε -factor, except in the term where a gradient hits the test function. Thus, the two-scale limit (u_j, u_j^1) satisfies the equation

$$\int_0^T \int_{\Omega} \int_{Y_j} \sigma_j [\nabla_x u_j + \nabla_y u_j^1] \cdot \nabla_y \varphi_1 dx dy dt = 0,$$

for all $\varphi_1 \in C_0^\infty((0, T) \times \Omega; C_{\text{per}}^\infty(Y))$. This equation is satisfied by $u_j^1 = \chi_j \cdot \nabla_x u_j$, where χ_j is the first order corrector (1.6). Hence, for any y -independent function $\varphi_1(t, x, y) \equiv \psi(t, x) \in C_0^\infty((0, T) \times \Omega)$,

$$\begin{aligned} 0 &= \int_0^T \int_{\Omega} \int_{Y_j} \sigma_j [\nabla_x u_j + \nabla_y u_j^1] \cdot \nabla_x \psi dx dy dt \\ &= \int_0^T \int_{\Omega} \left(\int_{Y_j} \sigma_j + \nabla_y \chi_j dy \right) \nabla_x u_j \cdot \nabla_x \psi dx dy dt, \end{aligned}$$

so (1.5) is indeed the homogenized conductivity tensor.

Up to the convergences of the nonlinear terms, it is now clear that (4.5), (4.6), and (4.7) constitute the weak formulation of the bidomain system (1.4), (4.1) (in the sense of Definition 4.1).

4.2. The nonlinear terms and strong convergence. To finalize the proof of Theorem 4.3, it remains to identify the limits

$$(4.8) \quad \begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi dS dt &= |\Gamma| \int_0^T \int_{\Omega} I(v, w) \varphi dx dt, \\ \lim_{\varepsilon \rightarrow 0} \varepsilon \int_0^T \int_{\Gamma^\varepsilon} H(v^\varepsilon, w^\varepsilon) \varphi dS dt &= |\Gamma| \int_0^T \int_{\Omega} H(v, w) \varphi dx dt, \end{aligned}$$

for all $\varphi \in C_0^\infty((0, T) \times \Omega)$. We note that the unfolding operator (2.14) allows us to transform the oscillating surface integral

$$\varepsilon \int_0^T \int_{\Gamma^\varepsilon} I(v^\varepsilon, w^\varepsilon) \varphi_i dS dt$$

into

$$\int_0^T \int_{\Omega} \int_{\Gamma} I(\mathcal{T}_\varepsilon^b(v^\varepsilon), \mathcal{T}_\varepsilon^b(w^\varepsilon)) \mathcal{T}_\varepsilon^b(\varphi_i) dS(y) dx dt,$$

coming from the integration formula (2.14) for $\mathcal{T}_\varepsilon^b$. Additionally, we have here used (2.13) and the fact $\mathcal{T}_\varepsilon^b(I(v^\varepsilon, w^\varepsilon)) = I(\mathcal{T}_\varepsilon^b(v^\varepsilon), \mathcal{T}_\varepsilon^b(w^\varepsilon))$. The smoothness of φ implies that $\mathcal{T}_\varepsilon^b(\varphi) \rightarrow \varphi$ in $L^2((0, T) \times \Omega \times \Gamma)$ as $\varepsilon \rightarrow 0$, cf. (2.16), so to identify the limits (4.8) it suffices to show

$$I(\mathcal{T}_\varepsilon^b(v^\varepsilon), \mathcal{T}_\varepsilon^b(w^\varepsilon)) \rightarrow I(v, w), \quad \text{weakly in } L^2((0, T) \times \Omega \times \Gamma),$$

where v, w are identified in (4.4). Besides, since w^ε appears linearly in I and H , cf. (GFHN), the weak convergence of $\{\mathcal{T}_\varepsilon^b(w^\varepsilon)\}_{\varepsilon>0}$ in $L^2((0, T) \times \Omega \times \Gamma)$ is enough to pass to the limit in (4.8), if we establish strong convergence of $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$.

As a step towards verifying the required strong convergence, we need to show that $\{(t, y) \mapsto \mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ is strongly precompact in $L^2_{t,y}$, for fixed $x \in \Omega$. As a result of Lemma 3.8, $\mathcal{T}_\varepsilon^b(v)$ is bounded in $L^2_t L^2_x H_y^{1/2}$, uniformly in ε . However, according to Lemma 3.8, the time derivative $\partial_t v^\varepsilon$ is merely of order $1/\varepsilon$ in the $L^2_t H_x^{-1/2}$ norm. Therefore, we cannot expect $\partial_t \mathcal{T}_\varepsilon^b(v^\varepsilon)$ (assuming that this object is meaningful) to be bounded in $L^2_t L^2_x H_y^{-1/2}$, uniformly in ε . As a consequence, strong $L^2_{t,y}$ compactness of $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ is not deducible from the classical Aubin-Lions theorem. Instead of attempting to control (in a negative space) the whole derivative $\partial_t \mathcal{T}_\varepsilon^b(v^\varepsilon)$, we will make use of a temporal translation estimate with respect to the L^2 norm (cf. lemma below). The $L^2_{t,y}$ compactness will then be a consequence of the Simon lemma (cf. Subsection 2.1).

The rest of this section is devoted to the detailed proof that $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ is strongly precompact in the fixed space $L^2((0, T) \times \Omega \times \Gamma)$. We begin with

Lemma 4.5. *There exists a constant C , independent of ε , such that*

$$(4.9) \quad \|\mathcal{T}_\varepsilon^b(v^\varepsilon)\|_{L^2(0,T;L^2(\Omega;H^{1/2}(\Gamma)))} \leq C$$

and

$$(4.10) \quad \|\mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot + \Delta_t, \cdot, \cdot) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot, \cdot)\|_{L^2(0,T-\Delta_t;L^2(\Omega \times \Gamma))} \leq C\Delta_t^{\frac{1}{2}},$$

for sufficiently small temporal shifts $\Delta_t > 0$.

Proof. By (3.11) and the linearity of $\mathcal{T}_\varepsilon^b(\cdot)$, we have $\mathcal{T}_\varepsilon^b(v^\varepsilon) = \mathcal{T}_\varepsilon^b(u_\varepsilon^\xi|_{\Gamma^\varepsilon}) - \mathcal{T}_\varepsilon^b(u_\varepsilon^\eta|_{\Gamma^\varepsilon})$. Thus, (4.9) follows by integrating (2.21) over $(0, T)$ and using estimates (a), (b) in (3.20). Regarding (4.10), we use the linearity of $\mathcal{T}_\varepsilon^b$, (2.15), and (3.25) to obtain

$$\begin{aligned} & \int_0^{T-\Delta_t} \int_\Omega \int_\Gamma |\mathcal{T}_\varepsilon^b(v^\varepsilon)(t + \Delta_t, x, y) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot, \cdot)|^2 dS(y) dx dt \\ &= \varepsilon \int_0^{T-\Delta_t} \int_{\Gamma^\varepsilon} |v^\varepsilon(t + \Delta_t, x) - v^\varepsilon(t, x)|^2 dS(x) dt \leq C\Delta_t. \end{aligned}$$

□

Let us think of $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ as a function of $x \in \Omega$, with values in $L^2(0, T; L^2(\Gamma))$. Fixing x , in view of Lemma 4.5 and Simon's compactness criterion, the sequence $\{(t, y) \mapsto \mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, y)\}_{\varepsilon>0}$ is precompact in $L^2((0, T) \times \Gamma)$. The x -variable is more difficult. As a matter of fact, since $\mathcal{T}_\varepsilon^b(v^\varepsilon)$ is piecewise constant as a function of x and thus does not belong to any Sobolev space, strong compactness in x is not immediately clear. We address this issue by deriving a translation estimate in the x -variable. To be more precise, we make use of a convenient Simon-type compactness criterion (x playing the role of time) established in [19, Corollary 2.5] (see also [3, Section 5]), which is recalled next.

For $Q \subset \mathbb{R}^n$ and $\xi \in \mathbb{R}^n$ ($n \geq 1$), we set $Q_\xi := Q \cap (Q - \xi) := \{x \in Q : x + \xi \in Q\}$, $\Sigma := \{-1, 1\}^n$ ($|\Sigma| = 2^n$), and $\xi_\sigma := (\xi_1 \sigma_1, \dots, \xi_n \sigma_n) \in \mathbb{R}^n$ for $\sigma \in \Sigma$. If $Q = (a, b)$ ($:= \prod_{\ell=1}^n (a_\ell, b_\ell)$) for $a, b \in \mathbb{R}^n$, $a < b$) is an open rectangle, then

$$(4.11) \quad Q = \bigcup_{\sigma \in \Sigma} Q_{\xi_\sigma}, \quad \text{for any } \xi \in \mathbb{R}^n \text{ such that } Q_\xi \neq \emptyset.$$

Let B be Banach a space. For $f : Q \rightarrow B$ and $\Delta \in \mathbb{R}^n$, we define the translation operator $\tau_\Delta : (Q - \Delta) \rightarrow B$ by

$$\tau_\Delta f(x) = f(x + \Delta).$$

The following theorem, due to Gahn and Neuss-Radu [19], is a multi-dimensional generalization of Simon's main result [37, Theorem 1].

Theorem 4.6 ([19]). *Let $\mathcal{F} \subset L^p(Q; B)$ for some Banach space B , open rectangle $Q = (a, b) \subset \mathbb{R}^n$, and $p \in [1, \infty)$. Then \mathcal{F} is precompact in $L^p(Q; B)$ if and only if*

- i. $\{\int_A f dx \mid f \in \mathcal{F}\}$ is precompact in B , for every open rectangle $A \subset Q$;*
- ii. for each $z \in \mathbb{R}^n$ with $0 < z < b - a$,*

$$(4.12) \quad \sup_{f \in \mathcal{F}} \|\tau_z f - f\|_{L^p(Q_z; B)} \rightarrow 0, \quad \text{as } z \rightarrow 0.$$

Recall (4.11), this time specifying $Q = (a, b) \subset \mathbb{R}^n$, and $\xi = (b - a)/2 \in \mathbb{R}^n$. Condition (4.12) in Theorem 4.6 is equivalent to [19]

$$(4.13) \quad \sup_{f \in \mathcal{F}} \|\tau_{z_\sigma} f - f\|_{L^p(Q_{\xi_\sigma}; B)} \xrightarrow{z \rightarrow 0} 0, \quad z \in \mathbb{R}^n, z \geq 0, \forall \sigma \in \Sigma.$$

The difference between (4.12) and (4.13) is the fixed domain that is utilized in the latter (it does not depend on the shift z). We make use of (4.13) in the proof of Lemma 4.8 below.

We now verify that the sequence $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon > 0}$ of unfolded membrane potentials satisfies the assumptions of Theorem 4.6, with $B = L^2((0, T) \times \Gamma)$, $Q = \Omega$, $p = 2$.

Lemma 4.7 (verification of *i*). *Given an arbitrary open rectangle $A \subset \Omega$, define the function $v_A^\varepsilon(t, y)$ by*

$$v_A^\varepsilon(t, y) = \int_A \mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, y) dx, \quad (t, x) \in (0, T) \times \Gamma.$$

Then the sequence $\{v_A^\varepsilon\}_{\varepsilon > 0}$ is precompact in $L^2((0, T) \times \Gamma)$.

Proof. In view of Jensen's inequality, it follows that

$$\begin{aligned} \|v_A^\varepsilon\|_{L^2(0, T; H^{1/2}(\Gamma))}^2 &= \int_0^T \left\| \int_A \mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, \cdot) dx \right\|_{H^{1/2}(\Gamma)}^2 dt \\ &\leq |A| \int_0^T \int_A \|\mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, \cdot)\|_{H^{1/2}(\Gamma)}^2 dx dt \\ &\leq |A| \|\mathcal{T}_\varepsilon^b(v^\varepsilon)\|_{L^2(0, T; L^2(\Omega; H^{1/2}(\Gamma^\varepsilon)))}^2 \stackrel{(4.9)}{\leq} C. \end{aligned}$$

Let $\Delta_t > 0$ be a small temporal shift. Again using Jensen's inequality,

$$\begin{aligned} &\|v_A^\varepsilon(\cdot + \Delta_t, \cdot) - v_A^\varepsilon(\cdot, \cdot)\|_{L^2(0, T - \Delta_t; L^2(\Gamma))}^2 \\ &\leq |A| \int_0^{T - \Delta_t} \int_A \|\mathcal{T}_\varepsilon^b(v^\varepsilon)(t + \Delta_t, x, \cdot) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(t, x, \cdot)\|_{L^2(\Gamma)}^2 dx dt \\ &\leq |A| \|\mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot + \Delta_t, \cdot, \cdot) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot, \cdot)\|_{L^2(0, T - \Delta_t, L^2(\Omega \times \Gamma))}^2 \stackrel{(4.10)}{\leq} C \Delta_t. \end{aligned}$$

Summarizing, there exists an ε -independent constant C such that

$$\|v_A^\varepsilon\|_{L^2(0, T; H^{1/2}(\Gamma))} \leq C, \quad \|v_A^\varepsilon(\cdot + \Delta_t, \cdot) - v_A^\varepsilon(\cdot, \cdot)\|_{L^2(0, T - \Delta_t; L^2(\Gamma))} \leq C \Delta_t^{1/2}.$$

The lemma follows from these estimates and Simon's compactness criterion. \square

In the next lemma we verify (4.13) with $\xi = (1/2, 1/2, 1/2) \in \mathbb{R}^3$, which is equivalent to condition *ii* in Theorem 4.6.

Lemma 4.8 (verification of *ii*). *Given any $\delta > 0$, there exists $h > 0$ such that for any $\Delta_x \in \mathbb{R}^3$ with $|\Delta_x| < h$ and for all $\varepsilon \in (0, 1]$,*

$$(4.14) \quad \|\mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot + (\Delta_x)_\sigma, \cdot) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot, \cdot)\|_{L^2(\Omega_{\xi_\sigma}; L^2((0, T) \times \Gamma))} < \delta, \quad \forall \sigma \in \Sigma.$$

Proof. We wish to estimate the quantity

$$\begin{aligned} J(\Delta_x; \varepsilon) &:= \left\| \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot + (\Delta_x)_\sigma, \cdot) - \mathcal{T}_\varepsilon^b(v^\varepsilon)(\cdot, \cdot, \cdot) \right\|_{L^2(\Omega_{\varepsilon\sigma}; L^2((0,T) \times \Gamma))}^2 \\ &= \int_0^T \int_{\Omega_{\varepsilon\sigma}} \int_\Gamma \left| v^\varepsilon \left(t, \varepsilon \left\lfloor \frac{x + (\Delta_x)_\sigma}{\varepsilon} \right\rfloor + \varepsilon y \right) - v^\varepsilon \left(t, \varepsilon \left\lfloor \frac{x}{\varepsilon} \right\rfloor + \varepsilon y \right) \right|^2 dS(y) dx dt. \end{aligned}$$

Recall that ε takes values in a sequence $\subset (0, 1]$ tending to zero. Fix any $\varepsilon_0 > 0$. Since the translation operation is continuous in L^2 , there exists $h_0 = h_0(\varepsilon_0) > 0$ such that $J(\Delta_x; \varepsilon) < \delta$ for any $|\Delta_x| < h_0$, for all $\varepsilon \in [\varepsilon_0, 1]$. The rest of the proof is devoted to arguing that this holds also for $\varepsilon \in (0, \varepsilon_0)$, thereby proving (4.14).

Choose $K_{\xi_\sigma}^\varepsilon \subset \mathbb{Z}^3$ such that $\Omega_{\xi_\sigma} = \text{interior} \left(\bigcup_{k \in K_{\xi_\sigma}^\varepsilon} \overline{\varepsilon Y^k} \right)$, $Y^k := k + Y$. Then $J(\Delta_x, \varepsilon)$ becomes

$$\sum_{k \in K_{\xi_\sigma}^\varepsilon} \int_0^T \int_{\varepsilon Y^k} \int_\Gamma \left| v^\varepsilon \left(t, \varepsilon \left\lfloor \frac{x + (\Delta_x)_\sigma}{\varepsilon} \right\rfloor + \varepsilon y \right) - v^\varepsilon \left(t, \varepsilon \left\lfloor \frac{x}{\varepsilon} \right\rfloor + \varepsilon y \right) \right|^2 dS(y) dx dt.$$

If $x \in \varepsilon Y^k$, then $\lfloor \frac{x}{\varepsilon} \rfloor = k$, but we have no useful information about $\left\lfloor \frac{x + (\Delta_x)_\sigma}{\varepsilon} \right\rfloor$. To address this issue, we make use of a favorable decomposition of the cells εY^k proposed in [32] (and also utilized in e.g. [19, 18]).

We decompose each cell εY^k as

$$\varepsilon Y^k = \bigcup_{m \in \{0,1\}^3} \varepsilon Y_\sigma^{k,m}, \quad \varepsilon Y_\sigma^{k,m} := \left\{ x \in \varepsilon Y^k : \varepsilon \left\lfloor \frac{x + \varepsilon \left\{ \frac{(\Delta_x)_\sigma}{\varepsilon} \right\}}{\varepsilon} \right\rfloor = \varepsilon(k + m_\sigma) \right\},$$

for $k \in K_{\xi_\sigma}^\varepsilon$ and $\sigma \in \Sigma$. Regarding the translation, for $x \in \varepsilon Y_\sigma^{k,m}$, we write $(\Delta_x)_\sigma = \varepsilon \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor + \varepsilon \left\{ \frac{(\Delta_x)_\sigma}{\varepsilon} \right\}$, and note that

$$\begin{aligned} \varepsilon \left\lfloor \frac{x + (\Delta_x)_\sigma}{\varepsilon} \right\rfloor &= \varepsilon \left\lfloor \frac{x + \varepsilon \left\{ \frac{(\Delta_x)_\sigma}{\varepsilon} \right\}}{\varepsilon} + \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor \right\rfloor \\ &= \varepsilon \left\lfloor \frac{x + \varepsilon \left\{ \frac{(\Delta_x)_\sigma}{\varepsilon} \right\}}{\varepsilon} \right\rfloor + \varepsilon \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor \\ &= \varepsilon(k + m_\sigma) + \varepsilon \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor. \end{aligned}$$

As a result of this,

$$\begin{aligned} J &= \sum_{k \in K_{\xi_\sigma}^\varepsilon} \sum_{m \in \{0,1\}^3} \int_0^T \int_{\varepsilon Y_\sigma^{k,m}} \int_\Gamma \\ &\quad \times \left| v^\varepsilon \left(t, \varepsilon k + \varepsilon m_\sigma + \varepsilon \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor + \varepsilon y \right) - v^\varepsilon(t, \varepsilon k + \varepsilon y) \right|^2 dS(y) dx dt \\ &\stackrel{\left(\begin{smallmatrix} x := \varepsilon k + \varepsilon y \\ dS(x) = \varepsilon^2 dS(y) \end{smallmatrix} \right)}{\leq} \varepsilon \sum_{k \in K_{\xi_\sigma}^\varepsilon} \sum_{m \in \{0,1\}^3} \int_0^T \int_{\varepsilon(k+\Gamma)} \\ &\quad \times \left| v^\varepsilon \left(t, x + \varepsilon \left(m_\sigma + \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor \right) \right) - v^\varepsilon(t, x) \right|^2 dS(x) dt, \end{aligned}$$

where we have also used $\int_{\varepsilon Y_\sigma^k, m} dx \leq \int_{\varepsilon Y^k} dx = \varepsilon^3$ to arrive at the final line. Since $\sum_{k \in K_{\xi_\sigma}^\varepsilon} \int_{\varepsilon(k+\Gamma)} = \int_{(\Gamma^\varepsilon)_{\xi_\sigma}}$, we conclude that

$$J \leq \varepsilon \sum_{m \in \{0,1\}^3} \int_0^T \int_{(\Gamma^\varepsilon)_{\xi_\sigma}} |v^\varepsilon(t, x+z) - v^\varepsilon(t, x)|^2 dS(x) dt,$$

where the shift $z = z(\Delta_x, \varepsilon, m)$ is $\varepsilon \left(m_\sigma + \left\lfloor \frac{(\Delta_x)_\sigma}{\varepsilon} \right\rfloor \right)$, i.e., z is an integer-multiple of ε . Note that $x+z \in (\Gamma^\varepsilon)_{\xi_\sigma}$ whenever Δ_x and ε are sufficiently small. Recalling the definition (3.11) of v^ε , the trace inequality (2.8) implies

$$\begin{aligned} & \varepsilon \int_0^T \int_{(\Gamma^\varepsilon)_{\xi_\sigma}} |v^\varepsilon(t, x+z) - v^\varepsilon(t, x)|^2 dS(x) dt \\ & \leq C \sum_{j=i,e} \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |u_j^\varepsilon(t, x+z) - u_j^\varepsilon(t, x)|^2 dx dt \\ & \quad + C\varepsilon^2 \sum_{j=i,e} \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |\nabla u_j^\varepsilon(t, x+z) - \nabla u_j^\varepsilon(t, x)|^2 dx dt, \end{aligned}$$

where the last term is bounded by a constant times ε^2 because of (3.20)-(a).

It remains to estimate the term on the second line, which will be done utilizing the well-known characterization of Sobolev spaces by means of translation (difference) operators. Recalling the standard proof of this characterization, a problem that arises (due to the geometry of Ω_j^ε) is that parts of the line segment between x and z may leave Ω_j^ε . To avoid this problem we make use of the interpolation operators (2.22) to obtain functions $Q_\varepsilon^j(u_j^\varepsilon)$ defined on the whole of Ω .

Using the triangle inequality and recalling the estimates (2.23), we obtain

$$\begin{aligned} & \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |u_j^\varepsilon(t, x+z) - u_j^\varepsilon(t, x)|^2 dx dt \\ & \leq \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |u_j^\varepsilon(t, x+z) - Q_\varepsilon^j(u_j^\varepsilon)(t, x+z)|^2 dx dt \\ & \quad + \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |Q_\varepsilon^j(u_j^\varepsilon)(t, x+z) - Q_\varepsilon^j(u_j^\varepsilon)(t, x)|^2 dx dt \\ & \quad + \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |Q_\varepsilon^j(u_j^\varepsilon)(t, x) - u_j^\varepsilon(t, x)|^2 dx dt \\ & \leq C_1 \varepsilon \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |\nabla u_j^\varepsilon|^2 dx dt + C_2 |z| \int_0^T \int_{(\Omega_j^\varepsilon)_{\xi_\sigma}} |\nabla Q_\varepsilon^j(u_j^\varepsilon)|^2 dx dt \\ & \leq C_3 (\varepsilon + |z|) \|\nabla u_j^\varepsilon\|_{L^2(0,T;L^2(\Omega_j^\varepsilon))}^2 \stackrel{(3.20)-(a)}{\leq} C_4 (\varepsilon + |z|). \end{aligned}$$

Hence,

$$J \leq C_5 \varepsilon + C_6 |\Delta_x|,$$

where the constants C_5, C_6 are independent of Δ_x, ε . We select the ε_0 introduced earlier sufficiently small, such that the first term on the right-hand side is $< \delta^2/2$ for all $\varepsilon < \varepsilon_0$. We pick $h_1 > 0$ such that the second term is $< \delta^2/2$ for all $|\Delta_x| < h_1$ (for any $\varepsilon \in (0, 1]$). Specifying $h := \min(h_0, h_1)$, the claim (4.14) now follows. \square

4.3. Concluding the proof of Theorem 4.3. Summarizing, we know that

$$\mathcal{T}_\varepsilon^b(w^\varepsilon) \stackrel{\varepsilon \downarrow 0}{\rightarrow} w \quad \text{in } L^2((0, T) \times \Omega \times \Gamma),$$

because $w^\varepsilon \xrightarrow{2-S} w$, cf. (4.4) and (2.17). Besides, w^ε appears linearly in I and H , cf. (GFHN). We have shown that $\{\mathcal{T}_\varepsilon^b(v^\varepsilon)\}_{\varepsilon>0}$ is strongly precompact in $L^2((0, T) \times \Omega \times \Gamma)$. It then follows that $\mathcal{T}_\varepsilon^b(v^\varepsilon) \rightarrow v$ in $L^2((0, T) \times \Omega \times \Gamma)$ and a.e. in $(0, T) \times \Omega \times \Gamma$, along a subsequence as $\varepsilon \rightarrow 0$ (not relabelled), where v is the two-scale limit of $\{v^\varepsilon\}_{\varepsilon>0}$ identified in (4.4). By way of estimate (d) in (3.20) and (the L^p version of) (2.15),

$$\|\mathcal{T}_\varepsilon^b(v^\varepsilon)\|_{L^4((0, T) \times \Omega \times \Gamma)} = \varepsilon^{1/4} \|v^\varepsilon\|_{L^4((0, T) \times \Gamma^\varepsilon)} \leq C,$$

where C is independent of ε . In view of this estimate and the Vitali convergence theorem, we conclude the validity of (4.8). This finishes the proof of Theorem 4.3.

REFERENCES

- [1] G. Allaire. Homogenization and two-scale convergence. *SIAM J. Math. Anal.*, 23(6):1482–1518, 1992.
- [2] G. Allaire, A. Damlamian, and U. Hornung. Two-scale convergence on periodic surfaces and applications. In A. Bourgeat et al., editor, *Proceedings of the International Conference on Mathematical Modelling of Flow through Porous Media (May 1995)*, pages 15–25. World Scientific Pub., Singapore, 1996.
- [3] H. Amann. Compact embeddings of vector-valued Sobolev and Besov spaces. *Glas. Mat. Ser. III*, 35(55)(1):161–177, 2000.
- [4] M. Amar, D. Andreucci, P. Bisegna, and R. Gianni. A hierarchy of models for the electrical conduction in biological tissues via two-scale convergence: the nonlinear case. *Differential Integral Equations*, 26(9-10):885–912, 2013.
- [5] B. Andreianov, M. Bendahmane, K. H. Karlsen, and C. Pierre. Convergence of discrete duality finite volume schemes for the cardiac bidomain model. *Netw. Heterog. Media*, 6(2):195–240, 2011.
- [6] M. Bendahmane and K. H. Karlsen. Analysis of a class of degenerate reaction-diffusion systems and the bidomain model of cardiac tissue. *Netw. Heterog. Media*, 1(1):185–218, 2006.
- [7] M. Boulakia, M. A. Fernández, J.-F. Gerbeau, and N. Zemzemi. A coupled system of PDEs and ODEs arising in electrocardiograms modeling. *Appl. Math. Res. Express. AMRX*, (2):Art. ID abn002, 28, 2008.
- [8] Y. Bourgault, Y. Coudière, and C. Pierre. Existence and uniqueness of the solution for the bidomain model used in cardiac electrophysiology. *Nonlinear Analysis: Real World Applications*, 10(1):458–482, 2009.
- [9] F. Boyer and P. Fabrie. *Mathematical Tools for the Study of the Incompressible Navier-Stokes Equations and Related Models*. Applied Mathematical Sciences. Springer New York, 2012.
- [10] D. Cioranescu, A. Damlamian, and G. Griso. The periodic unfolding method in homogenization. *SIAM J. Math. Anal.*, 40(4):1585–1620, 2008.
- [11] D. Cioranescu, A. Damlamian, P. Donato, G. Griso, and R. Zaki. The periodic unfolding method in domains with holes. *SIAM J. Math. Anal.*, 44(2):718–760, 2012.
- [12] D. Cioranescu and P. Donato. *An introduction to homogenization*, volume 17 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, New York, 1999.
- [13] P. Colli Franzone, L. F. Pavarino, and S. Scacchi. *Mathematical cardiac electrophysiology*, volume 13 of *MS&A. Modeling, Simulation and Applications*. Springer, Cham, 2014.
- [14] P. Colli Franzone and G. Savaré. Degenerate evolution systems modeling the cardiac electric field at micro- and macroscopic level. In *Evolution equations, semigroups and functional analysis (Milano, 2000)*, volume 50 of *Progr. Nonlinear Differential Equations Appl.*, pages 49–78. Birkhäuser, Basel, 2002.
- [15] P. Donato and K. H. Le Nguyen. Homogenization of diffusion problems with a nonlinear interfacial resistance. *NoDEA Nonlinear Differential Equations Appl.*, 22(5):1345–1380, 2015.
- [16] P. Donato, K. H. Le Nguyen, and R. Tardieu. The periodic unfolding method for a class of imperfect transmission problems. *J. Math. Sci. (N.Y.)*, 176(6):891–927, 2011. Problems in mathematical analysis. No. 58.
- [17] R. FitzHugh. Mathematical models of threshold phenomena in the nerve membrane. *The bulletin of mathematical biophysics*, 17(4):257–278, Dec 1955.
- [18] M. Gahn, M. Neuss-Radu, and P. Knabner. Homogenization of reaction–diffusion processes in a two-component porous medium with nonlinear flux conditions at the interface. *SIAM Journal on Applied Mathematics*, 76(5):1819–1843, 2016.

- [19] M. Gahn and M. Neuss-Radu. A characterization of relatively compact sets in $L^p(\Omega, B)$. *Stud. Univ. Babeş-Bolyai Math.*, 61(3):279–290, 2016.
- [20] I. Graf and M. A. Peter. Diffusion on surfaces and the boundary periodic unfolding operator with an application to carcinogenesis in human cells. *SIAM J. Math. Anal.*, 46(4):3025–3049, 2014.
- [21] E. Grandelius. The bidomain equations of cardiac electrophysiology. Master’s thesis, University of Oslo, 2017.
- [22] C. S. Henriquez and W. Ying. *The Bidomain Model of Cardiac Tissue: From Microscale to Macroscale*. Springer US, Boston, MA, 2009.
- [23] A. L. Hodgkin and A. F. Huxley. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J. Physiol.*, 117(4):500–544, 1952.
- [24] U. Hornung and W. Jäger. Diffusion, convection, adsorption, and reaction of chemicals in porous media. *J. Differential Equations*, 92(2):199–225, 1991.
- [25] J. P. Keener and A. V. Panfilov. A biophysical model for defibrillation of cardiac tissue. *Biophysical Journal*, 71(3):1335–1345, 1996.
- [26] J. P. Keener. The effect of gap junctional distribution on defibrillation. *Chaos*, 8(1):175–187, 1998.
- [27] J. Lions and E. Magenes. *Non-homogeneous boundary value problems and applications*. Number v. 3 in Non-homogeneous Boundary Value Problems and Applications. Springer-Verlag, 1972.
- [28] D. Lukkassen, G. Nguetseng, and P. Wall. Two-scale convergence. *Int. J. Pure Appl. Math.*, 2(1):35–86, 2002.
- [29] A. Marciniak-Czochra and M. Ptashnyk. Derivation of a macroscopic receptor-based model using homogenization techniques. *SIAM J. Math. Anal.*, 40(1):215–237, 2008.
- [30] W. McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, 2000.
- [31] J. C. Neu and W. Krassowska. Homogenization of syncytial tissues. *Crit. Rev. Biomed. Eng.*, 21(2):137–199, 1993.
- [32] M. Neuss-Radu and W. Jäger. Effective transmission conditions for reaction-diffusion processes in domains separated by an interface. *SIAM J. Math. Anal.*, 39(3):687–720, 2007.
- [33] G. Nguetseng. A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.*, 20(3):608–623, May 1989.
- [34] M. Pennacchio, G. Savaré, and P. Colli Franzone. Multiscale modeling for the bioelectric activity of the heart. *SIAM J. Math. Anal.*, 37(4):1333–1370, 2005.
- [35] G. Richardson. A multiscale approach to modelling electrochemical processes occurring across the cell membrane with application to transmission of action potentials. *Mathematical Medicine and Biology: A Journal of the IMA*, 26(3):201–224, 2009.
- [36] G. Richardson and S. J. Chapman. Derivation of the bidomain equations for a beating heart with a general microstructure. *SIAM J. Appl. Math.*, 71(3):657–675, 2011.
- [37] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl. (4)*, 146:65–96, 1987.
- [38] J. Sundnes, G. T. Lines, X. Cai, B. r. F. Nielsen, K.-A. Mardal, and A. Tveito. *Computing the electrical activity in the heart*, volume 1 of *Monographs in Computational Science and Engineering*. Springer-Verlag, Berlin, 2006.
- [39] L. Tung. *A bi-domain model for describing ischemic myocardial D-C potentials*. PhD thesis, MIT, Cambridge, MA, 1978.
- [40] A. Tveito, K. H. Jæger, M. Kuchta, K.-A. Mardal, and M. E. Rognes. A cell-based framework for numerical modeling of electrical conduction in cardiac tissue. *Frontiers in Physics*, 5:48, 2017.
- [41] M. Veneroni. Reaction-diffusion systems for the microscopic cellular model of the cardiac electric field. *Math. Methods Appl. Sci.*, 29(14):1631–1661, 2006.
- [42] M. Veneroni. Reaction-diffusion systems for the macroscopic bidomain model of the cardiac electric field. *Nonlinear Anal. Real World Appl.*, 10(2):849–868, 2009.
- [43] Z. Yang. The periodic unfolding method for a class of parabolic problems with imperfect interfaces. *ESAIM Math. Model. Numer. Anal.*, 48(5):1279–1302, 2014.

(Erik Granelius)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSLO
P.O. BOX 1053, BLINDERN
N-0316 OSLO, NORWAY
E-mail address: erikgran@math.uio.no

(Kenneth Hvistendahl Karlsen)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSLO
P.O. BOX 1053, BLINDERN
N-0316 OSLO, NORWAY
E-mail address: kennethkarlsen@me.com