

# A consistent species richness–climate relationship for oaks across the Northern Hemisphere

Xiaoting Xu<sup>1,2</sup>  | Dimitar Dimitrov<sup>3,4,5</sup>  | Nawal Shrestha<sup>2</sup> | Carsten Rahbek<sup>4,6</sup> | Zhiheng Wang<sup>2</sup>

<sup>1</sup>Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of Life Sciences, Sichuan University, Chengdu, China

<sup>2</sup>Department of Ecology and Key Laboratory for Earth Surface Processes of the Ministry of Education, College of Urban and Environmental Sciences, Peking University, Beijing, China

<sup>3</sup>Natural History Museum, University of Oslo, Oslo, Norway

<sup>4</sup>Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, Copenhagen Ø, Denmark

<sup>5</sup>Department of Natural History, University Museum of Bergen, University of Bergen, Bergen, Norway

<sup>6</sup>Grand Challenges in Ecosystems & the Environment Initiative, Imperial College London, Silwood Park Campus, Ascot, UK

## Correspondence

Zhiheng Wang, Department of Ecology and Key Laboratory for Earth Surface Processes of the Ministry of Education, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China.  
Email: Zhiheng.wang@pku.edu.cn

## Funding information

National Natural Science Foundation of China, Grant/Award Number: #31400467, #31522012, #31470564, #31621091, #31650; Fundamental Research Funds for the Central Universities, Grant/Award Number: #YJ201721; National Key Research Development Program of China, Grant/Award Number: #2017YFA0605101; Danmarks Grundforskningsfond, Grant/Award Number: DNRF96

Editor: Greg Jordan

## Abstract

**Aim:** Although the effects of climate on species richness are known, regional processes may lead to different species richness–climate relationships across continents resulting in species richness anomalies, especially for tropical groups. Phylogenetic niche conservatism may also influence species richness–climate relationships of different lineages. Here, we tested whether regional effects also exist for temperate lineages using the genus *Quercus*.

**Location:** Northern Hemisphere.

**Time period:** Present day.

**Major taxa studied:** *Quercus* (Fagaceae).

**Methods:** We used a dated phylogeny and distribution data for *Quercus* to evaluate its global species richness patterns and phylogenetic niche conservatism. To evaluate the consistency in species richness–climate relationships across continents of the genus *Quercus* as a whole and the temperate subgenus *Quercus*, we conducted analyses of covariance with continent as the categorical variable and climate variables as the covariate. We calibrated four widely used models using the global data or data from each continent separately and evaluated the predictive power of each model for different continents using the root mean squared error.

**Results:** The relationships between species richness and climate were not significantly different among continents for both the genus *Quercus* as a whole and the subgenus *Quercus*. Unlike the models based on European data, those based on North American and eastern Asian data predicted both the global species richness and the richness in other continents. The species richness of a subtropical subgenus *Cyclobalanopsis* was most influenced by water availability, while that of a temperate subgenus *Quercus* was most influenced by environmental temperature.

**Main conclusions:** In contrast to the subtropical subgenus *Cyclobalanopsis*, our results showed a consistent richness–climate relationship and absence of regional effects on species richness across continents for the temperate subgenus *Quercus*, likely suggesting that the species richness of temperate lineages, for example subgenus *Quercus*, may have reached equilibrium with the contemporary climate in the Northern Hemisphere.

## KEYWORDS

climate, eastern Asia, energy, historical effects, niche conservatism, North America, *Quercus*, regional effects, richness anomaly, water

## 1 | INTRODUCTION

Large-scale species richness patterns have intrigued ecologists for more than two centuries (Lomolino, Riddle, Whittaker, & Brown, 2010). The underlying mechanisms of these patterns are not fully understood despite the long history of studies (Currie et al., 2004; Gaston, 2000; Qian & Ricklefs, 2016; Ricklefs, 2004). Environmental variables, including contemporary climate, habitat heterogeneity and past climate change, have been proposed previously as the drivers of contemporary species richness patterns (Currie & Francis, 2004; Kerr & Packer, 1997; Rahbek & Graves, 2001; Tews et al., 2004). However, whether the relationships between species richness and environmental variables are consistent across different continents remains controversial (Currie & Francis, 2004; Latham, Ricklefs, & Schluter, 1994; Qian, Jin, & Ricklefs, 2017; Qian & Ricklefs, 2000; Ricklefs, 1987, 2004; Ricklefs & He, 2016; Ricklefs, Latham, & Qian, 1999).

Based on the strong correlation between species richness and environmental variables, ecologists have argued that contemporary species richness patterns are determined primarily by contemporary climate and habitat heterogeneity (Field, O'Brien, & Whittaker, 2005; O'Brien, Field, & Whittaker, 2000) and that there is a globally consistent species richness–climate relationship. For example, Currie and Paquin (1987) built a logistic model between species richness and actual annual evapotranspiration using the distributions of North American trees, and they demonstrated that this model could predict the tree species richness in Great Britain and Ireland. O'Brien and colleagues (Field et al., 2005; O'Brien, 1998, 2006; O'Brien et al., 2000) proposed the water–energy dynamic hypothesis to predict tree species richness in South Africa. Francis and Currie (2003) further evaluated this hypothesis and showed that it could explain over 87% of the variation in angiosperm family richness and is consistent globally and independent of evolutionary history and plant functional assemblages. Wang, Fang, Tang, and Lin (2011a) indicated that winter coldness was the primary driver of woody plant richness in eastern Asia because winter coldness exerted strong effects on species with tropical affinity and proposed a model including winter coldness and other variables representing water, temperature seasonality and habitat heterogeneity. This model predicted the tree species richness in North America with high accuracy. Although these models differ in the choice of variables, they all have high explanatory power in describing species richness variation and suggest a globally consistent correlation between species richness and climate.

Species richness anomalies occur when regions with similar environments show significantly different species richness (Latham & Ricklefs, 1993; Latham et al., 1994). A typical example is eastern Asia, which has a similar climate to that of eastern North America but has significantly higher species richness after accounting for environmental effects (Guo & Ricklefs, 2000; Qian et al., 2017; Qian & Ricklefs, 2000;

Ricklefs, Qian, & White, 2004). The mechanisms underlying the eastern Asian anomaly have been attributed to its distinct geological, climatic and evolutionary history (Qian et al., 2017; Ricklefs et al., 1999; Ricklefs & Renner, 1994; Xiang et al., 2004). For example, the uplift of the Qinghai-Tibetan Plateau and the resulting heterogeneous topography in south-west China and monsoon system in East Asia (Molnar, Boos, & Battisti, 2010) have provided ample opportunities for species in eastern Asia to diversify and survive the dramatic climatic fluctuations since the mid-Cenozoic (Wen, Zhang, Nie, Zhong, & Sun, 2014; Xing & Ree, 2017). Lineage speciation rates have been found to be higher in eastern Asia than in North America due to complex topography and long-term vegetation stability in the former (Hughes & Atchison, 2015; Qian & Ricklefs, 2000; Xiang et al., 2004). In contrast, the increased seasonality and aridity in North America during the Tertiary (i.e., the Eocene–Oligocene climate cooling) and the Quaternary may have led to the extinction of many tropical groups due to their phylogenetic niche conservatism constraining them to warm and wet climate (Latham et al., 1994; Prothero, 1994; Qian, 2002; Qian & Ricklefs, 2000). The large extent of ice sheets in North America during the Last Glacial Maximum (LGM) could also have lasting effects on the equilibrium between species richness and climate. The species richness anomaly in eastern Asia has been found primarily in old tropical groups (Latham et al., 1994; Qian et al., 2017), and whether it also exists for temperate groups remain controversial (but see Latham et al., 1994).

Here, we used the widely distributed temperate genus *Quercus* L. to test the consistency of the species richness–climate relationships and the species richness anomaly among continents after accounting for environmental effects. The genus *Quercus* is classified into two subgenera, the subtropical subgenus *Cyclobalanopsis* and temperate subgenus *Quercus* (Wu, Raven, & Hong, 1999). The subgenus *Cyclobalanopsis* has c. 150 species with distribution restricted to East and Southeast Asia. The temperate subgenus *Quercus* has the highest species richness in North America and includes c. 300 species widely distributed in the Northern Hemisphere. The subgenus *Quercus* includes five groups: group *Quercus*, group *Lobatae*, group *Protobalanus*, group *Cerris* and group *Ilex* (Nixon, 1997; Wu et al., 1999). Phylogenetic studies suggest that the group *Quercus*, group *Lobatae* and group *Protobalanus* form a monophyletic clade, which is sister to the clade containing group *Cerris*, group *Ilex* and subgenus *Cyclobalanopsis* (Deng, Jiang, Hipp, Manos, & Hahn, 2018; Denk & Grimm, 2009, 2010; Hipp et al., 2018; Manos & Stanford, 2001). Many *Quercus* species are important components of forest ecosystems in temperate regions in the Northern Hemisphere (Nixon, 1997; Wu et al., 1999). Previous studies on the species richness patterns of *Quercus* indicate that the species richness–climate relationship of this group is rooted in its evolutionary history due to niche conservatism (Xu, Wang, Rahbek, Lessard, & Fang, 2013) and that a water–energy interaction significantly influences *Quercus* species richness (Xu, Wang, Rahbek, Sanders, & Fang, 2016).

In the present study, we analysed the species richness patterns of the genus *Quercus*, especially of the temperate subgenus *Quercus* in different continents across the Northern Hemisphere. We assessed the effects of contemporary climate, climate change since the LGM and topography on richness patterns at continental and global scales as well as testing the consistency of the species richness–climate relationship across continents. We assessed the regional effects on species richness using continents as a variable to reflect the long isolation and independent geological history of each continent.

## 2 | DATA AND METHODS

### 2.1 | Species distribution data

*Quercus* species distribution data were obtained from the global *Quercus* distribution database (Xu et al., 2013, 2016), which includes over 20,000 distribution records for c. 400 *Quercus* species collected from published floras, checklists, online databases and herbarium specimens. The intraspecific taxa were merged to species level. Island species and hybrids [hybridization status according to Govaerts et al. (2011)] were excluded. The final dataset included distribution data for 376 species (Supporting Information Appendix S1). Among the 376 species, 91 belong to the subgenus *Cyclobalanopsis* and 285 belong to the subgenus *Quercus* (Denk & Grimm, 2009, 2010; Menitsky, 2005; Nixon, 1997; Rangel, 2006; Rangel, Carlos, Zenteno, de Lourdes, & Enríquez, 2002; Wu et al., 1999). Of the 285 species in the subgenus *Quercus*, 220 species occur in North America and 65 species occur in Eurasia. The species distributed in North America belong to the endemic group *Lobatae* (108 species), group *Protobalanus* (5 species) and group *Quercus* (107 species). Similarly, the species distributed in Eurasia belong to group *Quercus* (25 species), group *Ilex* (29 species) and group *Cerris* (12 species). All the 91 species in the subgenus *Cyclobalanopsis* are restricted to eastern Asia (Figure 1).

Species distributions were compiled based on global administrative divisions, also known as country subdivisions. Global administrative areas (GADM) were downloaded from the GADM database (<https://gadm.org/data.html>, accessed June 2010). To control the effects of area on species richness, neighbouring divisions with area <100,000 km<sup>2</sup> were merged into larger units according to data availability. Cambodia, which shows high data deficiency, was excluded from our analysis. The final map included 164 geographical units with a mean area of 210,616 ± 49,728.96 km<sup>2</sup>, and 70% areas between 180,000 and 250,000 km<sup>2</sup> (Figure S1a in Supporting Information Appendix S2). Among the 164 regions, 56 geographical units are in eastern Asia, 52 are in Europe and 56 are in North America (Figure 1). Area of geographical units is not significantly correlated with *Quercus* species richness at continental or global scales ( $p > 0.05$ , Figure S1b in Supporting Information Appendix S2). Species richness was estimated as the total number of species in each geographical unit.

### 2.2 | Environmental data

To estimate the effects of contemporary climate, habitat heterogeneity and past climate change on *Quercus* species richness, we


selected variables that have been widely used in previous studies and are easily interpretable in a biological context. Contemporary climate variables were grouped into three types representing environmental energy availability, water availability and climate seasonality (Table 1). Energy availability variables included mean annual temperature (MAT, °C), mean temperature of the coldest quarter (MTCQ, °C), potential evapotranspiration (PET, mm) and minimum monthly potential evapotranspiration (PETmin, mm). Water availability variables included mean annual precipitation (MAP, mm), actual evapotranspiration (AET, mm), rainfall (Rainfall, mm) and water deficit (WD, mm). Climate seasonality variables included annual range of temperature (ART, °C), temperature seasonality (TSN) and precipitation seasonality (PSN). Habitat heterogeneity included range of altitude (RALT, m), range of mean annual temperature (RMAT, °C) and range of mean precipitation (RMAP, mm) within each geographical unit. Past climate change was represented by temperature anomaly (TA, °C) and precipitation anomaly (PA, mm).

MAT, MTCQ, MAP, altitude and LGM climate data were downloaded from the WorldClim database ([www.worldclim.org](http://www.worldclim.org)) at a spatial resolution of 2.5 × 2.5 arc minutes. PET and AET with spatial resolutions of 30 arc seconds were obtained from the Consortium of International Agricultural Research Centers consortium for spatial information (<http://www.cgiar-csi.org/>). PET was calculated using the Hargreaves method, which requires less parameterization and reduces sensitivity to climatic input errors (Fisher, Whittaker, & Malhi, 2011; Trabucco, Zomer, Bossio, Straaten, & Verchot, 2008). WD was calculated as the difference between PET and AET. PETmin was calculated as monthly minimum PET of a year. Rainfall was calculated as the sum of monthly precipitation values when mean monthly temperature was above 0°C (Francis & Currie, 2003). RALT, RMAT and RMAP were calculated as the range of altitude, mean annual temperature and mean annual precipitation within each geographical unit, respectively. RALT, RMAT and RMAP were log transformed for analysis to improve the linearity and normality of the model residuals. TA and PA were calculated as the modern minus the LGM MAT and MAP, respectively. Due to the uncertainty of past climate models, we used the mean and median value of three available LGM simulations based on the MIROC-ESM (The Earth system model, Model for Interdisciplinary Research on Climate-Earth system model), MPI-ESM-P (Max-Planck-Institute Earth System Model) and CCSM4 (Community Climate System Model version 4) models. The mean and median values showed consistent results. The results based on median values are shown in Table S1 in Supporting Information Appendix S2. Correlation coefficients between variables are shown in Table S2 in Supporting Information Appendix S2.

## 3 | DATA ANALYSES

### 3.1 | Phylogenetic niche conservatism of *Quercus*

To determine whether the species richness–climate relationship is mediated by climatic niche conservatism, as suggested by Xu et al. (2013), we reconstructed ancestral climatic niches



**FIGURE 1** (a) Global patterns of *Quercus* species richness and species composition on each continent (the species richness data were updated from Xu et al., 2016). (b–g) Global patterns of the species richness for the six groups of *Quercus* [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

and estimated the phylogenetic niche conservatism for each species group present on the same continent. A consensus supertree of *Quercus* was built by merging two recently published *Quercus* phylogenies (Deng et al., 2018; Hipp et al., 2018) based on RAD sequences. In a genus like *Quercus* where hybridization is common, the use of restriction site associated DNA sequencing (RAD seq) approach likely provides more reliable estimates

of phylogenetic relationships among species than traditional approaches targeting a handful of loci (Wagner et al., 2013). The tree published by Hipp et al. (2018) includes 146 North American species, and 4 Eurasian species. The other tree published by Deng et al. (2018) includes 42 Eurasian species (including 35 species of subgenus *Cyclobalanopsis*) and 3 North American species. These two trees were merged directly based on overlapping species



**TABLE 1** Climate variables and their abbreviations used in the analyses

Groups	Abbreviations	Environmental variables
Energy	MAT	Mean annual temperature (°C)
	MTCQ	Mean temperature of coldest quarter (°C)
	PET	Annual potential evapotranspiration (mm)
	PETmin	Minimum monthly potential evapotranspiration (mm)
Water	MAP	Mean annual precipitation (mm)
	AET	Annual evapotranspiration (mm)
	WD	Mean annual precipitation (mm)
	Rainfall	Sum of monthly precipitation values for which mean monthly temperature was above 0 (°C)
	PCQ	Precipitation of coldest quarter (mm)
Seasonality	TSN	Temperature seasonality
	ART	Annual range of temperature (°C)
	PSN	Precipitation seasonality
Past climate change	TA	Temperature anomaly (°C)
	PA	Precipitation anomaly (mm)
Habitat heterogeneity	RALT	Range of altitude (m) within each geographical unit
	RMAT	Range of mean annual temperature (°C) within each geographical unit
	RMAP	Range of mean annual precipitation (mm) within each geographical unit

(Bininda-Emonds, 2004; Gordon, 1986). Two species from the North American tree were removed in the consensus supertree because one species was not identified and the other (*Quercus baronii*) showed a conflicting branch length with the Eurasian tree. In total, our consensus supertree contains 188 species (Supporting Information Appendix S4). The crown age of genus *Quercus* and the divergence time between one clade containing group *Quercus*, group *Lobatae*, group *Protobalanus* and the other clade containing group *Cerris*, group *Ilex* and subgenus *Cyclobalanopsis* in the consensus supertree were extracted from the North American tree, which covers more species and was dated using older fossils compared to the Eurasian tree.

Ancestral climatic niches were reconstructed for group *Lobatae*, group *Quercus*, group *Cyclobalanopsis* and the whole North American

clade (group *Lobatae*, group *Quercus* and group *Protobalanus*) using the “ace” function in the R package ape (Paradis, Claude, & Strimmer, 2004). Blomberg’s *K* (Blomberg, Garland, & Ives, 2003), as implemented by phyloSignal in the R package phylosignal (Keck, Rimet, Bouchez, & Franc, 2016), was employed to quantify the phylogenetic signal of each climatic variable because it is insensitive to sample size (Munkemuller et al., 2012). Blomberg’s *K* was estimated using the subtrees for each *Quercus* group. Higher *K*-values indicate stronger phylogenetic niche conservatism (i.e., close relatives are more similar in their climatic niches than expected based on phylogenetic relationships). The phylogenetic signal of all climatic variables was compared among clades/groups. Because evolution mode could influence the conservatism of phenotypic traits, we fitted three evolution models to each climatic variable to investigate the mode of evolution of climatic niches (Blomberg et al., 2003) using the R package Geiger (Harmon, Weir, Brock, Glor, & Challenger, 2008). The three models were the (a) Brownian motion model (BM) (Felsenstein, 1973), (b) Ornstein–Uhlenbeck model (OU) (Butler & King, 2004) and (c) early-burst model (EB) (Harmon et al., 2010). Climatic variables of each species were calculated as the mean values within their distribution ranges (Supporting Information Appendix S3). The mean values of mean annual temperature, mean annual precipitation, annual range of temperature and mean elevational range for each species were calculated and shown on the tree using the barplot.phylo4d function in the R package phylosignal (Keck et al., 2016).

### 3.2 | Richness–climate relationships among continents

To evaluate the consistency of species richness–climate relationships among North America, Europe and eastern Asia, we built both bivariate and multiple regression models for each continent separately and at global level. Bivariate regression models help detect the best single variable for explaining *Quercus* species richness on each continent. The variables that showed a significant effect on *Quercus* species richness in bivariate regressions at global level were further tested for their significant differences in slopes and intercepts among continents using analysis of variance (ANOVA). In particular, ANOVA was performed for each climatic variable, in which species richness was used as the response variable, continent (represented by eastern Asia, North America and Europe) as the categorical variable, and each climate variable as the covariate, respectively. We also included the interaction term of the climate variable and continent in the ANOVA models. The spatial autocorrelation in species richness could inflate type I error and, consequently, the significance levels of the statistical tests for regression and ANOVA models. Therefore, a modified *t* test was conducted in R following Dutilleul, Clifford, Richardson, and Hemon (1993) for significance testing of the models. We also conducted simultaneous autoregressive (SAR) models with spatially structured errors to account for spatial autocorrelation in our data (Kissling & Carl, 2008). A series of models with spatial weights at distances ranging

from 400 to 2,500 km at 100-km intervals were conducted, and the one with the lowest spatial autocorrelation in the residuals and minimum Akaike information criterion (AIC) value was selected as the best model. Spatial autocorrelation in model residuals was estimated using Moran's *I*. Spatial statistics were performed with the "spdep" library in R (Bivand & Wong, 2018; R Core Team, 2018). SAR models showed consistent results (see Table S3 in Supporting Information Appendix S2) with the bivariate regression models.

We constructed separate multiple regression models based on our global *Quercus* dataset and the datasets for Europe, North America and eastern Asia. We selected four previously proposed models for the relationships between species richness and climate. The first three models were (a) Richness ~ Rainfall + PETmin - PETmin<sup>2</sup> + log(RALT) proposed by O'Brien et al. (2000), (b) Richness ~ WD + PETmin - PETmin<sup>2</sup> proposed by Francis and Currie (2003), and (c) Richness ~ MTCQ + WD + TSN + MATR proposed by Wang, Fang, Tang, and Lin (2011b) (Table 1). The fourth model included one water variable, one energy variable and their interaction term according to previous findings that the interaction of water and energy have a significant effect on *Quercus* species richness (Xu et al., 2016). To identify the best water and energy variable for the fourth model, we tested all combinations of water and energy variables and selected the one with the highest explanatory power globally and at the continent level. Finally, we constructed the fourth model as Richness ~ Rainfall + PETmin + Rainfall × PETmin. Due to the importance of mean annual precipitation on spatial patterns of species richness, we also built two models in which Rainfall was replaced by mean annual precipitation (i.e., precipitation models). The results of the precipitation models were consistent with those of the Rainfall models and are shown in Tables S4 and S5 in Supporting Information Appendix S2.

To test for consistency in the richness–climate relationship, we calibrated each of the six models at the global scale and separately for each continent and used them to predict the global species richness and that of the other continents. We then calculated the root mean squared error (RMSE) and adjusted  $r^2$  for each prediction to assess the predictive power of the models. Lower RMSEs indicate higher predictive power and a higher level of consistency in the richness–climate relationship between the predicted continent and the continent used to build the model.

## 4 | RESULTS

The six groups of oaks showed distinct distributions and species richness patterns in eastern Asia, Europe and North America (Figure 1). The oak species richness in eastern Asia was dominated by the subtropical subgenus *Cyclobalanopsis*, but was dominated by temperate groups in North America and Europe. The total number of species in North America was higher than in eastern Asia. Ancestral climate reconstruction indicated that the ancestral MAT for the group *Cyclobalanopsis* was approximately 19°C, which was c. 3–5°C higher than that of the other groups (Figure 2). Compared to the climatic niche of the subgenus *Cyclobalanopsis*, the climatic niches of

the group *Quercus* and group *Lobatae* were significantly conserved with most *K* values larger than 1 (Table 2).

Simple regressions showed that water availability explained most of the variations in the species richness of the subtropical subgenus *Cyclobalanopsis* (Table 3). For the temperate subgenus *Quercus*, climate seasonality and past temperature change (TA) explained most variations in species richness in eastern Asia, while energy and habitat heterogeneity explained most variations in species richness in North America and Europe, respectively (Table 3). After accounting for the spatial autocorrelation in species richness, the slopes or intercepts of these simple regressions were not significantly different among continents for both the subgenus *Quercus* and the whole genus, which suggests that there is a consistent species richness–climate relationship among continents for oaks (Figure 3).


Multiple-regression models showed that all models have a lower  $r^2$  at the global scale than at the continental scale (Table 4). All four models showed a relatively high explanatory power in different continents ( $r^2_{\text{adj}}$  from 0.571 to 0.854). Moreover, the  $r^2$  of the four models was higher in eastern Asia ( $r^2_{\text{adj}} \geq 0.721$ ) than in North America ( $r^2_{\text{adj}} \geq 0.571$ ) and Europe ( $r^2_{\text{adj}} \geq 0.691$ ). The Francis & Currie model (the second model) best predicted species richness of eastern Asia ( $r^2_{\text{adj}}$  0.822). The water–energy interaction model best predicted species richness in North America ( $r^2_{\text{adj}}$  0.799), and the water–energy dynamic model best predicted richness in Europe ( $r^2_{\text{adj}}$  0.854). However, for the subgenus *Quercus*, the  $r^2$  of the four models ( $r^2_{\text{adj}} \geq 0.343$ ) are lower than that of the whole genus. In eastern Asia, the  $r^2$  of the four models for subgenus *Quercus* are higher than those for subgenus *Cyclobalanopsis* (Table S5 in Supporting Information Appendix S2).

Among the four models, the best global model was the Xu et al. (2016) model, which gave the highest explanatory power of species richness variation (whole genus,  $r^2_{\text{adj}} = 0.710$ ,  $p < 0.001$ , subgenus *Quercus*,  $r^2_{\text{adj}} = 0.581$ ,  $p < 0.001$ , Table 4 in maintext and Table S5 in Supporting Information Appendix S2) and could predict species richness in other continents with the lowest RMSEs compare to other three models (Figure 4). However, when we used EU models (the model built with European data) to predict eastern Asia, North America and global *Quercus* species richness, we found that the models built by Xu et al. (2016), O'Brien et al. (2000) and Francis and Currie (2003) showed larger RMSE values than did the Wang et al. (2011b) model (Figure 4). This difference is probably due to the narrower range of climate regimes in Europe, which limits the extrapolative power of nonlinear regression models based on the European data alone (Figure 5).

## 5 | DISCUSSION

### 5.1 | Consistent oak richness–climate relationships across the Northern Hemisphere

Despite differences in the species composition of *Quercus* flora among continents, we did not find significant regional effects on the species richness of the entire genus and the temperate



**FIGURE 2** Dated phylogenetic tree of the genus *Quercus* compiled from Hipp et al. (2000) and Deng et al. (2009). Species group membership, continental distribution, the mean annual temperature (MAT, °C), mean annual precipitation (MAP, mm), annual range of temperature (ART, °C) and range of altitude (RALT, m) are shown for each species. Range of altitude of a species was calculated as the mean of the altitudinal ranges of all geographical units in which the species occurs. The red figures besides the deep nodes represent the reconstructed ancestral temperature of these nodes [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Environmental variables		<i>Cyclobalanopsis</i>	<i>Lobatae</i>	<i>Quercus</i>	<i>Cerris + Ilex</i>	Full tree
		<i>n</i> = 35	<i>n</i> = 54	<i>n</i> = 87	<i>n</i> = 8	<i>n</i> = 188
Energy	MAT	0.223	2.063***	1.405***	0.486	0.462***
	MTCQ	0.216	2.738***	1.383***	0.831	0.503***
	PET	0.233	3.001***	1.921***	0.337	0.782***
	PETmin	0.229	3.841***	2.015***	0.858*	0.732***
Water	MAP	0.199	0.780**	1.002***	0.902	0.566***
	AET	0.271	0.820*	1.189***	1.003	0.791***
	Rainfall	0.200	0.799***	1.067***	0.928	0.593***
	PCQ	0.250	1.073***	0.807***	1.032*	0.556***
Seasonality	TSN	0.150	0.518	0.576*	0.870	0.177
	ART	0.215	2.617***	1.086***	0.856*	0.540***
	PSN	0.250	5.133***	1.459***	0.469	1.118***

Note. The North America groups, group *Lobatae* and group *Quercus*, showed significant phylogenetic signals in their climatic niches. *n* indicates the number of species sampled in the phylogenetic tree. Please see Table 1 for definitions of abbreviations.

\*\*\**p*-value < 0.001; \*\**p*-value < 0.01; \**p*-value < 0.1.

subgenus *Quercus*. This finding suggests that the eastern Asian species richness anomaly is not present in this group. Species anomalies between eastern Asia and eastern North America mainly exist for Tertiary relict lineages, which were extirpated from North America in the late Tertiary due to climate cooling and extensive glaciation but have survived in eastern Asia (Guo & Ricklefs, 2000; Qian et al., 2017; Qian & Ricklefs, 2000; Xiang et al., 2004). Temperate lineages, such as the subgenus *Quercus*, which diversified during this cooling period, have relatively higher species richness in North America than in eastern Asia. Other temperate genera (e.g., *Carya*) also have higher species richness in North America than in eastern Asia (Latham et al., 1994; Xiang et al., 2004). Our results, combined with previous findings, suggest that the species richness anomaly between North America and eastern Asia might not exist for temperate lineages.

Our results showed that *Quercus* species richness was a function of energy, water and their interaction (Table 4), which conforms with the tree species richness–climate models proposed in previous studies (Field et al., 2005; Francis & Currie, 2003; Hawkins et al., 2003; O'Brien, 1998; O'Brien et al., 2000; Whittaker, Nogues-Bravo, & Araujo, 2007; Xu et al., 2016). For example, in a series of classic studies, O'Brien and colleagues found that the species richness of woody plants in South Africa was a function of rainfall, PETmin and PETmin<sup>2</sup> due to higher evapotranspiration rates in warm regions (Field et al., 2005; O'Brien, 1998; O'Brien et al., 2000). However, we acknowledge that our study was limited to a typical temperate woody plant group in the Northern Hemisphere and therefore, further studies are needed to better understand the regional effects on species richness patterns of temperate clades. Although these studies support the importance of water, energy and their interaction on individual growth and distribution, the underlying physiological processes and the mechanistic link between individual growth and the general species richness pattern remain unknown.

**TABLE 2** Phylogenetic signals for different climatic variables evaluated using Bloomberg's *K*

## 5.2 | Phylogenetic niche conservatism and species richness–climate relationships

The current consistent *Quercus* richness–climate relationship across continents may be the result of the evolutionary history of this genus (Xu et al., 2013). According to the niche conservatism hypothesis, species tend to track their ancestral niches and have difficulties in adapting to new environments (Wiens et al., 2010). When climate deviates from its ancestral climate, the constraints on species richness induced by climate tend to be strengthened, and the species richness of a clade normally decreases. Moreover, groups with stronger niche conservatism have difficulties in adapting to new climates and may be more strongly limited by climate (Wiens & Donoghue, 2004). Recent studies on China's woody plants have found that winter temperature determines the latitudinal gradient of species richness by filtering out species with tropical affinities from the floras north of the tropics due to the tropical niche conservatism of these groups. The explanatory power of winter temperature on species with tropical affinity is higher than for those with temperate affinity (Wang et al., 2011b). Previous studies on *Quercus* have also found that the effect of environmental energy on the species richness of different groups depends on their ancestral niches; environmental energy limits species richness of tropical lineages at high latitudes but of temperate groups at both low and high latitudes (Xu et al., 2013). Here, we found stronger phylogenetic conservatism in the climatic niches represented by energy, water and seasonality variables for the temperate groups *Quercus* and *Lobatae* and the entire genus than for tropical group *Cyclobalanopsis*. Thus, niche conservatism may be one explanation for the globally consistent *Quercus* richness–climate relationship.


For a group with strong niche conservatism, adaptation to similar climate across its species tends to lead to a consistent species richness–climate relationship across continents. For the temperate



**TABLE 3** Adjusted  $r^2$  for the predictors of species richness patterns of oaks in eastern Asia, North America, Europe and the whole world evaluated by bivariate regression models

Environmental variables	Eastern Asia			North America		Europe		Global	
	Genus <i>Quercus</i>	Subgenus <i>Quercus</i>	Subgenus <i>Cyclobalanopsis</i>	Subgenus <i>Quercus</i>	Subgenus <i>Quercus</i>	Subgenus <i>Quercus</i>	Subgenus <i>Quercus</i>	Genus <i>Quercus</i>	Subgenus <i>Quercus</i>
Energy									
MAT	0.480**	0.220	0.170*	0.512**	0.185	0.426***	0.314***		
MTCQ	0.606**	0.291	0.233*	0.413***	0.325*	0.460***	0.342***		
PET	0.331*	0.117	0.043	0.540**	0.096	0.389***	0.331***		
PETmin	0.435	0.125	0.156	0.374*	0.106	0.413***	0.278**		
Water									
MAP	0.424*	0.140	0.335***	0.006	0.054	0.213**	0.098*		
AET	0.586**	0.281	0.552***	0.117	0.107	0.354***	0.214**		
WD	0.010	0.012	0.154***	0.111	0.002	0.010	0.027		
Rainfall	0.456*	0.158	0.369	0.039	0.233**	0.268**	0.131**		
PCQ	0.186*	0.0646	0.077	0	0.354***	0.010	0.023		
Seasonality									
TSN	0.682*	0.344	0.249**	0.242	0.390**	0.413***	0.307**		
ART	0.750**	0.394	0.438	0.158	0.286*	0.345***	0.227**		
PSN	0.257*	0.211*	0.168	0.093*	0.007	0.003	-0.006		
Past climate change									
TA	0.734**	0.554**	0.101**	0.339	0.269*	0.112	0.045		
PA	0.104**	0.057*	0.104	0.040	0.283**	0.043*	0.025		
Habitat heterogeneity									
log(RALT)	0.062	0.079	-0.026	-0.010	0.553***	0.111*	0.088*		
log(RMAT)	0.049	0.067	-0.007	-0.009	0.579***	0.115*	0.094*		
log(RMAP)	0.392	0.156	-0.004	0.021	0.527***	0.260***	0.166**		

Note. Subgenus *Cyclobalanopsis* was only distributed in Eastern Asia. Corrected  $p$ -values calculated using modified  $t$  tests (Dutilleul et al., 1993). Please see Table 1 for definitions of abbreviations. \*\*\* $p$ -value < 0.01; \*\*corrected  $p$ -value < 0.05; \* $p$ -value < 0.1.



**FIGURE 3** Comparisons of the intercepts and slopes of species richness–climate regressions among continents for the genus *Quercus* and subgenus *Quercus*. Different letters on top of the bars indicate significant differences in slopes or intercepts between continents. After accounting for spatial autocorrelation, all slopes of richness–climate regressions were not significant among continents. In Europe, the intercepts of the regressions between richness and annual temperature range and range of mean annual precipitation are significantly lower than those in North America. MAT = mean annual temperature (°C); MTCQ = mean temperature of coldest quarter (°C); PET = annual potential evapotranspiration (mm); PETmin = minimum monthly potential evapotranspiration (mm); MAP = mean annual precipitation (mm); AET = annual evapotranspiration (mm); Rainfall = rainfall (mm); TSN = temperature seasonality; ART = annual range of temperature; log(RMAP) = log-transformed mean annual precipitation

subgenus *Quercus*, fossil records suggested its diversification in the Oligocene in North America when climate became cooler and dryer. Species from this subgenus might have dispersed from North America to Eurasia through the Bering and the potential Atlantic land bridges during glaciation periods (Denk, Grimsson, & Zetter, 2010, 2012). Its strong niche conservatism may have constrained the species of this group to similar climates across continents, leading to consistent richness–climate relationships in different continents.

The high species richness of the groups *Quercus* and *Lobatae* is likely the result of the expansion of temperate forests and grasslands in North America due to continuous cooling and drying since the Oligocene (Axelrod, 1983). Dating results showed that the crown age of these two groups was c. 34 Ma, which corresponds to the Eocene–Oligocene climate cooling and drying. Fossil records also provide evidence that species from group *Quercus* and group *Lobatae* were widely distributed in high latitude regions of the Northern Hemisphere (Denk, Grimsson, & Zetter, 2010, 2012). When compared to *Cyclobalanopsis*, the species from these two groups show generally higher phylogenetic niche conservatism, which probably

reflects their adaptations to seasonal climate, resulting in a strong species–climate relationship.



Extinctions within a group would lead to low species richness of this group in some continents, and hence species diversity anomalies across continents. The earliest fossils with affinities to the group *Cyclobalanopsis* were found in North America in deposits from the Middle Eocene, when forests covered North America and the climate was warm and wet (Manchester, 1994). Subsequent cooling and drying during the Oligocene probably led to the regional extirpation of this group from North America (Axelrod, 1983). Currently, this group is confined to wet and warm subtropical and tropical eastern Asia where water availability is the main limiting factor for its species richness. It seems that adaptations to wet and warm climate may have give rise to the richness–climate relationship for subgenus *Cyclobalanopsis*. However, controversy exists between fossil and molecular evidence on the origination of the subgenus *Cyclobalanopsis* (e.g., Deng et al., 2018) and further studies are needed to solve the evolutionary history of this clade.

We did not detect significant phylogenetic signals for climatic niche variables in the subgenus *Cyclobalanopsis*. Currently, we have

**TABLE 4** The regression coefficients (Coef.), *p*-values (*p*) and adjusted  $r^2$  (adj.  $r^2$ ) of the four multiple regression models built using global data or data from each continent

Model type	Predicts	Global			Eastern Asia			North America			Europe		
		Coef.	<i>p</i>	adj. $r^2$	Coef.	<i>p</i>	adj. $r^2$	Coef.	<i>p</i>	adj. $r^2$	Coef.	<i>p</i>	adj. $r^2$
O'Brien et al. (2000)	Intercept	0.138	0.823	0.610	-0.415	0.740	0.755	5.895	<10 <sup>-5</sup>	0.695	-2.110	<10 <sup>-4</sup>	0.854
	Rainfall	0.001	<10 <sup>-4</sup>		0.001	0.045		-0.0004	0.068		0.002	<10 <sup>-4</sup>	
	PETmin	0.078	<10 <sup>-5</sup>		0.083	<10 <sup>-5</sup>		0.097	<10 <sup>-5</sup>		0.109	<10 <sup>-5</sup>	
	PETmin <sup>2</sup>	-0.001	<10 <sup>-5</sup>		-0.001	<10 <sup>-5</sup>		-0.001	<10 <sup>-5</sup>		-0.213	<10 <sup>-5</sup>	
	log(RALT)	-0.012	0.893		0.068	0.681		-0.745	<10 <sup>-5</sup>		0.002	0.015	
Francis and Currie (2003)	Intercept	0.709	<10 <sup>-5</sup>	0.665	0.794	<10 <sup>-5</sup>	0.822	0.674	0.012	0.571	0.252	0.040	0.819
	WD	-0.001	<10 <sup>-5</sup>		-0.001	<10 <sup>-5</sup>		-0.001	0.151		-0.002	<10 <sup>-3</sup>	
	PEmin	0.102	<10 <sup>-5</sup>		0.094	<10 <sup>-5</sup>		0.099	<10 <sup>-5</sup>		0.186	<10 <sup>-5</sup>	
	PEmin <sup>2</sup>	-0.001	<10 <sup>-5</sup>		-0.001	<10 <sup>-5</sup>		-0.001	<10 <sup>-5</sup>		-0.002	<10 <sup>-5</sup>	
Wang et al. (2011b)	Intercept	1.175	0.044	0.498	3.775	<10 <sup>-5</sup>	0.783	0.756	0.609	0.572	0.195	0.803	0.691
	MTCQ	0.046	<10 <sup>-5</sup>		0.046	0.032		0.162	0.001		0.121	0.020	
	WD	-0.001	0.001		-0.002	<10 <sup>-5</sup>		0.001	0.186		-0.002	0.003	
	TSN	0.001	0.183		-0.001	0.083		0.002	0.142		0.001	0.353	
	RMAT	0.026	0.007		0.012	0.206		-0.073	0.025		0.077	<10 <sup>-5</sup>	
Xu et al. (2016)	Intercept	-0.939	<10 <sup>-5</sup>	0.710	-0.611	0.025	0.721	-0.961	0.004	0.799	-0.333	0.308	0.707
	Rainfall	0.003	<10 <sup>-5</sup>		0.003	<10 <sup>-5</sup>		0.003	<10 <sup>-5</sup>		0.001	0.044	
	PEmin	0.058	<10 <sup>-5</sup>		0.054	<10 <sup>-5</sup>		0.067	<10 <sup>-5</sup>		-0.004	0.706	
	Rainfall × PEmin	0.000	<10 <sup>-5</sup>		0.000	<10 <sup>-5</sup>		0.000	<10 <sup>-5</sup>		0.000	<10 <sup>-4</sup>	

Note. All four models show a high explanatory power for all continents. Please see Table 1 for definitions of abbreviations.

Genus *Quercus*Subgenus *Quercus*

**FIGURE 4** The model built based on European data predicted species richness in other continents with large errors. Root mean squared errors of the four multiple models were assessed when they are used to predict *Quercus* species richness at the global scale and in eastern Asia (EA), North America (NA) and Europe (EU) (colour bars). Each of the four models is calibrated separately using the global data, eastern Asian data, North American data and European data (x axis) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

only sampled 35 out of 150 species in this subgenus and incomplete sampling may probably lead to biases in our phylogenetic signal estimates. The mode of climatic niche evolution follows the BM model for the groups *Quercus* and *Lobatae* but the OU model for the subgenus *Cyclobalanopsis* (Table S6 in Supporting Information Appendix S2). The OU mode of climatic niche evolution suggests constrained evolution under directional selection that may lead to a loss of phylogenetic signal (Blomberg et al., 2003), which may account for the apparent lack of phylogenetic signal in climatic variables in the subgenus *Cyclobalanopsis*. Further studies on the evolutionary history


of *Cyclobalanopsis* could help to understand the origin of the species richness–climate relationship of this group.

### 5.3 | Effects of Quaternary climate change on oak species richness

Climate cooling and glaciation since the Quaternary has contributed to current species richness patterns likely through local extinctions and species range contraction. Species extinction during the Quaternary in eastern Asia was less severe than that in Europe and



**FIGURE 5** Latitudinal patterns of (a) the log-transformed species richness, (b) minimum monthly potential evapotranspiration (PET<sub>min</sub>) and (c) rainfall. (d) The changes in species richness along the gradients of both minimum monthly potential evapotranspiration and rainfall. The size of dots is scaled by species richness [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



North America, which has been regarded as one of the reasons for high species richness in eastern Asia (Latham et al., 1994; Qian & Ricklefs, 2000). In contrast, our study showed that the explanatory power of temperature anomalies since the LGM for species richness of the subgenus *Quercus* was higher in eastern Asia than in North America and Europe. One possible explanation of this pattern is that it was easier for species to track climate change through the south–north mountain ranges in North America during the Quaternary climatic oscillations compared to the east–west mountain ranges in eastern Asia and Europe. Pollen fossil studies showed that *Quercus* and other dominant tree species have expanded their ranges following climate change since the LGM with a lag of <1,500 years (Jackson et al., 2000). This probably also explains the stronger species richness–climate relationship for the subgenus *Quercus* in North America than eastern Asia and suggests that species richness in North America is more likely to be influenced by recent climate change.

#### 5.4 | Effects of habitat heterogeneity on oak species richness

The greater physiological heterogeneity in eastern Asia compared to other parts of the world is often regarded as the main driver of the region's higher species richness because of the role of this heterogeneity in providing refuges and cradles for species evolution and offering greater habitat variability, enabling species coexistence (Andrews & O'Brien, 2000; Kerr & Packer, 1997; Tews et al., 2004). However, our results revealed that habitat heterogeneity represented by elevation and climatic range explains the species richness variation only in Europe but not in eastern Asia or North America. The relatively high isolation of the mountains in southern Europe from southern landmasses limits the southward distribution of species and increases the species pool, possibly contributing to

high species richness in southern Europe and the significant effect of habitat heterogeneity on species richness in Europe (Svenning, Normand, & Skov, 2008).

## 6 | CONCLUSIONS

In contrast to the subtropical subgenus *Cyclobalanopsis*, our results support a consistent species richness–climate relationship for oaks in the temperate subgenus *Quercus* across the Northern Hemisphere, representing the effects of water, energy and their interaction. We did not find significant regional effects on species richness patterns of the temperate subgenus *Quercus*, which suggests that the species distribution and richness of temperate groups (e.g., subgenus *Quercus*) may have reached an equilibrium with the contemporary climate. The high level of niche conservatism and adaptation to cool and dry climate adopted by the temperate groups of *Quercus* may have played an important role in establishing its globally consistent richness–climate relationship across the Northern Hemisphere. Our results suggest that the distributions of groups with stronger niche conservatism are more strongly limited by climate, and hence are more likely to be affected by future climate change.

## ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (#31400467, #31522012, #31470564, #31621091, #31650110471), National Key Research Development Program of China (#2017YFA0605101), and Fundamental Research Funds for the central Universities (#YJ201721). D.D. and C.R. received additional support from the Danish National Research Foundation grant DNR96 to the Center for Macroecology, Evolution and Climate.

## AUTHOR CONTRIBUTIONS

Z.W., X.X. and C.R. designed the research; X.X. collected the data; X.X. and Z.W. performed the analyses; X.X. led the writing with contributions from all of the authors.

## DATA ACCESSIBILITY

All data necessary to reproduce the analyses presented in this study are included in the Supporting Information (Appendices S1 to S4).

## ORCID

Xiaoting Xu  <https://orcid.org/0000-0001-8126-614X>

Dimitar Dimitrov  <https://orcid.org/0000-0001-5830-5702>

## REFERENCES

- Andrews, P., & O'Brien, E. M. (2000). Climate, vegetation, and predictable gradients in mammal species richness in southern Africa. *Journal of Zoology*, 251, 205–231. <https://doi.org/10.1111/j.1469-7998.2000.tb00605.x>
- Axelrod, D. I. (1983). Biogeography of oaks in the Arcto-Tertiary province. *Annals of the Missouri Botanical Garden*, 70, 629–657. <https://doi.org/10.2307/2398982>
- Bininda-Emonds, O. R. P. (2004). The evolution of supertrees. *Trends in Ecology and Evolution*, 19, 315–322. <https://doi.org/10.1016/j.tree.2004.03.015>
- Bivand, R. S., & Wong, D. W. S. (2018). Comparing implementations of global and local indicators of spatial association. *TEST*, 27, 716–748. <https://doi.org/10.1007/s11749-018-0599-x>
- Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. *Evolution*, 57, 717–745. <https://doi.org/10.1111/j.0014-3820.2003.tb00285.x>
- Butler, M. A., & King, A. A. (2004). Phylogenetic comparative analysis: A modeling approach for adaptive evolution. *The American Naturalist*, 164, 683–695. <https://doi.org/10.1086/426002>
- Currie, D. J., & Francis, A. P. (2004). Taxon richness and climate in angiosperms: Is there a globally consistent relationship that precludes region effects? Reply. *The American Naturalist*, 163, 780–785.
- Currie, D. J., Mittelbach, G. G., Cornell, H. V., Field, R., Guegan, J. F., Hawkins, B. A., ... Turner, J. R. G. (2004). Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic richness. *Ecology Letters*, 7, 1121–1134. <https://doi.org/10.1111/j.1461-0248.2004.00671.x>
- Currie, D. J., & Paquin, V. (1987). Large-scale biogeographical patterns of species richness of trees. *Nature*, 329, 326–327. <https://doi.org/10.1038/329326a0>
- Deng, M., Jiang, X.-L., Hipp, A. L., Manos, P. S., & Hahn, M. (2018). Phylogeny and biogeography of East Asian evergreen oaks (*Quercus* section *Cyclobalanopsis*; Fagaceae): Insights into the Cenozoic history of evergreen broad-leaved forests in subtropical Asia. *Molecular Phylogenetics and Evolution*, 119, 170–181. <https://doi.org/10.1016/j.ympev.2017.11.003>
- Denk, T., & Grimm, G. W. (2009). Significance of pollen characteristics for Infrageneric classification and phylogeny in *Quercus* (Fagaceae). *International Journal of Plant Sciences*, 170, 926–940.
- Denk, T., & Grimm, G. W. (2010). The oaks of western Eurasia: Traditional classifications and evidence from two nuclear markers. *Taxon*, 59, 351–366. <https://doi.org/10.1002/tax.592002>
- Denk, T., Grimsson, F., & Zetter, R. (2010). Episodic migration of oaks to Iceland: Evidence for a North Atlantic “Land Bridge” in the latest Miocene. *American Journal of Botany*, 97, 276–287. <https://doi.org/10.3732/ajb.0900195>
- Denk, T., Grimsson, F., & Zetter, R. (2012). Fagaceae from the early Oligocene of Central Europe: Persisting New World and emerging Old World biogeographic links. *Review of Palaeobotany and Palynology*, 169, 7–20. <https://doi.org/10.1016/j.revpalbo.2011.09.010>
- Dutilleul, P., Clifford, P., Richardson, S., & Hemon, D. (1993). Modifying the t test for assessing the correlation between two spatial processes. *Biometrics*, 49, 305–314. <https://doi.org/10.2307/2532625>
- Felsenstein, J. (1973). Maximum-likelihood estimation of evolutionary trees from continuous characters. *American Journal of Human Genetics*, 25, 471–492.
- Field, R., O'Brien, E. M., & Whittaker, R. J. (2005). Global models for predicting woody plant richness from climate: Development and evaluation. *Ecology*, 86, 2263–2277. <https://doi.org/10.1890/04-1910>
- Fisher, J. B., Whittaker, R. J., & Malhi, Y. (2011). ET come home: Potential evapotranspiration in geographical ecology. *Global Ecology and Biogeography*, 20, 1–18. <https://doi.org/10.1111/j.1466-8238.2010.00578.x>
- Francis, A. P., & Currie, D. J. (2003). A globally consistent richness-climate relationship for angiosperms. *The American Naturalist*, 161, 523–536. <https://doi.org/10.1086/368223>
- Gaston, K. J. (2000). Global patterns in biodiversity. *Nature*, 405, 220–227. <https://doi.org/10.1038/35012228>
- Gordon, A. (1986). Consensus supertrees: The synthesis of rooted trees containing overlapping sets of labeled leaves. *Journal of Classification*, 3, 335–348. <https://doi.org/10.1007/BF01894195>
- Govaerts, R., Andrews, S., Coombes, A., Gilbert, M., Hunt, D., Nixon, K., & Thomas, M. (2011). *World checklist of Fagaceae*. Facilitated by the Royal Botanic Gardens, Kew. Retrieved from <http://apps.kew.org/wcsp/>
- Guo, Q., & Ricklefs, R. E. (2000). Species richness in plant genera disjunct between temperate eastern Asia and North America. *Botanical Journal of the Linnean Society*, 134, 401–423. <https://doi.org/10.1111/j.1095-8339.2000.tb00538.x>
- Harmon, L. J., Losos, J. B., Jonathan Davies, T., Gillespie, R. G., Gittleman, J. L., Bryan Jennings, W., ... Mooers, A. Ø. (2010). Early burst of body size and shape evolution are rare in comparative data. *Evolution*, 64, 2385–2396.
- Harmon, L. J., Weir, J. T., Brock, C. D., Glor, R. E., & Challenger, W. (2008). GEIGER: Investigating evolutionary radiations. *Bioinformatics*, 24, 129–131. <https://doi.org/10.1093/bioinformatics/btm538>
- Hawkins, B. A., Field, R., Cornell, H. V., Currie, D. J., Guegan, J. F., Kaufman, D. M., ... Turner, J. R. G. (2003). Energy, water, and broad-scale geographic patterns of species richness. *Ecology*, 84, 3105–3117. <https://doi.org/10.1890/03-8006>
- Hipp, A. L., Manos, P. S., González-Rodríguez, A., Hahn, M., Kaproth, M., McVay, J. D., ... Cavender-Bares, J. (2018). Sympatric parallel diversification of major oak clades in the Americas and the origins of Mexican species diversity. *New Phytologist*, 217, 439–452. <https://doi.org/10.1111/nph.14773>
- Hughes, C. E., & Atchison, G. W. (2015). The ubiquity of alpine plant radiations: From the Andes to the Hengduan Mountains. *New Phytologist*, 207, 275–282. <https://doi.org/10.1111/nph.13230>
- Jackson, S. T., Webb, R. S., Anderson, K. H., Overpeck, J. T., Webb, T., Williams, J. W., & Hansen, B. C. S. (2000). Vegetation and environment in Eastern North America during the Last Glacial Maximum. *Quaternary Science Reviews*, 19, 489–508. [https://doi.org/10.1016/S0277-3791\(99\)00093-1](https://doi.org/10.1016/S0277-3791(99)00093-1)

- Keck, F., Rimet, F., Bouchez, A., & Franc, A. (2016). phyloSignal: An R package to measure, test, and explore the phylogenetic signal. *Ecology and Evolution*, 6, 2774–2780. <https://doi.org/10.1002/ece3.2051>
- Kerr, J. T., & Packer, L. (1997). Habitat heterogeneity as a determinant of mammal species richness in high-energy regions. *Nature*, 385, 252–254. <https://doi.org/10.1038/385252a0>
- Kissling, W. D., & Carl, G. (2008). Spatial autocorrelation and the selection of simultaneous autoregressive models. *Global Ecology and Biogeography*, 17, 59–71.
- Latham, R. E., & Ricklefs, R. E. (1993). Global patterns of tree species richness in moist forests: Energy-diversity theory does not account for variation in species richness. *Oikos*, 67, 325–333. <https://doi.org/10.2307/3545479>
- Latham, R. E., Ricklefs, R. E., & Schluter, D. (1994). Continental comparisons of temperate-zone tree species diversity. In R. Ricklefs & D. Schluter (Eds.), *Species diversity in ecological communities: Historical and geographical perspectives* (chapter 26, pp. 294–314). Chicago, IL: University of Chicago Press.
- Lomolino, M. V., Riddle, B. R., Whittaker, R. J., & Brown, J. H. (2010). *Biogeography* (4th ed.). Sunderland, MA: Sinauer Associates.
- Manchester, S. R. (1994). Fruits and seeds of the middle eocene nut beds flora, clarno formation, oregon. Paleontological Research Institution.
- Manos, P. S., & Stanford, A. M. (2001). The historical biogeography of Fagaceae: Tracking the tertiary history of temperate and subtropical forests of the Northern Hemisphere. *International Journal of Plant Sciences*, 162, S77–S93. <https://doi.org/10.1086/323280>
- Menitsky, Y. L. (2005). *Oaks of Asia*. New Hampshire: Science Publishers, Inc.
- Molnar, P., Boos, W. R., & Battisti, D. S. (2010). Orographic controls on climate and paleoclimate of Asia: Thermal and mechanical roles for the Tibetan Plateau. *Annual Review of Earth and Planetary Sciences*, 38, 77–102. <https://doi.org/10.1146/annurev-earth-040809-152456>
- Munkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schifffers, K., & Thuiller, W. (2012). How to measure and test phylogenetic signal. *Methods in Ecology and Evolution*, 3, 743–756. <https://doi.org/10.1111/j.2041-210X.2012.00196.x>
- Nixon, K. C. (1997). Fagaceae. In *Flora of North America Editorial Committee (Eds.), 1993+ Flora of North America North of Mexico. 20+ vols (Vol. 4, pp. 314–400)*. New York and Oxford.
- O'Brien, E. M. (1998). Water-energy dynamics, climate, and prediction of woody plant species richness: An interim general model. *Journal of Biogeography*, 25, 379–398. <https://doi.org/10.1046/j.1365-2699.1998.252166.x>
- O'Brien, E. M. (2006). Biological relativity to water-energy dynamics. *Journal of Biogeography*, 33, 1868–1888. <https://doi.org/10.1111/j.1365-2699.2006.01534.x>
- O'Brien, E. M., Field, R., & Whittaker, R. J. (2000). Climatic gradients in woody plant (tree and shrub) diversity: Water-energy dynamics, residual variation, and topography. *Oikos*, 89, 588–600. <https://doi.org/10.1034/j.1600-0706.2000.890319.x>
- Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics*, 20, 289–290. <https://doi.org/10.1093/bioinformatics/btg412>
- Prothero, D. R. (1994). The Late Eocene-Oligocene extinctions. *Annual Review of Earth and Planetary Sciences*, 22, 145–165. <https://doi.org/10.1146/annurev.ea.22.050194.001045>
- Qian, H. (2002). Floristic relationships between Eastern Asia and North America: Test of Gray's hypothesis. *The American Naturalist*, 160, 317–332.
- Qian, H., Jin, Y., & Ricklefs, R. E. (2017). Phylogenetic diversity anomaly in angiosperms between eastern Asia and eastern North America. *Proceedings of the National Academy of Sciences USA*, 43, 11452–11457. <https://doi.org/10.1073/pnas.1703985114>
- Qian, H., & Ricklefs, R. E. (2000). Large-scale processes and the Asian bias in species diversity of temperate plants. *Nature*, 407, 180–182. <https://doi.org/10.1038/35025052>
- Qian, H., & Ricklefs, R. E. (2016). Out of the tropical lowlands: Latitude versus elevation. *Trends in Ecology and Evolution*, 31, 738–741. <https://doi.org/10.1016/j.tree.2016.07.012>
- R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>.
- Rahbek, C., & Graves, G. R. (2001). Multiscale assessment of patterns of avian species richness. *Proceedings of the National Academy of Sciences USA*, 98, 4534–4539. <https://doi.org/10.1073/pnas.071034898>
- Rangel, S. R. (2006). Revisión taxonómica del complejo Acutifoliae de Quercus (Fagaceae) con énfasis en su representación en México. *Acta Botanica Mexicana*, 76, 1–45. <https://doi.org/10.21829/abm76.2006.1016>
- Rangel, S., Carlos, E., Zenteno, R., De Lourdes, M., & Enríquez, A. (2002). El género quercus (Fagaceae) en el estado de México. *Annals of the Missouri Botanical Garden*, 89(4), 551–593.
- Ricklefs, R. E. (1987). Community diversity—Relative roles of local and regional processes. *Science*, 235, 167–171. <https://doi.org/10.1126/science.235.4785.167>
- Ricklefs, R. E. (2004). A comprehensive framework for global patterns in biodiversity. *Ecology Letters*, 7, 1–15. <https://doi.org/10.1046/j.1461-0248.2003.00554.x>
- Ricklefs, R. E., & He, F. L. (2016). Region effects influence local tree species diversity. *Proceedings of the National Academy of Sciences USA*, 113, 674–679. <https://doi.org/10.1073/pnas.1523683113>
- Ricklefs, R. E., Latham, R. E., & Qian, H. (1999). Global patterns of tree species richness in moist forests: Distinguishing ecological influences and historical contingency. *Oikos*, 86, 369–373. <https://doi.org/10.2307/3546454>
- Ricklefs, R. E., Qian, H., & White, P. S. (2004). The region effect on mesoscale plant species richness between eastern Asia and eastern North America. *Ecography*, 27, 129–136. <https://doi.org/10.1111/j.0906-7590.2004.03789.x>
- Ricklefs, R. E., & Renner, S. S. (1994). Species richness within families of flowering plants. *Evolution*, 48, 1619–1636. <https://doi.org/10.1111/j.1558-5646.1994.tb02200.x>
- Svenning, J.-C., Normand, S., & Skov, F. (2008). Postglacial dispersal limitation of widespread forest plant species in nemoral Europe. *Ecography*, 31, 316–326. <https://doi.org/10.1111/j.0906-7590.2008.05206.x>
- Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M. C., Schwager, M., & Jeltsch, F. (2004). Animal species diversity driven by habitat heterogeneity/diversity: The importance of key-stone structures. *Journal of Biogeography*, 31, 79–92. <https://doi.org/10.1046/j.0305-0270.2003.00994.x>
- Trabucco, A., Zomer, R. J., Bossio, D. A., van Straaten, O., & Verchot, L. V. (2008). Climate change mitigation through afforestation/reforestation: A global analysis of hydrologic impacts with four case studies. *Agriculture, Ecosystems and Environment*, 126, 81–97. <https://doi.org/10.1016/j.agee.2008.01.015>
- Wagner, C. E., Keller, I., Wittwer, S., Selz, O. M., Mwaiko, S., Greuter, L., ... Seehausen, O. (2013). Genome-wide RAD sequence data provide unprecedented resolution of species boundaries and relationships in the Lake Victoria cichlid adaptive radiation. *Molecular Ecology*, 22, 787–798. <https://doi.org/10.1111/mec.12023>
- Wang, Z., Fang, J., Tang, Z., & Lin, X. (2011a). Patterns, determinants and models of woody plant diversity in China. *Proceedings of the Royal Society B: Biological Sciences*, 278, 2122–2132.
- Wang, Z., Fang, J., Tang, Z., & Lin, X. (2011b). Relative role of contemporary environment versus history in shaping diversity patterns of China's woody plants. *Ecography*, 34, 1–10.
- Wen, J., Zhang, J. Q., Nie, Z. L., Zhong, Y., & Sun, H. (2014). Evolutionary diversifications of plants on the Qinghai-Tibetan Plateau. *Frontiers in Genetics*, 5, 4. <https://doi.org/10.3389/fgene.2014.00004>
- Whittaker, R. J., Noguez-Bravo, D., & Araujo, M. B. (2007). Geographical gradients of species richness: A test of the water-energy conjecture of Hawkins et al. (2003) using European data for five taxa. *Global Ecology and Biogeography*, 16, 76–89.

- Wiens, J. J., Ackerly, D. D., Allen, A. P., Anacker, B. L., Buckley, L. B., Cornell, H. V., ... Stephens, P. R. (2010). Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters*, 13, 1310–1324. <https://doi.org/10.1111/j.1461-0248.2010.01515.x>
- Wiens, J. J., & Donoghue, M. J. (2004). Historical biogeography, ecology and species richness. *Trends in Ecology and Evolution*, 19, 639–644. <https://doi.org/10.1016/j.tree.2004.09.011>
- Wu, Z., Raven, P. H., & Hong, D. (1999). *Flora of China, vol. 4 (Cycadaceae through Fagaceae)*. Beijing, China: Science Press; St. Louis, MO: Missouri Botanic Garden.
- Xiang, Q.-Y., Zhang, W. H., Ricklefs, R. E., Qian, H., Chen, Z. D., Wen, J., & Li, J. H. (2004). Regional differences in rates of plant speciation and molecular evolution: A comparison between Asia and Eastern North America. *Evolution*, 58, 2175–2184.
- Xing, Y., & Ree, R. H. (2017). Uplift-driven diversification in the Hengduan Mountains, a temperate biodiversity hotspot. *Proceedings of the National Academy of Sciences*, 114, E3444–E3451.
- Xu, X., Wang, Z., Rahbek, C., Lessard, J.-P., & Fang, J. (2013). Evolutionary history influences the effects of water–energy dynamics on oak diversity in Asia. *Journal of Biogeography*, 40, 2146–2155. <https://doi.org/10.1111/jbi.12149>
- Xu, X., Wang, Z., Rahbek, C., Sanders, N. J., & Fang, J. (2016). Geographical variation in the importance of water and energy for oak diversity. *Journal of Biogeography*, 43, 279–288. <https://doi.org/10.1111/jbi.12620>

## BIOSKETCHES

Dr XIAOTING XU is an associate professor at Sichuan University, China. She is interested in understanding the origin, maintenance and conservation of biological diversity.

ZHIHENG WANG has interests in macroecology and biogeography, particularly the ecological and evolutionary mechanisms underlying large-scale species richness patterns and the impacts of climate change on species distributions.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Xu X, Dimitrov D, Shrestha N, Rahbek C, Wang Z. A consistent species richness–climate relationship for oaks across the Northern Hemisphere. *Global Ecol Biogeogr.* 2019;28:1051–1066. <https://doi.org/10.1111/geb.12913>