



REVIEW

Standardized monitoring of permafrost thaw: a user-friendly, multiparameter protocol¹

Julia Boike, Sarah Chadburn, Julia Martin, Simon Zwieback, Inge H.J. Althuisen, Norbert Anselm, Lei Cai, Stéphanie Coulombe, Hanna Lee, Anna K. Liljedahl, Martin Schneebeli, Ylva Sjöberg, Noah Smith, Sharon L. Smith, Dmitry A. Streletskiy, Simone M. Stuenzi, Sebastian Westermann, and Evan J. Wilcox

Abstract: Climate change is destabilizing permafrost landscapes, affecting infrastructure, ecosystems, and human livelihoods. The rate of permafrost thaw is controlled by surface and subsurface properties and processes, all of which are potentially linked with each other. However, no standardized protocol exists for measuring permafrost thaw and related processes and properties in a linked manner. The permafrost thaw action group of the Terrestrial Multidisciplinary distributed Observatories for the Study of the Arctic Connections (T-MOSAIC) project has developed a protocol, for use by non-specialist scientists and technicians, citizen scientists, and indigenous groups, to collect standardized metadata and data on permafrost thaw. The protocol introduced here addresses the need to jointly measure permafrost thaw and the associated surface and subsurface environmental conditions. The parameters measured along transects include: snow depth, thaw depth, vegetation height, soil texture, and water level. The metadata collection includes data on

Received 2 February 2021. Accepted 21 June 2021.

J. Boike* and **S.M. Stuenzi.** Alfred Wegener Institute Helmholtz Center for Polar and Marine Research (AWI), Telegrafenberg, 14473 Potsdam, Germany; Humboldt University Berlin, Geography Department, Unter den Linden 6, 10099 Berlin, Germany.

S. Chadburn* and **N. Smith.** College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter EX4 4QF, UK.

J. Martin.* Alfred Wegener Institute Helmholtz Center for Polar and Marine Research (AWI), Telegrafenberg, 14473 Potsdam, Germany; WSL Institute for Snow and Avalanche Research SLF, Flüelastrasse 11, 7260 Davos Dorf, Switzerland.

S. Zwieback.* Geophysical Institute, University of Alaska Fairbanks, Fairbanks 99775, Alaska, USA.

I.H.J. Althuisen and H. Lee. NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Nygårdsgaten 112, 5008 Bergen, Norway.

N. Anselm. Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Mailbox 12 01 61, 27515 Bremerhaven, Germany.

L. Cai. Department of Atmospheric Sciences, Yunnan University, North Cuihu Road 2, Kunming, 650000, China.

S. Coulombe. Canadian High Arctic Research Station, Polar Knowledge Canada, Cambridge Bay, NU X0B 0C0, Canada.

A.K. Liljedahl. Woodwell Climate Research Center, 149 Woods Hole Road, Falmouth, MA, 02540-1644, USA.

M. Schneebeli. WSL Institute for Snow and Avalanche Research SLF, Flüelastrasse 11, 7260 Davos Dorf, Switzerland.

Y. Sjöberg. Department of Geosciences and Natural Resource Management, Centre for Permafrost (CENPERM), University of Copenhagen, Øster Voldgade 10, 1350 Copenhagen, Denmark.

S.L. Smith. Geological Survey of Canada, Natural Resources Canada, Ottawa, ON K1A 0E8, Canada.

D.A. Streletskiy. Department of Geography, The George Washington University, 2036 H St NW, Washington DC, 20052, USA.

S. Westermann. Department of Geosciences, University of Oslo, Sem Sælands vei 1, 0371 Oslo, Oslo, Norway.

E.J. Wilcox. Cold Regions Research Centre, Wilfrid Laurier University, 75 University Avenue West, Waterloo, N2L 3C5, Canada.

Corresponding author: Julia Boike (e-mail: julia.boike@awi.de).

*These authors are the co-lead authors.

¹This article is part of a Virtual Special Issue entitled: Terrestrial Geosystems, Ecosystems, and Human Systems in the Fast-Changing Arctic.

© 2021 The Author(s). This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

timing of data collection, geographical coordinates, land surface characteristics (vegetation, ground surface, water conditions), as well as photographs. Our hope is that this openly available dataset will also be highly valuable for validation and parameterization of numerical and conceptual models, and thus to the broad community represented by the T-MOSAIc project.

Key words: snow depth, vegetation height, soil characteristics, active layer thaw depth, permafrost monitoring protocol.

Résumé : Le changement climatique déstabilise les paysages de pergélisol, affectant les infrastructures, les écosystèmes et les moyens de subsistance des populations. Le taux de dégel du pergélisol est contrôlé par les propriétés et les processus en surface et en subsurface, qui sont tous potentiellement liés les uns aux autres. Pourtant, il n'existe pas de protocole normalisé pour mesurer le dégel du pergélisol et les processus et propriétés connexes de manière liée. Le groupe d'action sur le dégel du pergélisol *Terrestrial Multidisciplinary distributed Observatories for the Study of the Arctic Connections (T-MOSAIc)* a élaboré un protocole destiné à être utilisé par des scientifiques et des techniciens non spécialisés, des scientifiques citoyens et des groupes autochtones, pour collecter des métadonnées et des données normalisées sur le dégel du pergélisol. Le protocole présenté ici répond à la nécessité de mesurer conjointement le dégel du pergélisol et les conditions environnementales associées en surface et en subsurface. Les paramètres mesurés le long des transects sont : l'épaisseur de la neige, l'épaisseur du dégel, la hauteur de la végétation, la texture du sol et le niveau de l'eau. La collection de métadonnées comprend des données sur le moment de la collecte des données, les coordonnées géographiques, les caractéristiques de la surface émergée (végétation, surface du sol, conditions de l'eau), ainsi que des photographies. Les auteurs espèrent que ce jeu de données librement accessible sera également très utile pour la validation et le paramétrage de modèles numériques et conceptuels, et donc pour la vaste communauté représentée par le projet T-MOSAIc. [Traduit par la Rédaction]

Mots-clés : hauteur de neige, hauteur de végétation, caractéristiques du sol, profondeur de dégel de la couche active, protocole de surveillance du pergélisol.

Background and general introduction

Northern landscapes and infrastructure are affected by the thaw of permafrost, especially in regions of ice-rich permafrost, because thawing can lead to surface subsidence and slope instability. Permafrost thaw has profound implications for Arctic ecosystems and their inhabitants, through changes to surface drainage and water resources (Osterkamp et al. 2009; Kokelj and Jorgenson 2013), vegetation and wildlife habitats (Sturm et al. 2001b; Jorgenson et al. 2010), and through the positive feedback to global warming via the emission of greenhouse gases (Burke et al. 2017; Burke et al. 2017; Hugelius et al. 2020; Turetsky et al. 2020).

There is an urgent need for standardized monitoring of permafrost conditions. The impacts of permafrost thaw on ecosystems are expected to increase with climate warming, changes in precipitation, and increasing surface disturbance (Kokelj and Jorgenson 2013; Rasmussen et al. 2018). For 2020, the Arctic Report Card highlights the highest recorded surface air temperatures, record lows of June snow cover, opposing trends of tundra greenness, and extreme wildfires (Arctic Program 2020). Permafrost temperature and active layer thickness are increasing, but there is considerable spatial variability in the magnitude of the change, owing to local variations in snow, vegetation, and soil characteristics (Romanovsky et al. 2020). These local variabilities are critical for the evaluation of permafrost thaw. Not only do the rate and nature of permafrost thaw depend on factors such as snow depth, the thickness of the organic layer, and vegetation height, but permafrost thaw will in turn influence these variables (Vincent et al. 2017). For example, increases over time in the density and height of shrubs have been reported from tundra regions across

Table 1. Summary of existing protocols for the parameters presented in this paper.

Spheres	Existing protocols, organization
Snow	Essential Climate Variables products and requirements for snow, The Global Climate Observing System (GCOS) (The Global Climate Observing System 2016a) Estimating the snow water equivalent from snow depth data, International Commission for Snow and Ice Hydrology (ICSH) (Jonas and Marks 2016) The international classification for seasonal snow on the ground, International Association of Cryospheric Sciences (IACS) (Fierz et al. 2009) European Snow Booklet, WSL Institute for Snow and Avalanche Research SLF (Haberkmorn 2019) Chapter 5: Snow and Ice, International Tundra Experiment (ITEX) Manual, Danish Polar Center (Molau 1996)
Permafrost	Global Terrestrial Network for Permafrost, International Permafrost Association (IPA) (Streletskiy et al. 2017b) Methods for measuring active-layer thickness; A handbook on periglacial field methods; IPA; Circumpolar Active Layer Monitoring Network (CALM) (Nelson and Hinkel 2004) Essential climate variables (ECVs) products and requirements for permafrost, GCOS (The Global Climate Observing System 2016b) Active layer monitoring standard protocol, Arctic Development and Adaptation to Permafrost in Transition (ADAPT) (Arctic Development and Adaptation to Permafrost in Transition) Chapter 6: Active layer protocol, (ITEX) manual (Nelson et al. 1996) Assessment of the status of the development of the standards for the Terrestrial Essential Climate Variables, Permafrost (Smith and Brown 2009)
Vegetation	Chapter 11: Community baseline measurements, ITEX manual (Molau and Edlund 1996) Vegetation standard description protocol, ADAPT (Grogan et al. n.d.) New handbook for standardised measurement of plant functional traits worldwide (Pérez-Harguindeguy et al. 2013)
Water	Guide to Hydrological Parameters – Volume 1, World Meteorological Organization (World Meteorological Organization 2008) Soil moisture content, CALM (Circumpolar Active Layer Monitoring Network)
Soil	Sampling protocols for permafrost-affected soils (Ping et al. 2013) Soil survey fields and laboratory methods, U.S. Department of Agriculture, Natural Resources Conservation Service (Soil Survey Staff 2014) Active layer sampling standard protocol for C/H/N determination, ADAPT (Arctic Development and Adaptation to Permafrost in Transition ^a) Planning and making a soil survey, Food and Agriculture Organization of the United Nations (Food and Agriculture Organization of the United Nations) Terrestrial instrument system (TIS) soil pit sampling protocol, The National Ecological Observatory Network (NEON) (The National Ecological Observatory Network 2021) The United Nations Terminology Database, United Nations (United Nations 2012)

Note: The parameters are grouped into five spheres: snow, permafrost, vegetation, water, and soil. Citable references are given in the table; some projects (example ADAPT, CALM) provide protocols online for which we provide links in the reference section of this paper.

^a<https://www.cen.ulaval.ca/adapt/protocols/adapt.php>.

the Arctic, and locally, shrub expansion may also be driven by permafrost degradation ([Sturm et al. 2001b](#)). Shrub growth can in turn reduce ([Blok et al. 2010](#)) or promote ([Wilcox et al. 2019](#)) permafrost thaw, depending on how shrub height affects snow accumulation and snow melt. The hydrological conditions in ice-rich permafrost lowlands determine the thawing of permafrost; inundated and wetter areas favour degradation, whereas drainage and drier soil conditions favour stabilization ([Nitzbon et al. 2020](#)).

No common protocol exists that simultaneously considers both permafrost thaw and the key environmental variables that affect permafrost thaw. A number of protocols have already been created by specialized research communities ([Table 1](#)), but each is dedicated to only a small subset of parameters. Collocated and consistent measurements of multiple variables are needed to explain changes in permafrost conditions, and therefore to upscale or to make future projections of future permafrost thaw. In addition, particular parameters are required as inputs for numerical and conceptual models (including Earth system

models and specialized permafrost models, such as CryoGrid; Nitzbon et al. 2020). The focus of our study was to design such a multiparameter protocol.

Here we developed simple protocols and an associated mobile app that will enable a wide range of non-expert users to make high-quality, standardized, and accessible measurements. Our protocols address the need for consistent collection and integration of data from across the permafrost region to: (i) better monitor and understand permafrost thaw; (ii) establish a baseline against which future change can be measured; and (iii) support the integration of field measurements within pan-Arctic geospatial datasets developed through remote sensing analyses or modelling. The app guides the user through the observation process, ensures that the observations are consistent and well documented, and transfers the observations to an accessible database.

We developed the protocol in the Terrestrial Multidisciplinary distributed Observatories for the Study of the Arctic Connections (T-MOSAiC) action group on permafrost thaw. T-MOSAiC is an International Arctic Science Committee (IASC) pan-Arctic, land-based programme that extends the activities of the sea-based programme Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAiC; <https://mosaic-expedition.org/>). Originally T-MOSAiC was planned to run concomitantly with MOSAiC to achieve simultaneous measurements of biogenic, hydrological, and atmospheric fluxes by extending the work to the lands surrounding the Arctic Ocean. Because the COVID pandemic limited travel to field sites, T-MOSAiC was extended to the end of 2021. Intense monitoring is proposed for 2021 to kick-start a longer term observational program to monitor the progression of thaw in permafrost and other associated environmental changes.

In this paper, we detail the rationale behind the protocol and choice of measurements; the detailed protocol is available in [Appendix A](#).

Protocol overview — choice of parameters and scale issue


Protocols for everyone

The protocol's primary target group is not the permafrost experts, but persons with limited prior field experience. The users comprise professionals and students from a wide range of backgrounds, including ecology, hydrology, and geology. In-depth expertise in permafrost ecosystems is not required. Citizen scientists form the protocol's secondary target audience, ideally under guidance from an experienced user. For instance, a high-school class could continually monitor the permafrost conditions with support from a biology teacher.

The protocol is geared towards nonspecialists in three ways. First, no specialized knowledge or skills are needed. The measurements are simple, and an app has been developed to guide the user through the measurement process. In addition, videos are provided to illustrate key steps. The app also takes care of data handling, ensuring data quality and usability by enforcing the compilation of required metadata and homogenizing data transmission, and storage. Second, no specialized equipment is needed. The protocol only requires simple tools, namely a ruler, camera, tape measure, shovel, and a steel rod. Finally, the protocol has been streamlined so as not to take up too much of the nonspecialist's time.

[Appendix A](#) provides further details of the app for data collection, as well as instructional videos. One was recorded at a permafrost site in northern Norway in autumn 2020 by fine-art students. Another one was recorded at the permafrost long-term observatory site Bayelva on Svalbard, Norway, in spring 2021 by the permafrost thaw action group and the Alfred Wegener Institute Helmholtz Center for Polar and Marine Research (AWI).

Fig. 1. Spheres with the associated parameters, measurement modes, and observation timings along one transect over one seasonal cycle.



Parameters

We grouped the parameters for which we provide protocols into 5 spheres as follows:


1. Snow: snow depth;
2. Permafrost: thaw depth;
3. Vegetation: vegetation height;
4. Water: water level;
5. Soil: organic layer depth, soil texture, ground ice.

We chose the specific measurement parameters (Fig. 1) to cover the major controls of permafrost thaw with simple measurements that are accessible to non-experts, and in doing so we inevitably cannot include some commonly used parameters, such as soil temperature, owing to their need for specialized equipment.

Figure 1 gives an overview of the spheres we incorporate, the measurements described in this protocol, and their seasonality. Measurements start during the winter on snow, and are continued at the same transect points through the seasons of snowmelt, vegetation growth, deepening of the thawed layer, and development of a water level in summer. Measurements of soil properties, such as organic layer thickness and soil texture, are only done once along the transect — ideally during the later part of the season when the thawed layer has reached its maximum.

Our five parameters in these spheres can vary dramatically across the landscape, for example, snow depth on palsas is much shallower than on an adjacent mire (Martin et al. 2019). In addition, all these spheres interact with each other, and the landscape variability

Fig. 2. Example of landscape variability covering palsa mire, forest, and upland tundra (Iškoras; Finnmark, northern Norway). Typically, one 10 m long transect cannot cover all the characteristic features as shown in this figure. If timing and capacities allow, several transects can be established. If there is already an established transect at this site it can be used.



is sometimes driven by dynamic feedbacks between these parameters, which can amplify small variations into major sources of heterogeneity. For example, a small variation in surface elevation can lead to a positive feedback in which snow and water accumulate in the depression, warming the ground and leading to thaw and potential ground subsidence (if the permafrost is ice-rich), resulting in further accumulation of snow and water, and increasing permafrost thaw at this location (Kokelj and Jorgenson 2013; Nitzbon et al. 2020). Some features vary at the metre-scale, including microtopography such as hummocks, and vegetation. Others will vary on the scale of hundreds of metres, such as differences between valley bottoms and hillslopes. This protocol accounts for these issues of parameter interconnectivity and variability by using transects, with measurements of multiple parameters from different spheres conducted on the same transect.

Where to measure?

The protocol design aims to ensure that measurements capture the variability within a landscape. Because the overarching goal is to understand permafrost thaw on a pan-Arctic scale, we must consider the issues in scaling between a measurement at a single point to regional models/satellite data pixels (10s to a few 100s of metres to kilometres) and global models (10s to 100s of kilometres).

To ensure representation of variability within a landscape, and taking into account the target audience and time constraints in the field, we chose the scale of the measurements as a 10–30 m long transect to allow “typical microtopographic features” to be resolved by

Arctic Science Downloaded from cdnsciencepub.com by UNIVERSITY OF OSLO on 02/28/22
For personal use only.

sampling every 1 m. This means that the minimum effort (one 10 m long transect) can resolve a key aspect of variability and requires very little investment of time. Examples of typical microtopography captured by the sampling strategy include tundra polygons and peat plateaus/palsas, which are typical landforms in permafrost areas.

Time permitting, larger-scale variability will be captured with further transects in the local area, taking account of the landscape features that are present. For example, at the Iškoras site in northern Norway (Fig. 2), separate transects would ideally cover the palsa mire, the forest, and the nearby upland tundra. Furthermore, larger-scale topographic features, such as the slopes and the bottom of a valley, could be captured through multiple transects. In the protocol we urge the users to consider the landscape variability in and around their site, and to select “representative” locations for their transect (see the protocol described in Appendix A).

Details of the spheres’ parameters

The five measurement spheres are described below. Here we give details on the scientific importance of each sphere and its interactions with permafrost thaw, as well as the rationale behind the choice of parameter to measure and the chosen measurement technique.

Snow

Background

Snow cover exerts a fundamental control on the thermal and hydrological regime of permafrost. It acts as an insulator thanks to its low thermal conductivity, reducing heat loss in winter (Zhang 2005; Grünberg et al. 2020). The type of vegetation cover can significantly influence the insulating power of snow because plants affect the distribution of snow and its depth (Domine et al. 2018). In spring, snow strongly reflects the solar radiation (i.e., a high albedo) (Stiegler et al. 2016). The duration and extent of the snow cover in spring regulate the soil temperature and meltwater supply (Boike et al. 2003). Snow masses in Arctic regions are highly diverse and determined by regional conditions. Trend analyses point out an increase of snow masses in Siberian regions where others are likely to decrease (Callaghan et al. 2011; Pulliainen et al. 2020).

We focus here on snow depth, as the thermal resistance of the snowpack is in the first order a function of snow depth (Zhang et al. 1996). Crumley et al. (2020) show the usefulness of snow depth measurements for a citizen science approach for a different application. Snow depth is spatially variable due to land-cover characteristics (topography, vegetation) and wind-induced redistribution. For example, the snow cover on plains can experience drift (Sturm et al. 2001a; Parr et al. 2020), whereas local depressions, or an abundance of shrubs, trap snow (Wilcox et al. 2019). Critical observation times are the onset of snow accumulation at the beginning of the winter season, absolute maximum during winter, and maximum height just before spring melt. We recommend regular observations with a frequency of at least once per month, ideally once per week.

Measurement

Snow depth is the full height of a snowpack measured perpendicular to the underlying ground (Haberhorn 2019). Snow depth captures evolution of the snow cover over time with minimal effort but maximum information.

It is measured mechanically using either a simple ruler to record the depth or, if available, a snow rod with the measuring units already on the probe. Those tools are easy to obtain and user friendly. Snow depth measurements can be difficult if the snowpack is very hard or if the soil below the snow is very soft. In the first case, the probe may not reach

the ground (e.g., if there is a hard refrozen crust within the snowpack or in the presence of a basal ice layer). In the second case, the probe may penetrate the ground (e.g., unfrozen peat, deep grass, or moss hummock). The vegetation (e.g., bushes) within the snowpack can also influence the measurement.

Permafrost

Background

Thaw depth is the only variable for characterizing permafrost conditions that is included in the T-MOSAIc protocol. It is defined as the distance between the ground surface and the frost table (Brown et al. 2000). Thaw depth increases over the summer period, as the thaw front penetrates deeper into the ground. The most critical time for measuring thaw depth is at the end of the thaw season, when thaw depth is at or near its annual maximum (Brown et al. 2000). This timing typically ranges from mid-August to mid-September in the arctic and subarctic regions (Brown et al. 2000).

Thaw depth is an important variable for characterizing changing permafrost conditions because increasing air temperatures and ground warming often cause the maximum thaw depth to increase via thawing at the top of the permafrost (Brown et al. 2000). However, two additional factors have to be considered when using maximum thaw depth as an indicator of permafrost response to climate conditions. Firstly, the maximum annual thaw depth varies from year to year in response to interrelated variables such as soil moisture, vegetation, and snow (e.g., Walker et al. 2003; Shiklomanov et al. 2010; Grünberg et al. 2020). Secondly, the thawing of ice-rich permafrost primarily induces subsidence rather than increases in thaw depth (Osterkamp et al. 2009; O'Neill et al. 2019). Hence, a comprehensive quantification of permafrost thaw necessitates observations of subsidence (Streletskiy et al. 2017a). While direct observations of subsidence are not included in the protocol because of the lack of simple methods for measuring it, the measurements of vegetation and inundation (wetness) can indicate subsidence induced by the thaw of ice-rich permafrost (Kokelj and Jorgenson 2013).

Measurement

Multiple methods exist for measuring thaw depth in the field (Smith and Brown 2009). Mechanical probing is arguably the most popular method because it does not require sophisticated equipment (Brown et al. 2000), and for this same reason it is the method adopted for the T-MOSAIc protocol.

Thaw depth is measured by inserting a pointed metal rod (usually 1.0–1.5 m in length) into the soil down to the point of resistance against the frost table at each point along the transect. The depth that the rod has been inserted into the ground can then be determined using a measuring tape, or from graduated marks on the rod itself.

The measurements need to account for the substantial small-scale spatial variability in thaw depth. To ensure unbiased sampling and to facilitate comparisons over time, the measurement should be made in immediate proximity to the marked transect point. If standing water should make it too difficult to measure at the point, the measurement should be marked as “Water”.

Mechanical probing works best in organic and gravel-poor mineral soils that are ice bonded when frozen (Brown et al. 2000). The app guides the user through challenges that may arise for substrates that are less amenable to probing. The most commonly encountered limitations are:

- probing may be impossible in bedrock or gravel;
- it can be difficult to distinguish between subsurface stones and frozen substrate, for instance in soils that contain gravel;
- in locations of deep thaw, the thaw depth may exceed the length of the rod;
- the unusual mechanical properties of saline marine sediments or plastically frozen clays present a challenge to frost probing.

Vegetation

Background

Vegetation is an important component in influencing the surface energy balance and the thermal and hydrological regime of permafrost. At the same time it can also react to changes in the environment (Myers-Smith et al. 2011). Different vegetation types can have contrasting effects on permafrost ecosystems. Forests are usually considered to efficiently insulate the underlying permafrost (Chang et al. 2015) by altering the thermal regime, by intercepting snow, and promoting the accumulation of an organic surface layer (Bonan and Shugart 1989; Stuenzi et al. 2021). Low stature tundra vegetation can similarly alter thermal and hydrological conditions through differences in albedo between vegetation types (Juszak et al. 2016; Aartsma et al. 2020), as well as the effect of vegetation height on snow conditions, including snow depth, snowmelt, and the physical properties of snow (Domine et al. 2018; Wilcox et al. 2019). From a permafrost-thaw perspective, we consider the presence and the height of vegetation as the most important parameters for including vegetation in permafrost modelling. Vegetation height is commonly measured from the soil surface to the highest point of the vegetation. As multiple measurements are made within each quadrat, this will then provide representative average vegetation heights along the transect (similarly with height measurements of multiple trees).

Measurement

The measurement of vegetation height can provide a good estimate of the type of vegetation regime present, and requires little knowledge about actual plant species or plant functional types. Height measurements should be carried out in 1 m × 1 m quadrats (Molau and Edlund 1996) at each point along a 10–30 m transect. This transect should be established before taking any measurements at the site. Optionally, if the site is located in forest, a minimum of 10 individual trees in a 15 m × 15 m plot should also be measured (Pérez-Harguindeguy et al. 2013; Kruse et al. 2019). Most measurements therefore require a ruler or tape measure only, but in tall forests it might be necessary to give training in height estimation beforehand.

Water

Background

Permafrost has a primary influence on the movement of water through a landscape, and water, in turn, impacts the ground thermal regime and the rate of permafrost thaw (Riseborough et al. 2008; Woo 2012). The liquid water and ice content of a soil exerts a fundamental control on its thermal diffusivity, and thereby the transport of heat between the active layer and permafrost (Edlefsen and Anderson 1943; Kurylyk and Watanabe 2013). Furthermore, the water content influences the thawing and freezing rates of the ground because of the latent heat associated with melting or freezing (Outcalt et al. 1990). In addition to influencing the rate of thaw, surface and groundwater are also indicators of thaw of ice-rich permafrost, which can lead to impoundment in depressions (Jorgenson et al. 2010). Observations of wetness are thus critical for predicting and monitoring permafrost thaw (Jorgenson et al. 2010; Chadburn et al. 2015; Rasmussen et al. 2018).

Measurement

From a permafrost thaw perspective, we consider the spatial and temporal distribution of soil wetness indicated by the height of the water table the most important hydrological variable to record. Water table observations are most easily done in combination with measurement of thaw depth or soil pit, as it can be carried out with the same equipment and along the same transect. Acquiring observations of both wetness and thaw depth at the same locations and times helps in later interpreting the relationship between water level and soil thaw. Following our protocol, the height of the water table relative to the ground surface level is noted in the hole (using the frost probe, shovel, or your hands) as: “above the ground surface”, “within 10 cm below the ground surface”, or “more than 10 cm below the ground surface”. This very simple classification, carried out at points along transects, provides valuable information for characterizing soil wetness which can be used by permafrost modellers.

Soil

Background

Soil properties play a crucial role in the energy and water balance of permafrost systems, by affecting the exchange of heat and water between the atmosphere and the subsurface, and thus the rate of permafrost thaw (Shur and Jorgenson 2007; Chadburn et al. 2015; Walvoord and Kurylyk 2016). Permafrost-affected soil comprises a mixture of various media including organic matter, mineral particles ranging from gravel and sand to clay, as well as ice and unfrozen water. Organic matter insulates the permafrost from the air, the magnitude of the insulation depending on the organic layer thickness and organic matter content (Romanovsky et al. 2020). Soil texture also influences ground ice contents of permafrost, and together they control physical, thermal, and mechanical properties of permafrost and its behavior at thaw (French and Shur 2010; Jorgenson et al. 2010). Gravel or coarse sand show markedly different thermal and hydraulic properties compared with finer-grained soils (Shur and Jorgenson 2007). Soil texture also affects porosity, which determines the maximum amount of water that can be contained in a soil layer. Ice content and the form of the ice (such as ice lenses or massive ice) can affect energy transfer directly, as well as induce frost heave or subsidence of the ground surface in response to the formation or melting of the ice (Osterkamp et al. 2009; Kokelj and Jorgenson 2013; Romanovsky et al. 2020).

Measurement

Soil properties are documented as a one-time observation from a single measurement point near the transect. To characterize the soil profile (pedon), a soil pit is established close to the transect but set to the side to minimize disturbance. The pit should be approximately 1 m wide and 1 m deep, or until one can no longer dig due to frozen ground. The scale of 1 m was chosen to allow a clear soil profile to be revealed and the small-scale variability in soil properties to be accounted for. The best time is at the end of the growing season when thaw depth is greatest. If digging a pit is not allowed or possible, estimating the surface layer using a hand-held soil auger/drill is recommended.

The observations comprise a photograph of the clear profile and a description of visible characteristics, such as depth of organic layer, contents of ice and rocks, colour of the soil, and soil texture. For nonspecialists, we provide a simple hands-on flow chart within the the myThaw app that helps identification of soil texture (i.e., clay, silt, sand, gravel) adapting the protocol of the mySoil app (British Geological Survey 2021). Overall, the soil measurements are designed so that they do not require any specialist equipment or

laboratory analysis. To restore the site, the pit must be refilled and the organic mat reassembled.

Metadata, data quality, and storage


Metadata provide essential information about the quality, use, and genesis of the information being collected. Our metadata protocol complies with the standards of the Open Geospatial Consortium (OGC) ([Open Geospatial Consortium 2021](#)), and thus facilitates interoperability. Specifically, everything related to data processing and data management follows Observation to Archive (O2A; [Koppe et al. 2015](#); [Gerchow et al. 2017](#)), and in turn all instrumentation aspects of O2A follow sensorML specification ([OGC 2014](#)).

The protocol requests basic information about the site location, including latitude, longitude, elevation, and the location of the nearest weather station. This information is crucial for both mapping and modelling, and therefore adds greatly to the usability of the data collected. Land surface models require various forcing data, which can be obtained either from the nearest weather station, or in some cases from gridded products by using the nearest grid cell to the site. We then request an overview of the site characteristics as seen by eye, including whether the site is rocky, what type of soils are there, and how wet it is. For example, it may be a very wet or dry site, or it may be mixed, and these overview assessments, while providing similar information to the spheres themselves, will give an overview of the site as a whole. This also provides further information regarding how representative the transect measurements are. While vegetation height is covered in its own sphere, the dominant type of vegetation merits inclusion as metadata because it is a key indicator of the type of site. Basic information about any water features, such as ponds and rivers, as well as natural and anthropogenic disturbances are recorded because these will also affect the site, impacting the hydrology and permafrost thaw. Photos are required in the four cardinal directions in a standardized manner that provides a sense of scale, to give an overview of the site and clarify descriptions. An additional photo shows the placement of the transect.

The protocols are designed to ensure that the data and metadata meet scientific standards. We aim to provide quality-assured and data management over the whole data life cycle. Data should be findable, accessible, interoperable, and reusable according to the FAIR principles (Findability, Accessibility, Interoperability, Reusability; [Wilkinson et al. 2016](#)). Hence, measurement data and metadata need to be provided accurately and completely, have a persistent and unique identifier, and deposited in a trusted repository. It must follow the semantics of a standardized, controlled vocabulary to have broadly applicable language for machine access and processing. We apply the O2A dataflow framework, which includes the comprehensive description and management of all data with metadata, central data storage, and controlled data access ([Koppe et al. 2015](#); [Gerchow et al. 2017](#)). Through a standardized procedure, data uploads can be monitored in near-real time, and their spatial distribution visualized. The data can be accessed instantly as-is via the near-real time database ([Alfred Wegener Institute 2021](#)) while quality controlled and thematically curated datasets will be published in the PANGAEA long-term repositories ([Pangea 2021](#)), thus giving credit to the data provider in a data publication ([Schäfer et al. 2020](#)). A map-based search and visualization of the data with download link for the data (example: thaw depth) is planned. Data will be collected using a mobile app directly in the field. Data uplink occurs on-the-fly or whenever the data collector can upload it to an AWI server and will be automatically ingested into the O2A process chain ([Fig. 3](#)).

For quality control, a first quality check is done automatically using the O2A system, such as removing unphysical data (for example, negative snow depths) or implausible coordinates and times. This is managed by setting the measurement properties in

Fig. 3. Illustration showing the workflow of the data collection (myThaw app) and O2A (Alfred Wegener Institute 2021) process chain towards archiving into a repository. Data are collected offline and ingested into O2A in “delayed mode” (as soon as internet access is available) using full metadata annotation. A dashboard is used for visualization of the data once they are uploaded. Data can be visualized spatially on the Portal. Final publications take place in the repositories. Figure adapted after Koppe et al. (2015).



sensor.awi.de. Before archiving the data set in PANGAEA, an additional thorough manual data check will be done.

Description of the mobile myThaw app for data collection

The mobile app myThaw is freely available to everybody (Appendix A). The app allows the collected data to be exported to a central data storage for data analysis and reporting. One of the advantages of apps is the possibility of gathering data offline or while on-the-go. The offline form allows researchers to collect and store data while in the field and upload it once an internet connection is available (for example, at the field station). As nearly all researchers and citizens today own a smartphone or tablet, we see advantages in using a mobile over a field notebook or report-based archives. The app is designed for use in cold climates and is user friendly, with help/guidelines and “pop-up window” options when necessary. Because our protocol asks for measurements at multiple moments across time and spheres, at new and recurring locations (i.e., long term measurements at the same sites), the app can identify the recurring location, thus eliminating the need to re-enter the metadata. The app will be available under the CC BY 4.0 licence. Further maintenance and development, such as security updates and, if necessary, debugging, are planned for the future. In summary, we provide a method for secure and collaborative data entry, resulting in faster data analyses, visualization, access, and storage.

Next steps for the data: conclusions and outlook

The database that we will develop using this protocol and app will cover permafrost state and land-surface conditions. The value of this is not only in analysing the trends and relationships in this dataset alone, which can be used for model validation and parameterisation, but it can also be analysed in combination with other datasets, for example atmospheric conditions, permafrost types, and remote sensing data including vegetation maps and topography (Nitze et al. 2018; Raynolds et al. 2019).

Further developments could also link our protocol to water, soils, and sediment sampling. For example, the action group called Standardized methods across Permafrost

Arctic Science Downloaded from cdnsciencepub.com by UNIVERSITY OF OSLO on 02/28/22 For personal use only.

Landscapes: from Arctic Soils to Hydrosystems (SPLASH) is currently working on a standardized protocol for sampling mineral and organic components in soils, sediments, and water across permafrost landscapes (Bouchard et al. 2020).

We present a set of simple protocols for observing permafrost thaw and associated environmental conditions. The protocols cover permafrost, snow, vegetation, water, and soil. They are unique in that they:

- are for everyone: no knowledge or sophisticated equipment is needed;
- encompass multiple critical parameters, so that the drivers and controls of permafrost thaw can be quantified;
- come with an app that guides the user through the measurement process and guarantees data quality, consistency and accessibility.

The protocols address the urgent need for high-quality field observations of permafrost conditions and interlinked ecosystem parameters. The observations will be critical for understanding and predicting permafrost thaw and for establishing a baseline for quantifying future change. The consistency and accessibility of the observations is crucial for data-driven analyses. The dataset will serve to enhance and validate Earth system models and remote sensing methods that are indispensable for monitoring and projecting permafrost thaw across the Arctic.

The current protocol has already been implemented by some INTERACT sites and data will be collected in 2021. The next steps include sharing it with a wider group of scientists and the public, for example to colleagues, the Permafrost Young Researchers Network, Cryolist server, and sharing on social media. The protocol should be distributed to researchers and citizen scientists to obtain data on snow, vegetation, soil, and thaw depth at locations around the Arctic. Future work will include a linked higher level protocol that includes measurements, for example of ground subsidence and soil temperatures for which more advanced instruments, techniques, and expertise are required. More widely, similar integrated protocols that address carbon and nutrient cycling would also be of great value in monitoring the permafrost landscape. This will require coordination with recent calls for standardized monitoring initiatives of other aspects of Arctic environments, including the need for standardized protocols for Arctic freshwater initiatives (Heino et al. 2020) or for SPLASH (Bouchard et al. 2020).

Beyond these community-led initiatives, national infrastructure funding for permanent monitoring sites is needed to understand long term permafrost thaw.

Acknowledgements

We acknowledge the following grants and funding for this work:

J.B. and J.M.: Helmholtz Association in the framework of MOSES (Modular Observation Solutions for Earth Systems); D.S.: National Science Foundation (NSF) 1836377 and 2019691; I.A.: Research Council of Norway (RCN), KLIMAFORSK programme, RCN project number 294948 (2019–2022); H.L.: RCN, INTPART program, RCN project number 309625; S.M.S.: European Research Council (ERC) consolidator grant Glacial Legacy of Ulrike Herzschuh (grant no. 772852) and Federal Ministry of Education and Research (BMBF) of Germany grant to Moritz Langer (No. 01LN1709A); E.J.W.: W. Garfield Weston Award for Northern Research (PhD), Ontario Graduate Scholarship; S.E.C.: Natural Environment Research Council (NERC) grant No. NE/R015791/1; S.W.: European Space Agency (ESA) Permafrost CCI; Nunataryuk (EU grant agreement No. 773421); S.Z.: National Aeronautics and Space Administration (NASA) 80NSSC19K1494.

Valuable comments to the initial protocol were contributed by Annett Bartsch, Liane Benning, Bill Cable, Matthias Fuchs, Birgit Heim, Stefan Kruse, Susanne Liebner, Maribeth Murray, Jan Nitzbon, Peter Pulsifer, Gabriela Schaeppman-Strub, Warwick Vincent, and Caroline Duchesne. The app was developed by Ole Eckermann. The video tutorial from Işkoras, Norway, September 2021 was recorded by E. Audunsdottir and S. Azanova and edited by S. Azanova. The video tutorial from Bayelva, Svalbard, Norway, March 2021, was recorded by E. Horvath and edited by E. Horvath and L. Grübner. We thank the following people for the mySoil material: Lawley, R., B.A. Emmett, D.A. Robinson. 2014. Soil Observatory lets researchers dig deep. *Nature* 509, 427 and Shelley, W., R. Lawley and D.A. Robinson. 2013. Crowd-sourced soil data for Europe. *Nature*, 496:300. We acknowledge support by the Open Access Publication Funds of Alfred Wegener Institute Helmholtz Center for Polar and Marine Research.

Author contributions

J.B., S.C., and S.Z. conceived the idea and conceptualization for the protocol and paper. The original draft and outline of the paper and protocol were prepared by J.B., S.Z., S.C., and J.M. The individual sections with details of the spheres in the paper and protocol were contributed by the following: Snow, J.M. and M.S.; Permafrost, S.Z., J.B., E.J.W., D.S., and S.S.; Vegetation, I.A. and S.M.S.; Water, Y.S. and A.L.; Soil, J.B., S.C., and H.L.; Metadata, S.C., L.C., N.S., and S.W.; Data collection, transfer, and storage, J.B. and N.A. All of the other sections were written by J.B., S.C., and S.Z. The figures were drawn by J.M. with input from J.B., S.C., S.Z., and N.S.; J.B., S.C., S.Z., and J.M. organized the writing and contributions from the co-authors. Review and editing of the various versions of the paper were provided by J.B., S.Z., S.C., J.B., J.M., S.B., I.A., and N.S.; N.A. set up the O2A data flow with inputs from J.B. The video tutorial from Işkoras was organized by I.A. and H.L. The video tutorial from Bayelva was organized by J.B. and J.M. All of the co-authors approved the final version of the manuscript.

References

- Aartsma, P., Asplund, J., Odland, A., Reinhardt, S., and Renssen, H. 2020. Surface albedo of alpine lichen heaths and shrub vegetation. *Arctic, Antarctic, Alpine Res.* **52**(1): 312–322. doi: [10.1080/15230430.2020.1778890](https://doi.org/10.1080/15230430.2020.1778890).
- Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research. 2021. Observatory to Archive (O2A). Available from <https://data.awi.de/?site=home> [accessed 07 January 2021].
- Arctic Development and Adaptation to Permafrost in Transition (ADAPT). ADAPT Standard protocols. Available from <http://www.cen.ulaval.ca/adapt/protocols/adapt.php> [accessed 07 January 2021].
- Arctic Program 2020. Arctic Report Card. Available from <https://arctic.noaa.gov/Report-Card> [accessed 07 January 2021].
- Blok, D., Heijmans, M.M.P.D., Schaeppman-Strub, G., Kononov, A.V., Maximov, T.C., and Berendse, F. 2010. Shrub expansion may reduce summer permafrost thaw in Siberian tundra. *Global Change Biol.* **16**: 1296–1305. doi: [10.1111/j.1365-2486.2009.02110.x](https://doi.org/10.1111/j.1365-2486.2009.02110.x).
- Boike, J., Roth, K., and Ippisch, O. 2003. Seasonal snow cover on frozen ground: Energy balance calculations of a permafrost site near Ny-Ålesund, Spitsbergen. *J. Geophys. Res.: Atmosph.* **108**(D2): 8163. doi: [10.1029/2001JD000939](https://doi.org/10.1029/2001JD000939).
- Bouchard, F., Agnan, Y., Bröder, L., Fouché, J., Hirst, C., Sjöberg, Y., et al. 2020. The SPLASH Action Group – Towards standardized sampling strategies in permafrost science, *Adv. Polar Sci.* **31** (3): 153–155. doi: [10.13679/j.advps.2020.0009](https://doi.org/10.13679/j.advps.2020.0009).
- Bonan, G.B., and Shugart, H.H. 1989. Environmental Factors and Ecological Processes in Boreal Forests. *Annual Rev. Ecol. System.* **20**(1): 1–28. doi: [10.1146/annurev.es.20.110189.000245](https://doi.org/10.1146/annurev.es.20.110189.000245).
- British Geological Survey. 2021. mySoil Growing our knowledge (version 5.0) [Mobile App]. UKRI. Available from <https://www.bgs.ac.uk/mysoil> [accessed 16 February 2021].
- Brown, J., Hinkel, K.M., and Nelson, F.E. 2000. The circumpolar active layer monitoring (calm) program: Research designs and initial results. *Polar Geogr.* **24**(3): 166–258. doi: [10.1080/10889370009377698](https://doi.org/10.1080/10889370009377698).
- Burke, E.J., Ekici, A., Huang, Y., Chadburn, S.E., Huntingford, C., Ciais, P., et al. 2017. Quantifying uncertainties of permafrost carbon–climate feedbacks. *Biogeosciences*, **14**(12): pp. 3051–3066.
- Callaghan, T.V., Johansson, M., Brown, R.D., Groisman, P.Y., Labba, N., Radionov, V., et al. 2011. The changing face of arctic snow cover: a synthesis of observed and projected changes. *Ambio*, **40**: 17–31. doi: [10.1007/s13280-011-0212-y](https://doi.org/10.1007/s13280-011-0212-y).

- Chadburn, S.E., Burke, E.J., Essery, R.L.H., Boike, J., Langer, M., Heikenfeld, M., et al. 2015. Impact of model developments on present and future simulations of permafrost in a global land-surface model. *Cryosphere*, **9**(4): 1505–1521.
- Chang, X., Jin, H., Zhang, Y., He, R., Luo, D., Wang, Y., et al. 2015. Thermal impacts of boreal forest vegetation on active layer and permafrost soils in northern da Xing'Anling (Hinggan) Mountains, northeast China. *Arctic, Antarctic, and Alpine Research*, **47**(2): 267–279. doi: [10.1657/AAAR00C-14-016](https://doi.org/10.1657/AAAR00C-14-016).
- Circumpolar Active Layer Monitoring Network (CALM). [Online]. Soil moisture content. Available from https://www2.gwu.edu/~calm/research/soil_moisture.html [accessed 08 January 2021].
- Crumley, R.L., Hill, D.F., Wikstrom Jones, K., Wolken, G.J., Arendt, A.A., Aragon, C.M., et al. 2020. Assimilation of citizen science data in snowpack modeling using a new snow dataset: Community Snow Observations. *Hydrol. Earth Syst. Sci. Discuss.* 1–39. doi: [10.5194/hess-2020-321](https://doi.org/10.5194/hess-2020-321).
- Domine, F., Belke-Brea, M., Sarrazin, D., Arnaud, L., Barrere, M., and Poirier, M. 2018. Soil moisture, wind speed and depth hoar formation in the Arctic snowpack. *J. Glaciol.* **64**(248): 990–1002. doi: [10.1017/jog.2018.89](https://doi.org/10.1017/jog.2018.89).
- Edlefsen, N.E., and Anderson, A.B.C. 1943. Thermodynamics of soil moisture. *Hilgardia*, **15**(2) 31–298. doi: [10.3733/hilg.v15n02p031](https://doi.org/10.3733/hilg.v15n02p031).
- Fierz, C., Armstrong, R.L., Durand, Y., Etchevers, P., Greene, E., McClung, D.M. et al. 2009. The International classification for seasonal snow on the ground. Available from <https://unesdoc.unesco.org/ark:/48223/pf0000186462> [accessed 07 January 2021].
- Food and Agriculture Organization of the United Nations (FAO). [Online]. Planning and making a soil survey. Available from http://www.fao.org/tempref/FI/CDrom/FAO_Training/FAO_Training/General/x6706e/x6706e02.htm [accessed 07 January 2021].
- French, H., and Shur, Y. 2010. The principles of cryostratigraphy. *Earth Sci. Rev.* **101**(3-4): 190–206. doi: [10.1016/j.earscirev.2010.04.002](https://doi.org/10.1016/j.earscirev.2010.04.002).
- Gerchow, P., Koppe, R., Macario, A., Haas, A., Schäfer-Neth, C., Pfeiffenberger, H., and Schäfer, A. 2017. O2A — data flow framework from sensor observations to archives. epic3 digital infrastructures for research, di4r conference, Brussels. Available from <https://indico.egi.eu/indico/event/3455/session/1/contribution/114/material/slides/1.pdf> [accessed 18 May 2021].
- Grogan, P., Henry, G., Grant, R., and Levesque, H. [Online]. ADAPT vegetation standard description protocol. Available from <https://www.cen.ulaval.ca/adapt/protocols/adapt.php> [accessed 07 January 2021].
- Grünberg, I., Wilcox, E., Zwieback, S., Marsh, P., and Boike, J. 2020. Linking tundra vegetation, snow, soil temperature, and permafrost. *Biogeosciences*, **17**(16): 4261–4279. doi: [10.5194/bg-17-4261-2020](https://doi.org/10.5194/bg-17-4261-2020).
- Haberkorn, A. 2019. European snow booklet — an inventory of snow measurements in Europe. *EnviDat*. doi: [10.16904/envidat.59](https://doi.org/10.16904/envidat.59).
- Heino, J., Culp, J.M., Erkinaro, J., Goedkoop, W., Lento, J., Rühland, K.M., and Smol, J.P. 2020. Abruptly and irreversibly changing Arctic freshwaters urgently require standardized monitoring. *J. Appl. Ecol.* **57**: 1192–1198. doi: [10.1111/1365-2664.13645](https://doi.org/10.1111/1365-2664.13645).
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R.B., Jones, M., MacDonald, G., et al. 2020. Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. *Proceedings of the National Academy of Sciences*, **117**(34): 20438–20446. doi: [10.1073/pnas.1916387117](https://doi.org/10.1073/pnas.1916387117).
- Jonas, T., and Marks, D. 2016. Estimating the snow water equivalent from snow depth data. Available from https://iahs.info/uploads/ICSIH_upload/articles/no1_2016/ICSIH%20article%20no1.pdf [accessed 05 January 2021].
- Jorgenson, M.T., Romanovsky, V., Harden, J., Shur, Y., O'Donnell, J., Schuur, E.A.G., et al. 2010. Resilience and vulnerability of permafrost to climate change. *Can. J. For. Res.* **40**(7): 1219–1236. doi: [10.1139/X10-060](https://doi.org/10.1139/X10-060).
- Juszkak, I., Eugster, W., Heijmans, M.M.P.D., and Schaepman-Strub, G. 2016. Contrasting radiation and soil heat fluxes in Arctic shrub and wet sedge tundra. *Biogeosci.* **13**(13): 4049–4064. doi: [10.5194/bg-13-4049-2016](https://doi.org/10.5194/bg-13-4049-2016).
- Kokelj, S.V., and Jorgenson, M.T. 2013. Advances in thermokarst research. *Permafrost Periglacial Processes* **24**: 108–119. doi: [10.1002/ppp.1779](https://doi.org/10.1002/ppp.1779).
- Koppe, R., Gerchow, P., Macario, A., Haas, A., Schäfer-Neth, C., and Pfeiffenberger, H. 2015. O2A: A generic framework for enabling the flow of sensor observations to archives and publications. *OCEANS Genova conference*, Genoa, 18-21 May 2015, pp. 1–6. doi: [10.1109/OCEANS-Genova.2015.7271657](https://doi.org/10.1109/OCEANS-Genova.2015.7271657).
- Kruse, S., Bolshiyarov, D., Grigoriev, M.N., Morgenstern, A., Pstryakova, L., Tsibizov, L., and Udke, A. 2019. Russian–German Cooperation: Expeditions to Siberia in 2018. *Berichte zur Polar- und Meeresforschung (Reports on polar and marine research)*, **734**: 257. doi: [10.2312/BzPM_0734_2019](https://doi.org/10.2312/BzPM_0734_2019).
- Kurylyk, B.L., and Watanabe, K. 2013. The mathematical representation of freezing and thawing processes in variably-saturated, non-deformable soils. *Adv. Water Resour.* **60**: 160–177. doi: [10.1016/j.advwatres.2013.07.016](https://doi.org/10.1016/j.advwatres.2013.07.016).
- Liljedahl, A.K., Timling, I., Frost, G.V., and Daanen, R.P. 2020. Arctic riparian shrub expansion indicates a shift from streams gaining water to those that lose flow. *Communications Earth & Environment*, **1**(50): 1–9. doi: [10.1038/s43247-020-00050-1](https://doi.org/10.1038/s43247-020-00050-1).
- Martin, L.C.P., Nitzbon, J., Aas, K.S., Etzelmüller, B., Kristiansen, H., and Westermann, S. 2019. Stability conditions of peat plateaus and palsas in northern Norway. *J. Geophys. Res. Earth Surf.* **124**(3): 705–719. doi: [10.1029/2018JF004945](https://doi.org/10.1029/2018JF004945).
- Molau, U. 1996. *International Tundra Experiment (ITEX) manual*, 2nd ed. Chapter 5: Snow and Ice. Available from <https://www.gvsu.edu/itex/library-8.htm> [accessed 05 January 2021].

- Molau, U., and Edlund, S. 1996. International Tundra Experiment (ITEX) manual. 2nd ed. Chapter 8: Plant Response Variables. Available from <https://www.gvsu.edu/itex/library-8.htm> [accessed 05 January 2021].
- Myers-Smith, I.H., Forbes, B.C., Wilmling, M., Hallinger, M., Lantz, T., Blok, D., et al. 2011. Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environ. Res. Lett.* **6**(4): 1–15. doi: [10.1088/1748-9326/6/4/045509](https://doi.org/10.1088/1748-9326/6/4/045509).
- Nelson, F.E., and Hinkel, K.M. 2004. Methods for measuring active-layer thickness. In *A Handbook on Periglacial Field Methods*. Edited by O. Humlum, and N. Matsuoka. Available from <https://ipa.arcticportal.org/publications/handbook> [accessed 07 January 2021].
- Nelson, F., Brown, J., Lewkowicz, T., and Taylor, A. 1996. Chapter 6: Active Layer Protocol. In *International Tundra Experiment (ITEX) manual*, 2nd ed. Available from https://www2.gwu.edu/~calm/research/active_layer.html [accessed 05 January 2021].
- Nitzbon, J., Langer, M., Martin, L.C.P., Westermann, S., Schneider von Deimling, T., and Boike, J. 2020. Effects of multi-scale heterogeneity on the simulated evolution of ice-rich permafrost lowlands under a warming climate. *The Cryosphere*, **3**(15): 1399–1422. doi: [10.5194/tc-15-1399-2021](https://doi.org/10.5194/tc-15-1399-2021).
- Nitze, I., Grosse, G., Jones, B.M., Romanovsky, V.E., and Boike, J. 2018. Remote sensing quantifies widespread abundance of permafrost region disturbances across the Arctic and Subarctic. *Nat Commun.* **9**: 5423. doi: [10.1038/s41467-018-07663-3](https://doi.org/10.1038/s41467-018-07663-3).
- O'Connor, M.T., Cardenas, M.B., Ferencz, S.B., Wu, Y., Neilson, B.T., Chen, J., and Kling, G.W. 2020. Empirical models for predicting water and heat flow properties of permafrost soils. *Geophys. Res. Lett.* **47**(11). doi: [10.1029/2020GL087646](https://doi.org/10.1029/2020GL087646).
- O'Neill, H.B., Smith, S.L., and Duchesne, C. 2019. Long-term permafrost degradation and thermokarst subsidence in the Mackenzie Delta area indicated by thaw tube measurements. In *Proceedings of the 18th International Conference on Cold Regions Engineering and the 8th Canadian Permafrost Conference*. 18–22 August 2019. Edited by J.-P. Bilodeau, D.F. Nadeau, D. Fortier, and D. Conciatori. American Society of Civil Engineers, Quebec City, Canada. pp. 643–651.
- Open Geospatial Consortium. 2014. OGC SensorML: Model and XML Encoding Standard. Available from https://portal.opengeospatial.org/files?artifact_id=55939 [accessed 18 May 2021].
- Open Geospatial Consortium. 2021. OGC Standards. Available from <https://www.ogc.org/docs/ogc> [accessed 05 January 2021].
- Osterkamp, T.E., Jorgenson, M.T., Schuur, E.A., Shur, Y.L., Kanevskiy, M.Z., Vogel, J.G., and Tumskey, V.E. 2009. Physical and ecological changes associated with warming permafrost and thermokarst in Interior Alaska. *Permafrost Periglacial Proc.* **20**(3): 235–256. doi: [10.1002/ppp.656](https://doi.org/10.1002/ppp.656).
- Outcalt, S.I., Nelson, F.E., and Hinkel, K.M. 1990. The zero-curtain effect: Heat and mass transfer across an isothermal region in freezing soil. *Water Resour. Res.* **26**(7): 1509–1516. doi: [10.1029/WR026i007p01509](https://doi.org/10.1029/WR026i007p01509).
- Pangea. 2021. Data Publisher for Earth & Environmental Science. Available from <https://www.pangaea.de/> [accessed 05 January 2021].
- Parr, C., Sturm, M., and Larsen, C. 2020. Snowdrift Landscape Patterns: An Arctic Investigation. *Water Resour. Res.* **56**(12). doi: [10.1029/2020WR027823](https://doi.org/10.1029/2020WR027823).
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., et al. 2013. New handbook for standardised measurement of plant functional traits worldwide. *Aust. J. Bot.* **61**: 167–234. doi: [10.1071/bt12225](https://doi.org/10.1071/bt12225).
- Ping, C.-L., Clark, M.H., Kimble, J.M., Michaelson, G.J., Shur, Y., and Stiles, C. 2013. Sampling protocols for permafrost-affected soils. *Soil Horizons*, **54**(1): 13–19. doi: [10.2136/sh12-09-0027](https://doi.org/10.2136/sh12-09-0027).
- Pulliaainen, J., Luojus, K., Derksen, C., Mudryk, L., Lemmetyinen, J., Salminen, M., et al. 2020. Patterns and trends of Northern Hemisphere snow mass from 1980 to 2018. *Nature*, **581**(7808): 294–298. doi: [10.1038/s41586-020-2258-0](https://doi.org/10.1038/s41586-020-2258-0).
- Rasmussen, L.H., Zhang, W., Hollesen, J., Cable, S., Christiansen, H.H., Jansson, P.-E., and Elberling, B. 2018. Modelling present and future permafrost thermal regimes in Northeast Greenland. *Cold Reg. Sci. Technol.* **146**: 199–213. doi: [10.1016/j.coldregions.2017.10.011](https://doi.org/10.1016/j.coldregions.2017.10.011).
- Raynolds, M.K., Walker, D.A., Balsler, A., Bay, C., Campbell, M., Cherosov, M.M., et al. 2019. A raster version of the Circumpolar Arctic Vegetation Map (CAVM). *Remote Sens. Environ.* **232**: 111297.
- Riseborough, D., Shiklomanov, N., Eitzmüller, B., Gruber, S., and Marchenko, S. 2008. Recent advances in permafrost modelling. *Permafrost Periglacial Process.* **19**(2): 137–156. doi: [10.1002/ppp.615](https://doi.org/10.1002/ppp.615).
- Romanovsky, V.E., Smith, S.L., Isaksen, K., Nyland, K.E., Kholodov, A.L., Shiklomanov, N.I., et al. 2020. Terrestrial permafrost. In *State of the Climate in 2019*. *Bull. Am. Meteorol. Soc.* **101**(8): 265–271. doi: [10.1175/BAMS-D-20-0086.1](https://doi.org/10.1175/BAMS-D-20-0086.1).
- Schäfer, A., Anselm, N., Eilers, J., Frickenhaus, S., Gerchow, P., Glöckner, F.O., et al. 2020. Implementing FAIR in a Collaborative Data Management Framework. EGU Copernicus Meetings. Available from <https://meetingorganizer.copernicus.org/EGU2020/EGU2020-19631.html> [accessed 07 January 2021].
- Shiklomanov, N.I., Streletskiy, D.A., Nelson, F.E., Hollister, R.D., Romanovsky, V.E., Tweedie, C.E., et al. 2010. Decadal variations of active-layer thickness in moisture-controlled landscapes, Barrow, Alaska. *J. Geophys. Res.* **115**(G00104). doi: [10.1029/2009JG001248](https://doi.org/10.1029/2009JG001248).
- Shur, J., and Jorgenson, T. 2007. Patterns of permafrost formation and degradation in relation to climate and ecosystems. *Permafrost Periglacial Process.* **18**: 7–19. doi: [10.1002/ppp.582](https://doi.org/10.1002/ppp.582).
- Smith, S., and Brown, J. 2009. Assessment of the status of the development of the standards for the Terrestrial Essential Climate Variables - T7 - Permafrost and seasonally frozen ground. Available from <http://library.arcticportal.org/668/> [accessed 07 January 2021].
- Soil Survey Staff. 2014. Soil survey field and laboratory methods manual. Soil Survey Investigations Report No. 51, Version 2.0. Edited by R. Burt and Soil Survey Staff. U.S. Department of Agriculture, Natural Resources

- Conservation Service. Available from https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1244466.pdf [accessed 08 January 2021].
- Stiegler, C., Lund, M., Christensen, T.R., Mastepanov, M., and Lindroth, A. 2016. Two years with extreme and little snowfall: effects on energy partitioning and surface energy exchange in a high-Arctic tundra ecosystem. *Cryosphere*, **10**(4): 1395–1413. doi: [10.5194/tc-10-1395-2016](https://doi.org/10.5194/tc-10-1395-2016).
- Streletskiy D., Biskaborn, B., Smith, S., Noetzi, J., Viera, G., and Schoeneich, P. 2017a. Strategy and implementation plan 2016–2020 for the Global Terrestrial Network for Permafrost (GTN-P). Available from <http://library.arcticportal.org/1938/> [accessed 06 January 2021].
- Streletskiy, D.A., Shiklomanov, N.I., Little, J.D., Nelson, F.E., Brown, J., Nyland, K.E., and Klene, A.E. 2017b. Thaw subsidence in undisturbed tundra landscapes, Barrow, Alaska, 1962–2015. *Permafrost Periglacial Process*. **28**(3): 566–572. doi: [10.1002/ppp.1918](https://doi.org/10.1002/ppp.1918).
- Sturm, M., Liston, G.E., Benson, C.S., and Holmgren, J. 2001a. Characteristics and growth of a snowdrift in Arctic Alaska, USA. *Arctic, Antarctic, Alpine Res.* **33**(3): 319–329. doi: [10.1080/15230430.2001.12003436](https://doi.org/10.1080/15230430.2001.12003436).
- Sturm, M., Holmgren, J., McFadden, J.P., Liston, G.E., Chapin, F.S., and Racine, C.H. 2001b. Snow–shrub Interactions in Arctic Tundra: a hypothesis with climatic implications. *J. Clim.* **14**(3): 336–344. doi: [10.1175/1520-0442\(2001\)014<0336:SSIIAT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<0336:SSIIAT>2.0.CO;2).
- Stuenzi, S.M., Boike, J., Cable, W., Herzsuh, U., Kruse, S., Pestryakova, L.A., et al. 2021. Variability of the surface energy balance in permafrost-underlain boreal forest. *Biogeosciences*, **18**(2): 343–365. doi: [10.5194/bg-18-343-2021](https://doi.org/10.5194/bg-18-343-2021).
- The Global Climate Observing System. 2016a. ECV Products and Requirements for Snow. Available from <https://gcos.wmo.int/en/essential-climate-variables/snow/ecv-requirements> [accessed 05 January 2021].
- The Global Climate Observing System. 2016b. ECV Products and Requirements for Permafrost. Available from <https://gcos.wmo.int/en/essential-climate-variables/permafrost/ecv-requirements> [accessed 05 January 2021].
- The National Ecological Observing Network (NEON). 2021. TIS Soil Pit Sampling Protocol. Available from https://data.neonscience.org/documents/-/document_library_display/EygRkSpUBoq/view/1883155 [accessed 07 January 2021].
- Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A., et al. 2020. Carbon release through abrupt permafrost thaw. *Nat. Geosci.* **13**(2):138–143. doi: [10.1038/s41561-019-0526-0](https://doi.org/10.1038/s41561-019-0526-0).
- United Nations 2012. The United Nations terminology database. Available from <https://unterm.un.org/unterm/portal/welcome> [accessed 07 January 2021].
- Vincent, W.F., Lemay, M., and Allard, M. 2017. Arctic permafrost landscapes in transition: towards an integrated Earth system approach. *Arct. Sci.* **3**(2): 39–64. doi: [10.1139/as-2016-0027](https://doi.org/10.1139/as-2016-0027).
- Walker, D.A., Jia, G.J., Epstein, H.E., Reynolds, M.K., Iii, F.S.C., Copass, C., et al. 2003. Vegetation-soil-thaw-depth relationships along a low-arctic bioclimate gradient, Alaska: synthesis of information from the ATLAS studies. *Permafrost Periglacial Process*. **14**(2): 103–123. doi: [10.1002/ppp.452](https://doi.org/10.1002/ppp.452).
- Walvoord, M.A., and Kurylyk, B.L. 2016. Hydrologic impacts of thawing permafrost—a review. *Vadose Zone J.* **15**(6). doi: [10.2136/vzj2016.01.0010](https://doi.org/10.2136/vzj2016.01.0010).
- Wilcox, E.J., Keim, D., de Jong, T., Walker, B., Sonnentag, O., Sniderhan, A.E., et al. 2019. Tundra shrub expansion may amplify permafrost thaw by advancing snowmelt timing. *Arct. Sci.* **5**(4): 202–217. doi: [10.1139/as-2018-0028](https://doi.org/10.1139/as-2018-0028).
- Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J., Appleton, G., Axton, M., Baak, A., et al. 2016. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*, **3**(1): 160018. doi: [10.1038/sdata.2016.18](https://doi.org/10.1038/sdata.2016.18).
- Woo, M. 2012. *Permafrost Hydrology*. Springer, Berlin, Heidelberg, Luxembourg, pp. 576.
- World Meteorological Organization 2008. Guide to hydrological parameters – Volume 1. Available from https://www.wmo.int/pages/prog/hwrr/publications/guide/english/168_Vol_1_en.pdf [accessed 08 January 2021].
- Zhang, T. 2005. Influence of the seasonal snow cover on the ground thermal regime: An overview. *Rev. Geophys.* **43**(4). doi: [10.1029/2004RG000157](https://doi.org/10.1029/2004RG000157).
- Zhang, T., Osterkamp, T.E., and Stamnes, K. 1996. Influence of the depth hoar layer of the seasonal snow cover on the ground thermal regime. *Water Resour. Res.* **32**(7): 2075–2086. doi: [10.1029/96WR00996](https://doi.org/10.1029/96WR00996).

Appendix A

Protocol: T-MOSAIc permafrost thaw


We provide a user-friendly application (app) named myThaw for smartphones, tablets and personal computers along with this protocol, which you can use to enter your data and upload it to the T-MOSAIc permafrost thaw database. The app will guide you through the measurement.

Download the myThaw app here:

<https://play.google.com/store/apps/details?id=de.awi.permafrost>

<https://apps.apple.com/app/mythaw/id1578278222>

Fig. A1. An overview of the measurements to take along one transect over one seasonal cycle.



Video tutorials here:

Ískoras, Norway, September 2020: <https://youtu.be/zTsk5NWmkdk>


Bayelva, Svalbard, Norway, March 2021: <https://youtu.be/G5dbh6Pix8o>

Equipment needed (all measurements are in metric (SI) units)

- Your smartphone or any other device with the myThaw app installed.
- Alternatively, a weatherproof notebook and pencil (do not use a regular pen, the ink smears); you can enter the data from the field in the app later.
- Foldable ruler (1 or 2 m long) and tape measure (30 m long).
- Pointed metal rod (frost probe), around 1 cm thick, about 1 or 2 m long; if not graduated, additional measuring tape.
- Smaller poles to leave at the site to mark the beginning and end of the transect.
- Camera or mobile phone with a camera.
- A spade or shovel for digging. A hand saw or bread knife can also be very useful for this.

The measurement frequency will of course depend on your capacities. Please see our recommendations on the measurement intervals as a “best case” scenario! We appreciate any measurements — if you can’t take measurements as often as recommended, your data will still be valuable.

Fig. A2. An example of how you could label your transect photo.



Section 0. How to locate your measurements

The overall aim of this project is to map and monitor permafrost thaw at as many sites as possible around the Arctic. Before taking any measurements, you need to select a location for the transect. All of the measurements that are taken at more than one point (everything except the soil pit, Section on *Permafrost – thaw depth*), should be measured along a single transect that you can return to each time you take measurements (Fig. A1). We recommend that the transect has a minimum of 10 measurement points spaced 1 m apart (a 10 m transect), but preferably 30 measurement points spaced 1 m apart (a 30 m transect). Choose a transect by considering accessibility and representativeness: a place which is “typical for the site/landscape” and orient the transect to encompass the variability present at the site. Please take a photo of the transect, and if at all possible annotate this photo with numbered measurement points. *It is important to be very careful when taking measurements or walking near your transect, to avoid damaging the plants, soil, and snow.* The more often measurements are taken the more disturbance on the site might appear over time.

Note that you must keep the numbering of the measurement points consistent when entering data in the app. If you cannot obtain a measurement at any point, just leave the appropriate box empty.

Optionally, indicate any additional information on the photo that you can — such as higher/middle/lower ground (see Fig. A2). You can set up more than one transect if you want to, and assign a number to each one.

Actions: Mark the beginning and end of each transect with a pole or similar (Fig. A3). In presence of rocky ground or a basal ice layer it may be necessary to use drilling equipment. To prevent microplastic pollution, please do not use plastic tape or flags.

- Record the GPS coordinates of the middle of the transect using your phone. Alternatively, you could also use e.g., Google Maps on your smartphone by holding your finger in your current location to drop a pin, and then swiping up to see the coordinates of the dropped pin.
- Make sure you do not walk in your transect, especially in winter, for snow depth measurements.

Fig. A3. Measuring snow depth on the transect during winter. This illustrates what a transect might look like.



Metadata

When to fill out: All information needs to be provided once for each transect, and updated once per year if there is any change.

1. Date of measurements*:
2. Name of the site:
3. Plot ID (if there is more than one study location or transect at your site):
4. Latitude of your site (a decimal number between -90 degrees to 90 degrees):
Longitude of your site (between -180 to 180):
5. Elevation of your site (metres above sea level):
6. Are you aware of any nearest official or national weather service station
 Yes No
 If Yes, answer the following:
 Name/ID of station and name of monitoring network:
 Distance to this weather station (if known, in m):
 Latitude of this weather station (between -90 and 90):
 Longitude of this weather station (between -180 to 180):
7. Are there climate data available at the site itself? Yes No
 If Yes, answer the following:
 Distance from your transect to closest climate measurement location [m] :
 Latitude of climate measurement location:
 Longitude of climate measurement location:
 Elevation of the climate measurement location:
 Variables measured (tick all that apply)
 Air temperature
 Wind speed
 Air pressure

Arctic Science Downloaded from cdnsciencepub.com by UNIVERSITY OF OSLO on 02/28/22
For personal use only.

- Humidity
- Shortwave radiation
- Longwave radiation
- Rainfall
- Snowfall
- Snow depth

8. How do you access your site?

9. How far to the nearest road [m] ?

What do you see at the site (30 m × 30 m area)?

10. Ground surface (the layer below the vegetation)

- Rock
- Soil

If you ticked "soil" in the previous question, tick all that apply:

- Peat
- Gravel
- Sand
- Silt
- Clay
- Unknown

11. How wet is the ground?

- Wet (water above the surface)
- Moist (soils are damp)
- Dry
- Unknown

12. Water features (tick all that apply):

- Wetland
- Lake
- Wet depressions
- River/creek
- Water tracks
- None
- Unknown
- Other: _____

13. Are there trees at the site?


- Yes
- No

14a. Most dominant vegetation at the site:

You can use the flowchart to identify it (Fig. A4). Tick one checkbox.

- Grasses/sedges
- Forbs
- Deciduous Shrubs (e.g., *Vaccinium* sp.)
- Evergreen Shrubs (e.g., crowberry, *Empetrum* sp.)
- Moss
- Lichen
- Trees: Deciduous needle (e.g., larch)
- Trees: Evergreen needle (e.g., spruce)
- Trees: Broadleaf (e.g., birch)
- There is no vegetation

Fig. A4. Flowchart vegetation.



14b. Indicate any other vegetation types that are present at the site:

- Grasses/sedges
- Forbs
- Deciduous Shrubs (e.g., *Vaccinium* sp.)
- Evergreen Shrubs (e.g., crowberry (*Empetrum* sp.))
- Moss
- Lichen
- Trees: Deciduous needle (e.g., larch)
- Trees: Evergreen needle (e.g., spruce)
- Trees: Broadleaf (e.g., birch)
- There is no vegetation

In the wider area as far as you can see.

15. Disturbance

- No disturbance
- Natural disturbance (example: burned area, slumps)
- Disturbance by humans (example: pipeline, storage area, reservoir)
- Unknown

If there is disturbance, please describe it (for example, what type, how far from the site, how big):

16. Overview photos of the site

With your phone at highest resolution available (or another camera e.g., 50 mm lens, if no cell phone):

- One photo of the site (landscape or portrait) as close as possible, showing the location (e.g., snow pit, measurement spot, etc.)
- A set of four pictures from next to the site, one looking to the North, one to the East, one to the South, and one to the West.

These should be in landscape mode with about 10% of the photo above the horizon; in one of the shots it would be good to have a person standing about 20 m away (for scale) and looking away from the camera (to avoid privacy issues). Otherwise use a scale bar, shovel, or any other object with a distinct size to indicate the scale.

Section 1: Snow – snow depth [cm]

Where to measure: 10–30 m transect: This should be established before taking any measurements at the site. See Section 0 (Fig. A3), above, for details of how to select and mark the transect.

When to measure: Start preparing the transect before the first snowfall by marking its beginning and end (see Fig. A3). Take a photograph of the site with the transect. Start the snow depth measurements with the first day of snowfall (beginning of snow season) until the end of the melting season (less than half of the ground area covered by snow). To capture the change of snow depth over time, measure ideally once per week, or alternatively every second week. Monthly measurements are still valuable.

Instrument: stick and tape measure, or graded avalanche probe.

Time: 2–3 minutes per measurement.


Scale: 10–30 m transect, one measurement every metre.

Method: Snow depth measurements are made during the snow season with a pole and tape measure or, if available, with a graded avalanche probe. Put the pole through the snowpack at right angles to the snow surface until it reaches ground and record the snow surface depth (precision about 1 cm; Fig. A5).

Actions:

- Put the pole or the avalanche probe straight (90° to surface) through the snowpack.
- Make sure the pole reaches the ground (not just stuck on an ice layer or crust) – you may need to apply more pressure than you think or less pressure if on soft ground.
- If the ground cannot be reached, make a note including the point number (e.g., presence of a basal ice layer at point 4).
- Measure snow depth with the scale or tape measure, or read off from the scale on the probe.
- Record value [cm].
- Repeat every metre.
- Estimate how accurate the measurements are [cm].
- Take a photo of the transect.

Fig. A5. Snow depth measurement.



Section 2: Permafrost – thaw depth [cm]

Where to measure: Thaw depth is known to vary substantially over very short distances, and the site characteristics should be accounted for when deciding on the location and spacing of the points. The 10–30 m transect should be established before taking any measurements at the site. See Section 0, above, for details of how to select and mark the transect. Note that if your site is very rocky, you may find that thaw depth measurements are not possible.

When to measure: Ideally, thaw depth should be measured at regular intervals around once per month from the time of snowmelt until the annual freeze-up. However, if it can only be collected once, measuring during the end of summer/early fall when the thaw depth has reached its yearly maximum should be prioritized.


Instrument: Metal rod, measuring tape.

Time: 1 minute per measurement, depending on soil properties.

Scale: 10–30 m transect, one measurement every metre.

Method: At each measurement point, insert a frost probe vertically to the surface until the point of resistance (Fig. A6). Feel free to push against the resisting surface a couple of times to ensure that it is the frost table which you are hitting and not a rock suspended in the unfrozen soil. Measure the depth that the frost probe has gone into the soil. Make a note if the depth exceeds the length of the rod. If the observation is suspect (e.g., due to the probe hitting a stone) or inserting the rod is impossible altogether (gravel, bedrock, etc.), try to repeat the observation within a distance of less than 1 m, or leave the box blank.

Fig. A6. Thaw depth measurement: the frost probe is inserted into the ground; measure the length of the section of the probe that has gone into the ground.



Actions:

- Insert the frost probe into the soil vertically until resistance against frost table is met; gently press the back of your hand against the surface vegetation layer to determine the surface position.
- Record the depth that the frost probe has gone into the soil, noting if the measurement had to be made at another location due to an obstruction, or any other anomaly.
- Repeat every metre.
- Take a photo of the transect from either end.


Section 3: Vegetation – height [cm]

Where to measure

For all sites: take vegetation height measurements in 1 m × 1 m quadrats (squares) at each point along your 10–30 m transect. This transect should be established before taking any measurements at the site. See Section 0, above, for details of how to select and mark the transect.

If your site is a forest: measure a minimum of 10 individual trees in a 15 m × 15 m plot as well as the transect.

Fig. A7. Quadrat for measuring vegetation height. The four measurement locations are shown in red.



When to measure

Forest: at least once a year, preferably during growing season (June–August).

Tundra and other non-forested sites: at least once during peak growing season. Preferably a seasonal overview, i.e., once every month from spring (shoulder season) to autumn (shoulder season). Mark out quadrats and revisit the same locations for repeated measurements.

Instrument: camera and ruler (forest: preferably a tape measure, tundra: preferably a carpenter’s rule).

Time: Quadrats: 2 min per quadrat.

Trees: 1 min per tree.

Scale: At all sites: 10–30 1 m × 1 m quadrats at 1 m intervals along the transect (see Section 0 above).

Forest only: 15 m × 15 m area.

Method: Measurement of vegetation cover height.

Forest: measurement (for trees smaller than 2 m in height) or estimation of the tree height of 10 trees.

Actions:

1.1. All sites:


- Mark out 1 m × 1 m quadrats at each point along your transect.
- Mark two diagonals within the quadrant and measure vegetation (excluding moss/lichen) height at four locations along the diagonals as shown in [Figures A7 and A8](#).
- Measure vegetation height from the soil surface to the highest point of the vegetation, at each sample location without extending/pulling the plants. If there is no vegetation (other than moss/lichen) at the location, record 0. If there is moss or lichen at the point you measure then tick the “moss/lichen present” box.
- Take a photo of each quadrat.

1.2. Forest sites only:

Actions:

- Mark out/estimate a 15 m × 15 m plot that is representative of the bigger area.
- Take a picture from the middle of the plot in every direction, preferably with a tape measure or a person standing next to a tree for height reference.

Fig. A8. Measuring vegetation height along the transect.



- Estimate (trees taller than 2 m) and measure (trees smaller than 2 m) the height of 10 individual trees that are typical of the site. Select trees that cover the range of heights within the plot. If there are fewer than 10 trees in your plot, just measure every tree.

How to measure/estimate the tree height for trees taller than 2 m (simple estimation):

1. Especially in dense forest, it can be hard to go as far back as needed to use a geometric measurement method. A simple solution, which needs some practice but works well, is a height estimation using the help of objects close-by and your own height.
2. Step back and make sure you are able to see the tree top and the base.
3. Estimate the tree height based on 2 m increments using branches for guidance.

Section 4: Water – water level

Where to measure: 10–30 m transect: This should be established before taking any measurements at the site. See Section 0, above, for details of how to select and mark the transect.


When to measure: Try to capture the changes through the seasons, including the spring snowmelt season, (possible) drying over the summer, and any changes towards the fall/autumn. No observations are necessary when the ground and surface waters are completely frozen. *Ideally you could take these measurements at the same time as the permafrost thaw depth (see Section 2, above) because the same equipment is used. You can even use the same hole as for thaw depth, to reduce damage to the soil.*

Instrument: Camera (picture) and ruler, pointed metal rod (frost probe) or something to make holes (could also be your hands).

Time: 2 min per measurement.

Scale: 10–30 m transect, one measurement per metre.

Fig. A9. Measuring the water level below the surface and above the surface.



Method: At each point, note if the water level is found within the top 10 cm of the ground, below the top 10 cm of the ground, or above the ground surface.

Actions: If there is water above the surface, measure the depth of the water [m] with either the ruler or the frost probe.

- If there is no water above the surface, insert the frost probe 10 cm into the ground and note if the hole fills with water (i.e., the water level is within 10 cm of the ground surface) (Fig. A9). Tick the relevant box (“Water is less than 10 cm below ground – hole fills with water”; “Water level is deeper than 10 cm below ground – hole does not fill with water”). Repeat the procedure for all observation points.
- Take a photo of the transect from either end.

Section 5: Soil – soil pit


Where to measure: The soil pit should be located close to the other measurements but set to the side so that you do not dig up the ground close to where the other measurements are taken. The pit should be approximately 1 m wide.

When to measure: once only, at a time when the soil has thawed to its greatest depth (late summer, early autumn).

Instrument: Spade and (or) shovel, a ruler or tape measure, camera.

Time: 1 h, but may take one afternoon.

Fig. A10. Examining the soil pit during summer. This demonstrates what a soil pit might look like.




Scale: 1–2 m wide soil pit; ideally aim for 1 m depth, but you can stop once you hit the frozen layer.

Method: Determine the location and orientation of the soil pit. Decide which side of the pit you want to use to record the soil profile (Fig. A10). *To reduce melting in the pit, the side you choose should not be directly exposed to the sun.* Avoid walking on or otherwise disturbing the ground on this side of the pit. Dig the pit and carefully put the excavated materials aside so you can easily backfill the pit later. Describe the profile and take a picture of the soil face, including the ruler/tape measure for scale. Please refill the profile with soil and vegetation cover as well as you can afterwards.

Actions:

- Dig a soil pit.
 - Take a picture of the soil face including the ruler/tape measure for scale with the 0 level at the surface.
 - Describe the soil.
- Estimate the thickness of the upper organic layer — this is the depth to the boundary between organic (dark brown/peaty) and mineral soil.
 - Is there any ice (at the bottom of the profile)?
If yes, take a photo with a ruler included for scale.
 - Are there rocks in the soil?
If yes, take a photo with a ruler included for scale.
- Select the soil texture description which most closely describes your soil
- Clay soil
 - Sandy soil
 - Loamy soil
 - Silty soil
 - Peaty soil

Fig. A11. To determine the soil texture, take a handful of soil and follow the instructions in the flowchart (Fig. A11) adapted and reused with permission.



British Geological Survey© UKRI 2021

- Put the soil back into the pit.
- Cover with vegetation again.

For help with selecting soil texture, please follow the instructions in the flowchart (Fig. A11).